



Moving Headwater Streams to the Head of the Class



SCIENCE & MATHEMATICS

Search opportunities on the Oxford University Press **Journals Career Network**

Your perfect job is closer than you think

FREE

[View Metrics](#)

Email alerts

[New issue alert](#)

[New advance articles alert](#)

[Article activity alert](#)

W
i
n
s
o
r
H
.
L
o
w
e;
G
e
n
e
E
.
L
i
k
e
n
s

Receive exclusive offers and updates
from Oxford Academic

Related articles in

Web of Science

Citing articles via

Web of Science (119)

Google Scholar

CrossRef

Latest | Most Read | Most Cited

The Coming Era of Open Data

Long-Term Studies Contribute
Disproportionately to Ecology and Policy

Agriculture in 2050: Recalibrating Targets for
Sustainable Intensification

BioScience

(2005)

55

(3):

196-

197.

Nature Museums and Botanical Gardens for
Environmental Conservation in Developing
Countries

A Unified View of Development: Making Evo-
Devo Operational

DOI: [https://doi.org/10.1641/0006-3568\(2005\)055\[019](https://doi.org/10.1641/0006-3568(2005)055[019)

Published:

01 March
2005

**Issue
Section:**

N
e
w
s
&
F
e
a
t
u
r
e
s

**Everywhere
on
Earth,
streams
and
rivers
occur
in
hierarchical
networks
resembling
the**

branching
pattern
of
a
tree,
with
smaller
branches
joining
to
form
larger
branches
as
water
travels
from
uplands
to
lakes,
estuaries,
and
seas.
The
finest
branches
of
these
networks,
beginning
where

water
flowing
overland
first
coalesces
to
form
a
discernible
channel,
are
called
head-
water
streams.
Conservative
estimates
indicate
that
headwater
streams
account
for
more
than
70
percent
of
stream-
channel
length

in
the
United
States
(
L
e
o
p
o
l
d
e
t
a
l
.
1
9
6
4),
yet
because
of
their
small
size,
these
streams
are
often

missing
from
maps
that

 PDF

 Cite

 Share ▼

 Tools ▼




Volume 55, Issue 3

March 2005

Article Contents

Conservation challenges and opportunities

Research priorities

Protect the source

References cited

Author notes

[< Previous](#) [Next >](#)

management
of
natural
resources.

Relative
to
larger
streams
and
rivers
that

are
fed
by
upstream
networks

and
affected
by
cumulative
upstream
stressors,
the
small
drainage

areas
of
head-
water
streams
give
these
systems
high
levels
of
hydrologic
independence
and
ecological
autonomy.
This
independence
justifies
the
use
of
headwater
watersheds
as
building
blocks
in
the
construction
of

protected-
area
networks

(
S
a
u
n
d
e
r
s
e
t
a
l
.
2
0
0
2)

and
their
prioritization
in
management
and
regulatory
efforts
to
protect

many
of
the
ecosystem
services
we
value,
such
as
clean
water,
recreational
opportunities,
nutrient
removal,
and
biodiversity.

There
is
growing
evidence
that
the
water
quality,
biodiversity,
and
ecological
health
of
freshwater

systems
depend
on
functions
provided
by
headwater
streams,
which
are
similar
in
their
importance
to
the
fine
branches
of
the
human
respiratory
system
in
the
lung.
Among
the
functions
of
these

streams
are
the
maintenance
of
natural
discharge
regimes,
the
regulation
of
sediment
export,
the
retention
of
nutrients,
the
processing
of
terrestrial
organic
matter,
and
the
establishment
of
the
chemical
signature
for

water
quality
in
the
landscape.
High
levels
of
habitat
diversity
among
and
within
these
small
streams
create
niches
for
diverse
organisms,
including
headwater-
specialist
species
of
aquatic
invertebrates,
amphibians,
and
fish.

Headwaters

also

act

as

refugia

for

riverine

species

during

specific

life-

history

stages

and

critical

periods

of

the

year,

such

as

warm

summer

months.

Like

the

alveoli

(the

final

branches

of

the
respiratory
tree
that
serve
as
the
primary
gas
exchange
units
of
the
lungs),
headwater
streams
are
characterized
by
strong
and
vital
interactions
with
the
systems
that
surround
them.
Terrestrial
inputs

—
dissolved
nutrients,
toxins,
and
particulate
matter,
for
example
—
play
a
central
role
in
determining
the
physical
and
chemical
conditions
of
headwater
streams
(
L
i
k
e
n
s

a
n
d
B
o
r
m
a
n
n
1
9
7
4)
and
in
regulating
the
composition
and
productivity
of
biotic
communities
in
these
streams
(
W
a
l

l

a

c

e

e

t

a

l

.

1

9

9

7).

Because

of

this

close

terrestrial–

aquatic

linkage,

the

ecosystem

services

provided

by

head–

waters

and

the

species

they

support
tend
to
be
very
sensitive
to
natural
and
anthropogenic
disturbance
of
surrounding
lands.
Along
with
other
distinctive
qualities,
this
close
connection
creates
a
unique
set
of
challenges
and
opportunities
related

to
the
protection
of
head-
waters,
and
to
research
in
these
systems.

Conservation challenges and opportunities

It
could
be
argued
that
lowland
sites,
where
the
human
footprint

is
both
intense
and
expanding
quickly,
are
in
greater
need
of
formal
protection
than
upland,
headwater
areas.
There
is
no
doubt
that
it
is
important
to
safeguard
lowland
sites,
but
it

is
difficult
to
see
how
any
conservation
action
with
a
goal
of
protecting
the
long-
term
ecological
integrity
and
ecosystem
services
of
natural
systems,
whether
aquatic
or
terrestrial,
can
succeed
without

a
foundation
of
intact
and
functional
headwaters.
This
point
highlights
the
error
of
government
proposals
to
withdraw
the
protection
afforded
under
the
Clean
Water
Act
(33
U.S.C.,
chapter
26)
to
headwater

streams
and
other
“isolated”
waters.

The
high
sensitivity
of
ecological
processes
and
natural
communities
in
headwater
streams
to
atmospheric
and
terrestrial
disturbances
leads
to
low
thresholds
of
impact.
Consequently,
disturbances
that

are
spread
across
multiple
headwater
watersheds
—
as
might
result
from
road
networks,
air
pollution,
and
diffuse
patchworks
of
logging
sites
or
residential
development
—
are
likely
to
be
more
detrimental

than
disturbances
that
are
confined
to
few
or
to
individual
watersheds.
When
possible,
minimizing
the
spatial
extent
of
human
disturbance
in
headwater
areas
may
guard
against
the
widespread
degradation
of
physical

and
chemical
conditions
in
these
upland
stream
networks
and
the
subsequent
transmittal
of
impacts
there
to
downstream
systems.

Capitalizing
on
the
accessibility
and
natural
history
of
headwater
streams
to
generate
public

support
for
their
protection
is
another
conservation
strategy
with
high
potential
for
long-
term
benefits.
These
small
streams
run
through
many
backyards,
schoolyards,
and
public
parks.
They
can
be
home
to

net-
spinning
aquatic
insects
and
20-
centimeter-
long
salamanders,
and
can
serve
as
natural
mesocosms
for
observing
how
sediment
bars
and
dams
of
woody
debris
are
formed
and
function.
The
many

education
and
volunteer-
monitoring
initiatives
aimed
at
protecting
vernal
pools
show
that
this
combination
of
accessibility
and
compelling
natural
history,
when
in
the
hands
of
committed
and
energetic
people,
can
be

an
invaluable
conservation
tool.

Research priorities

The
article
by

B
e
r
n
h
a
r
d
t
a
n
d
c
o
l
l
e
a
g

u
e
s
(
2
0
0
5
)
in
this
issue
of
BioScience
spotlights
a
question
with
important
implications
for
the
conservation
of
headwater
streams:
To
what
extent
do
these

streams
act
to
modify
nutrients
exported
from
the
surrounding
watershed,
as
opposed
to
simply
being
passive
conduits
of
these
nutrients?
Although
more
work
on
this
topic
is
needed,
there
is
growing

evidence
that
in-
stream
processes
do
play
a
significant
role
in
modifying
the
nitrogen
input-
output
balance
of
headwater
watersheds.
These
findings
suggest
that
interpretations
of
nutrient
levels
in
head-
water

streams
must
account
for
both
terrestrial
and
in-
stream
processes,
which
may
act
independently
or
interactively
to
affect
watershed
export
values.
They
also
highlight
the
potential
for
recovery
times
of
both

terrestrial
and
in-
stream
processes
to
limit
the
resilience
of
head-
water
ecosystems
to
anthropogenic
disturbance.

There
is
general
understanding
of
the
role
of
headwaters
in
setting
the
chemical
signature
of

fresh
water
at
the
landscape
scale.
As
the
human
footprint
continues
to
expand
and
competition
among
conservation
priorities
strengthens,
spatially
explicit,
quantitative
understanding
of
how
cumulative
head-
water
impacts
affect
downstream

resources
is
likely
to
become
critical.
Especially
important
in
this
context
may
be
mechanistic
studies
of
how
headwater
watersheds
that
have
been
degraded
interact
with
undegraded
ones
to
affect
downstream
resources,

and
research
identifying
specific
thresholds
in
the
intensity
and
spatial
extent
of
headwater
impacts
beyond
which
degradation
of
downstream
resources
is
likely
to
occur.

We
believe
that
a
third
research
priority

should
be
on
investigations
of
the
spatial
population
dynamics
of
species
within
the
stream
networks
and
associated
matrices
of
small
watersheds
that
make
up
headwater
systems.
The
design
of
ecological
reserves

for
biodiversity
protection
is
largely
dependent
on
understanding
the
population
structure
and
dispersal
patterns
of
resident
species.
Knowledge
of
the
spatial
structure
of
populations
informs
estimates
of
the
minimum
area
required

to
prevent
local
extinction.
Maintaining
interpopulation
dispersal
enhances
ecological
resilience
by
increasing
the
likelihood
of
recolonization
if
local
extinctions
occur.
Using
a
combination
of
direct
and
indirect
methods
(e.g.,
mark-
recapture

and
population
genetic
analyses),
this
work
will
provide
critical
information
on
the
minimum
area
and
configuration
of
protected
headwater
areas
required
for
species
persistence.

Protect the source

Headwaters

are
a
source
of
life.
They
are
critical
habitat
for
rare
and
endangered
freshwater
species,
and
guardians
of
many
downstream
resources
and
ecosystem
services
on
which
humans
rely
(
M
e

y
e
r
e
t
a
l

.

2

0

0

3).

In
the
past
year,
deforestation
in
headwater
drainages
exacerbated
flooding
and
landslides
in
Haiti,
the
Philippines,
and
Indonesia,
destroying

lives
and
property
in
lowland
communities.
Fortunately,
opportunities
for
research,
education,
management,
and
legislation
that
can
lead
to
effective,
long-
term
protection
of
headwater
ecosystems
worldwide
are
as
clear
as
the

risks
of
failing
to
protect
these
critical
ecosystems.

References cited

- 1 Bernhardt
ES. 2005.
Can't see
the forest
for the
stream? In-
stream
processing
and
terrestrial
nitrogen
exports.
BioScience .
55:
219–230.
[Cross-
Ref](#)

- 2 Leopold LB

Wolman MG
Miller JP. 1964.
Fluvial
Processes in
Geomorphology.
San Francisco:
W. H. Freeman.

- 3 Likens GE
Bormann
FH. 1974.
Linkages
between
terrestrial
and aquatic
ecosystems.
BioScience .
24:
447–456.

[Cross-
Ref](#)

- 4 Meyer JL.
2003.
Where
Rivers Are
Born: The
Scientific
Imperative
for
Defending
Small
Streams
and

Wetlands.
Washington
(DC):
American
Rivers,
Sierra Club.

- 5 Saunders DL
Meeuwig JJ
Vincent ACJ.
2002.
Freshwater
protected
areas:
Strategies for
conservation.
*Conservation
Biology* . 16:
30–41.

[CrossRef](#)

- 6 Wallace
JB Eggert
SL Meyer
JL
Webster
JR. 1997.
Multiple
trophic
levels of a
stream
linked to
terrestrial
litter

inputs.
Science .
277:
102–104.
[CrossRef](#)

Author notes

1
Winsor
H.
Lowe
(email:
lowe@ecosystems
.)

o
r
g)
is
a
postdoctoral
associate
at
the
Institute
of
Ecosystem
Studies,
Millbrook,
NY
12545.
He
will
begin
work
as
an
assistant
professor
in
aquatic
ecology
at
the
University
of
Montana
in
June

2005.

2

Gene

E.

Likens

(email:

li

k

e

n

s

g

@

e

c

o

s

t

u

d

i

e

s

.

o

r

g)

is

the

president

and

director

of

the
Institute
of
Ecosystem
Studies,
where
he
holds
the
G.
Evelyn
Hutchinson
Chair
in
Ecology.
He
is
a
cofounder
of
the
Hubbard
Brook
Ecosystem
Study.

©
2005
American
Institute
of
Biological
Sciences

[About BioScience](#)

[Purchase](#)

[Editorial Board](#)

[Recommend to your Library](#)

[Author Guidelines](#)

[Advertising and Corporate Services](#)

[Facebook](#)

[Journals Career Network](#)

[Twitter](#)

Online ISSN 1525-3244

Print ISSN 0006-3568

Copyright © 2017 American Institute of Biological Sciences

[About Us](#)

Connect

Resources

Explore

[Contact Us](#)

[Join Our Mailing List](#)

[Authors](#)

[Shop OUP Academic](#)

[Careers](#)

[OUPblog](#)

[Librarians](#)

[Oxford Dictionaries](#)

[Help](#)

[Twitter](#)

[Societies](#)

[Oxford Index](#)

[Access & Purchase](#)

[Facebook](#)

[Sponsors & Advertisers](#)

[Epigeum](#)

[Rights & Permissions](#)

[YouTube](#)

[Press & Media](#)

[OUP Worldwide](#)

[Open Access](#)

[Tumblr](#)

[Agents](#)

[University of Oxford](#)

Oxford University Press is a department of the University of Oxford. It furthers the University's objective of excellence in research, scholarship, and education by publishing worldwide

OXFORD
UNIVERSITY PRESS

Copyright © 2017 Oxford University Press

[Privacy Policy](#)

[Cookie Policy](#)

[Legal Notices](#)

[Site Map](#)

[Accessibility](#)

[Get Adobe Reader](#)

