January 9, 2020

Russell Bacon Forest Supervisor Thunder Basin National Grassland Supervisor's Office 2468 Jackson St Laramie, WY 82070

> *Re: Thunder Basin Plan Amendment Comment Letter (EIS No. 20190250), sent via:* <u>https://cara.ecosystem-management.org/Public/CommentInput?Project=55479</u>

Dear Mr. Bacon:

We are scientists with decades of experience studying prairie dogs (*Cynomys* spp.) and prairie dog associated species. In this letter, we provide an overview of relevant science that demonstrates the essential role black-tailed prairie dogs play as native keystone species and ecosystem engineers in the Great Plains, including Thunder Basin National Grassland. We are concerned the proposed "2020 Amendment" Draft Environmental Impact Statement does not fully take this into account in its analysis of prairie dog impacts to rangeland vegetation.

This letter is in response to notice 84 Federal Register 54899.

Sincerely,



The Portrayal of Black-tailed Prairie Dog Activity in the Draft Environmental Impact Statement for Thunder Basin National Grassland's 2020 Plan Amendment

Thunder Basin National Grassland's Draft Environmental Impact Statement (DEIS) for the "2020 Plan Amendment" does not fully incorporate the role of black-tailed prairie dog (*Cynomys ludovicianus*; hereafter simply "prairie dog") activity as a natural disturbance process essential to the grassland ecosystem of the Thunder Basin and the wildlife this ecosystem supports. Though the DEIS acknowledges that prairie dogs are keystone species, the analysis of the effects of prairie dogs on rangeland vegetation in the DEIS makes the assumption that prairie dog activity is not part of the reference condition for the ecosystem. This assumption is wrong, yet underlies the analysis and conclusions, as evident in the following excerpts from the DEIS.

• <u>DEIS</u>, page 70: For this analysis, the "representative value" for production of *each ecological site's reference plant community was compared to the representative value for production of the vegetative state expected from long-term prairie dog occupation*, commonly the "increased bare ground" state or plant community with short-stature vegetation based on target acres for prairie dog occupancy. The *reference state and increased bare ground state* were chosen for analysis purposes to estimate differences in forage availability among alternatives. (emphasis added; footnote omitted)

This statement indicates that the "bare ground state" is not considered part of the "reference state." Prairie dogs can create areas of sparse vegetation and bare ground. We emphasize that these conditions are within the natural range of variation of the grassland ecosystem. Prairie dog colonies can occur in a given area for thousands of years (White 1985); thus, the colony is within the natural range of grassland variation.

• <u>DEIS</u>, page 80: Plant communities change as prairie dog colonies become more established through time and as population density grows and forage needs increase. In areas recently colonized (less than 10 years), there may be little difference in species composition and production, whereas repeated heavy grazing on older colonies often results in lower overall plant productivity and change in species composition (Johnson and Collinge 2004). *Removal of prairie dogs following short-term prairie dog occupation may allow the community to shift back toward the reference plant community, as long as a disturbance threshold has not been crossed*. (emphasis added)

Additionally, tables 11, 12, 13, and 14, include the "Indicator or Measure": "Target acres occupied by prairie dogs that would transition away from the vegetation reference state."

As native species, by definition, prairie dogs and their activities must be treated as essential components of the natural range of variation for Thunder Basin's grassland ecosystem. Prairie dog disturbance does not lead to departure from reference conditions. The following scientific overview supports this assertion.

The Essential Role of Black-tailed Prairie Dogs as Ecosystem Engineers and Keystone Species

It is hard to overstate the importance of prairie dogs to the ecology of grasslands. The role of prairie dogs as a keystone species is now well-established scientifically (Kotliar et al. 1999; Kotliar 2000; Miller et al. 2000, 2007; Reading 2009; Slobodchikoff et al. 2009; Ceballos et al. 2010; Cully et al. 2010; Davidson et al. 2012; Martínez-Estévez et al. 2013; Lacher et al. 2019, Augustine et al. In press). See our attached Table 1. Prairie dogs probably qualify under multiple categories of keystone species—as prey and for their modification of habitat (Mills et al. 1993, Davidson et al. 2012). The grassland areas that prairie dogs inhabit should probably be considered ecosystems unto themselves (Clark et al. 1989).

Keystone species enrich ecosystem function uniquely and significantly through their activities, and their impact is larger than predicted relative to their biomass (Paine 1980; Terborgh 1988; Mills et al. 1993; Power et al. 1996; Kotliar et al. 1999; Miller et al. 1998/1999). Prairie dogs are functionally unique; they perform roles within their ecosystem not performed by other species or processes (Kotliar 2000). The scientific literature clearly supports the argument that prairie dogs fulfill all the requirement of keystone species (Coppock et al. 1983a, b; Detling and Whicker 1988; Whicker and Detling 1988a, b; 1993; Reading et al. 1989; Kotliar et al. 1997; 1999; Wuerthner 1997; American Society of Mammalogists 1998; Kotliar 2000, Miller et al. 2000; Slobodchikoff et al. 2009; Ceballos et al. 2010; Cully et al. 2010; Delibes-Mateos et al. 2011; Davidson et al. 2012; Martínez-Estévez et al. 2013; Lacher et al. 2019; Augustine et al. In press).

Prairie dog activities and the changes made by these activities create a unique ecological system known as the "prairie dog ecosystem" (Clark et al. 1989; Miller et al. 1996). Over 200 vertebrate species have been observed on prairie dog colonies (Table 1) (Koford 1958; Tyler 1968; Campbell and Clark 1981, Clark et al. 1982; O'Meilia et al. 1982; Agnew et al. 1986; Reading et al. 1989; Sharps and Uresk 1990; Mellink and Madrigal 1993; Hoogland 1995; Barko 1996; Manzano 1996; Ceballos and Pacheco 1997; Ceballos et al. 1999; Kotliar et al. 1999, Reading 2009; Davidson et al. 2012, Lacher et al. 2019). Some of these species depend on prairie dog colonies for their survival and many appear to benefit, at least seasonally or opportunistically from their existence (Table 1) (Reading et al. 1989; Hoogland 1995; Manzano 1996; Ceballos et al. 1999; Kotliar et al. 2010).

Prairie dogs and other animals inhabiting prairie dog colonies represent a rich prey patch for a large number of predators (Table 1) (Reading et al. 1989; Miller et al. 1996; Plumpton and Anderson 1997; Berry et al. 1998; Kotliar et al. 1999; Reading 2009). A rich diversity of predators including prairie rattlesnakes (*Crotalus viridis*), golden eagles (*Aquila chrysaetos*), great horned owls (*Bubo virginianus*), weasels (*Mustela frenata*), bobcats (*Lynx rufus*), coyotes (*Canis latrans*), and others prey on prairie dogs and small mammals that have a higher abundance on prairie dog colonies (Agnew et al. 1986). Some predators, especially endangered black-footed ferrets (*Mustela nigripes*), are completely dependent on prairie dogs (Clark 1989; Miller et al. 1996, Jachowski et al. 2011a, b, Eads et al. 2014). Other species, such as badgers (*Taxidea taxus*), swift foxes (*Vulpes velox*), and ferruginous hawks (*Buteo regalis*), benefit

substantially from the presence of prairie dogs as prey (Uresk and Sharps 1986; Sharps and Uresk 1990; Allison et al. 1995; Plumpton and Andersen 1997, 1998; Berry et al. 1998; Goodrich and Buskirk 1998).

The benefits of prairie dogs extend well beyond simply being food for predators (Table 1) (Reading et al. 1989; Ceballos et al. 1999; Kotliar et al. 1999). Prairie dogs also substantially alter their environment, although the extent and nature of those changes vary by region and prairie dog species (Martínez-Estévez et al. 2013; Baker et al. 2013). Since prairie dogs excavate more burrows than they regularly utilize¹, they create hibernacula, dens, and nests for many animals, such as black-footed ferrets, swift foxes (Vulpes velox), American badgers (Taxidea taxus), cottontails (Sylvilagus spp.), burrowing owls (Athene cunicularia), shrews, other rodents, and several species of reptiles and amphibians (Reading et al. 1989; Sharps and Uresk 1990; Plumpton and Lutz 1993; Fitzgerald et al. 1994; Desmond et al. 1995; Kretzer and Cully 2001; Shipley and Reading 2006; van Nimegen et al. 2008; Reading 2009). These species and more also use the burrows as refugia from predators or temperature extremes. As a result, researchers have found that desert cottontails (S. audonbonii) and northern grasshopper mice (Onychomys leucogaster) exist in higher numbers on prairie dog colonies than in surrounding grasslands (O'Meilia et al. 1982; Agnew et al. 1988; Dano 1952 in Stapp 1998, Cully et al. 2010). Similarly, studies in Mexico found higher rodent species richness, density, and diversity, and higher avian species richness on prairie dog colonies compared with surrounding grasslands in Chihuahua, Mexico (Manzano 1996; Ceballos and Pacheco 1997; Ceballos, et al. 1999; Ceballos et al. 2010). Most of the research to date has focused on birds and mammals. Considerably less research has examined reptile and amphibian associations with prairie dogs, but those findings were similar; namely, that some species appear to benefit from prairie dog activities, while others are harmed, but overall richness is greater across the landscape (Kretzer and Cully 2001; Shipley and Reading 2006; and Shipley et al. 2008). Similarly, little is known about prairie invertebrates (but see Bangert and Slobodchikoff 2004, 2006; Davidson and Lightfoot. 2007; and Kenney et al. 2016), yet the burrows and altered vegetation in a prairie dog colony should offer habitat advantages to some invertebrates as well. Indeed, having both colonized and uncolonized areas of grasslands most likely increases beta diversity in nearly all taxa, creating a more biodiverse landscape (Goguen 2012).

Prairie dogs also have a large effect on vegetation structure, productivity, nutrient cycling, and ecosystem processes (Coppock et al. 1983a,b; Detling and Whicker 1988; Shalaway and Slobodchikioff 1988; Whicker and Detling 1988a, b; 1993; Weltzin et al. 1997a; Stapp 1998, Ceballos et al. 2010; Baker et al. 2013; Connell et al. 2019; Gervin et al. 2019; Geaumont et al. 2019). The activities of prairie dogs, especially their grazing and clipping of tall vegetation, result in changes in plant composition (Bonham and Lerwick 1976; Coppock et al. 1983a, Detling and Whicker 1988; Whicker and Detling 1988a, b; 1993, Weltzin et al. 1997a; Detling 1998; Ceballos et al. 2010; Gervin et al. 2019). In general, the vegetation on prairie dog colonies is characterized by lower biomass and a greater preponderance of annual forbs and short grasses compared to tall grasses and shrubs, but is higher in nitrogen (crude protein) concentration than

¹Despite the common belief that there are several prairie dogs per burrow entrance, there are actually several burrow entrances per prairie dog (Biggins et al. 1993; Hoogland 1995).

vegetation from surrounding areas (Bonham and Lerwick 1976; Coppock et al. 1983a, Weltzin et al. 1997a; Detling 1998, Baker et al. 2013; Connell et al. 2019; Gervin et al. 2019). Several species benefit from these changes and preferably use prairie dog colonies (Table 1), especially due to the short stature of the plants on them, such as Mountain Plover (*Charadrius montanus*) and Horned Larks (*Eremophila alpestris*) (Dinsmore et al. 2005).

Prairie dogs negatively impact some plant and animal species (Table 1), reducing the prevalence and controlling the spread of taller grasses and several shrubs, such as mesquite (*Prosopis* spp.), sagebrush (*Artemisia* spp.), and longleaf jointfir (*Ephedra trifurca*) (Bonham and Lerwick 1976; Coppock et al. 1983; List 1997; Weltzin et al. 1997b; Ceballos et al. 2010; Baker et al. 2013; Gervin et al. 2019). Ironically, prairie dogs are poisoned for livestock interests, but these shrubs reduce grass available for cattle, and mesquite makes roundups more difficult (Miller 1991).

Prairie dogs provide a number of ecosystem services. In northern Mexico, Martínez-Estévez et al. (2013) found consistently higher ground water recharge, lower rates of soil erosion, increased soil carbon storage, improved soil water potential, and higher forage availability compared with surrounding areas. Prairie dog burrowing activities modify ecosystem processes such as water, mineral and nutrient cycling. Prairie dogs turn over approximately 225 kg of soil per burrow system, which translates to several tons of soil per hectare (Whicker and Detling 1993). By mixing in nutrient-rich urine and manure, prairie dog digging can change soil composition, chemistry, and microclimate, facilitate below-ground herbivory, increase porosity of soil to permit deeper penetration of precipitation, and increase the incorporation of organic materials into the soil (Ingham and Detling 1984; Whicker and Detling 1988 a, b; Munn 1993; Outwater 1996; Martínez-Estévez et al. 2013). As a result, prairie dog colonies support higher numbers of nematodes and higher levels of soil nitrogen (Ingham and Detling 1984, Detling 1998). All of these processes contribute to above ground plants with a higher nutritional content, greater digestibility, and a larger live plant to dead plant ratio, creating favorable feeding habitat for other herbivores (Whicker and Detling 1993; Connell et al. 2019). Indeed, pronghorn (Antilocapra americana) and bison (Bison bison) preferentially graze on prairie dog colonies (Coppock et al. 1983; Krueger 1986; Detling and Whicker 1993, Detling 1998). Some evidence also suggests that cattle use prairie dog colonies for grazing (on the edges) and resting (centers) (Guenther and Detling 2003; Sierra-Corona et al. 2015). Foraging models predict that bison can gain weight faster by grazing on pastures with prairie dog colonies than on grasslands without prairie dogs (Vanderhyde 1985 in Whicker and Detling 1993).

Finally, burrowing owls, and likely other species, benefit from the vigilance of prairie dogs, who issue alarm calls when they spot potential predators (Bryan and Wunder 2014).

Kotliar et al. (1999:177) concluded that collectively these functions are large, not wholly duplicated by other species (either in form or extent), and that the loss of prairie dogs would lead to "substantial erosion of biological diversity and landscape heterogeneity across the prairie." They concluded that the prairie dog therefore fulfills the definition of keystone species (see also Kotliar 2000). We agree. The structure, form, and function of prairie dog colonies provide a keystone role in the prairie, and the role is large. Despite the difficulty in quantifying a role, we

contend that existing evidence indicates prairie dogs (and other associated species) provide important prey to predators, and their grazing and burrowing activities modify the environment in a manner beneficially used by numerous other prairie organisms (Whicker and Detling 1993; Kotliar et al. 1999, Reading 2009; Cully et al. 2010). Most importantly, those grazing and burrowing activities affect vegetative composition, vegetation quantity and quality, productivity, nutrient cycling, and soil quality (Bonham and Lerwick 1976; Coppock et al. 1983; Detling and Whicker 1988; Whicker and Detling 1988 a, b; 1993; Connell et al. 2019). Prairie dog declines and reintroduction attempts into certain regions, for example southern New Mexico (Facka et al. 2010, Davidson et al. 2014, 2018), are problematic owing to arid conditions exacerbated by drought (i.e., climate change). Consequently, maintaining existing prairie dog colonies in their native range is crucial to maintaining prairie dogs, the ecosystems they create, and the species that depend upon them.

Literature Cited

- Agnew, W., D. W. Uresk, and R. M. Hansen. 1986. Flora and fauna associated with prairie dog colonies and adjacent ungrazed mixed-grass prairie in western South Dakota. Journal of Range Management. 39: 135-139.
- Agnew, W., D. W. Uresk, and R. M. Hansen. 1988. Arthropod consumption by small mammals on prairie dog colonies and adjacent ungrazed mixed grass prairie in western South Dakota. USDA Forest Service General Technical Report RM-154: 8-87.
- Allison, P. S., A. W. Leary, and M. J. Bechard. 1995. Observations of wintering ferruginous hawks (Buteo regalis) feeding on prairie dogs (Cynomys ludovicianus) in the Texas panhandle. Texas Journal of Science. 47 (3): 235-237.
- American Society of Mammalogists. 1998. Resolution on the decline of prairie dogs and the grassland ecosystem in North America. 6-10 June, Blacksburg, Virginia.
- Anderson, E., S. C. Forrest, T. W. Clark, and L. Richardson. 1987. Paleobiology, biogeography, and systematics of the black-footed ferret, Mustela nigripes (Audubon and Bachman), 1851. Great Basin Naturalist Memoirs. 8: 11-62.
- Arrowood, P. C., C. A. Finley, and B. C. Thompson. 2001. Analyses of burrowing owl populations in New Mexico. J. Raptor Res. 35: 362-370.
- Augustine, D. J. 2011. Habitat selection by Mountain Plovers in shortgrass steppe. J. Wildl. Manage. 75(2): 297-304.
- Augustine, D. J. and B. W. Baker. 2013. Associations of grassland bird communities with blacktailed prairie dogs in the North American Great Plains. Conserv. Biol. 27(2): 324-334.
- Augustine, D. J. and J. D. Derner. 2012. Disturbance regimes and mountain plover habitat in shortgrass steppe: Large herbivore grazing does not substitute for prairie dog grazing or fire. J. Wildl. Manage. 76(4): 721–728;
- Augustine, D., A. Davidson, K. Dickinson, and B. Van Pelt. In press. Thinking like a grassland: Challenges and opportunities for biodiversity conservation in the Great Plains of North America. Rangeland Ecol. Manage.
- Bak, J. M., K. G. Boykin, B. C. Thompson, and D. L. Daniel. 2001. Distribution of wintering ferruginous hawks (*Buteo regalis*) in relation to black-tailed prairie dog (*Cyomys*

ludovicianus) colonies in southern New Mexico and northern Chihuahua. J. Raptor Res. 35: 142-129.

- Baker, B. W., D. J. Augustine, J. A. Sedgwick and B. C. Lubow. 2013. Ecosystem engineering varies spatially: a test of the vegetation modification paradigm for prairie dogs. Ecography 36: 230-239
- Bangert, R. and C. N. Slobodchikoff. 2004. Prairie dog engineering indirectly affects beetle movement behavior. Journal of Arid Environments 56:83-94.
- Bangert, R. and C. N. Slobodchikoff. 2006. Conservation of prairie dog ecosystem engineering may support beta and gamma diversity. Journal of Arid Environments 67: 100-115.
- Barko, V. A. 1996. Effect of the black-tailed prairie dog on avifaunal composition in southern shortgrass prairie. MS thesis. Oklahoma State University, Stillwater, OK.
- Bonham, C. D., and A. Lerwick. 1976. Vegetation changes induced by prairie dogs on shortgrass range. Journal of Range Management. 29: 221-225.
- Bryan, R. D. and M. B. Wunder. 2014. Western Burrowing Owls (*Athene cunicularia hypugaea*) eavesdrop on alarm calls of black-tailed prairie dogs (*Cynomys ludovicianus*). Ethology 120: 180-188.
- Campbell, T. M. and T. W. Clark. 1981. Colony characteristics and vertebrate associates of white-tailed and black-tailed prairie dogs in Wyoming. American Midland Naturalist. 105: 269-76.
- Cartron, J.-L. E., P. J. Polechla, Jr., and R. R. Cook. 2004. Prey of nesting ferruginous hawks in New Mexico. Southwest. Nat. 49 (2): 270-276.
- Ceballos, G. and J. Pacheco. 1997. The relationship of prairie dogs and regional biodiversity in Mexico. P. 52, in Seventh International Theriological Congress, Acapulco, Mexico (Abs.).
- Ceballos, G., J. Pacheco, and R. List. 1999. Influence of prairie dogs (*Cynomys ludovicianus*) on habitat heterogeneity and mammalian diversity in Mexico. Journal of Arid Environments. 41:161-172.
- Ceballos G., A. Davidson, R. List, J. Pacheco, P. Manzano-Fischer, G. Santos-Barrera, J. Cruzado1. 2010. Rapid decline of a grassland system and its ecological and conservation implications. PLoS ONE 5(1): e8562. doi:10.1371/journal.pone.0008562.
- Chipault, J. G. and J. K. Detling. 2013. Bison selection of prairie dog colonies on shortgrass steppe. W. N. Amer. Nat. 73: 168-176.
- Clark, T. W., D. Hinckley, and T. Rich (eds.). 1989. The prairie dog ecosystem: Managing for biological diversity. Montana BLM Wildlife Technical Bulletin No. 2.
- Clark, T. W., T. M. Campbell, D. G. Socha, and D. E. Casey. 1982. Prairie dog colony attributes and associated vertebrate species. Great Basin Naturalist. 42: 572-82.
- Collins, A. R., Workman, J. P., and D. W. Uresk. 1984. An Economic analysis of black-tailed prairie dog [*Cynomys ludovicianus*] Control. Journal of Range Management. 37(4): 358-61.
- Connell, L. C., L. M. Porensky, and J. D. Scasta. 2019. Prairie dog (*Cynomys ludovicianus*) influence on forage quantity and quality in a grazed grassland-shrubland ecotone. Rangeland Ecol. Manage. 72: 360–373.
- Cook, R. R., J. E. Carton, and P.J. Polechla, Jr. 2003. The importance of prairie dogs to nesting ferruginous hawks in grassland ecosystems. Wildl. Soc. Bull. 31: 1073-1082.
- Coppock, D. L., J. K. Detling, J. E. Ellis, and M. I. Dyer. 1983a. Plant-herbivore interactions in a North American mixed-grass prairie. I. Effects of black-tailed prairie dogs on intraseasonal

aboveground plant biomass and nutrient dynamics and plant species diversity. Oecologia. 56: 1-9.

- Coppock, D. L., J. K. Detling, J. E. Ellis, and M. I. Dyer. 1983b. Plant-herbivore interactions in a north american mixed-grass prairie: II. responses of bison to modification of vegetation by prairie dogs. Oecologia. 56: 10-15.
- Cully, Jack F. Jr. 1986. Gunnison's prairie dog: an important prey species for ferruginous hawks, red-tailed hawks, golden and bald eagles. Unpublished report, dated July 14, 1986. 11 pp. + tables, maps.
- Cully, Jr, J. F., S. K. Collinge, R. E. Vannimwegen, C. Ray, W. C. Johnson, B. Thiagarajan, D. B. Conlin and B. Holmes, 2010. Spatial variation in keystone effects: small mammal diversity associated with black-tailed prairie dog colonies. Ecography 33: 1-11.
- Dano, L. 1952. Cottontail rabbit (Sylvilagus audubonii baileyi) populations in relation to prairie dog (Cynomys ludovicianus ludovicianus) towns. M. S. thesis, Colorado State University, Fort Collins, CO.
- Davidson, A. D. and D. C. Lightfoot. 2007. Interactive effects of keystone rodents on the structure of desert grassland arthropod communities. Ecography 30: 515-525.
- Davidson, A. D., R. R. Parmenter, and J. R. Gosz. 1999. Responses of small mammals and vegetation to a reintroduction of Gunnison's prairie dogs. J. Mammal. 80: 1311-1324.
- Davidson, A. D., J. K. Detling, J. H. and Brown. 2012. Ecological roles and conservation challenges of social, burrowing, herbivorous mammals in the world's grasslands. Frontiers in Ecology and the Environment 10: 477e486.
- Davidson, A. D., M. T. Friggens, K. T. Shoemaker, C. L. Hayes, J. Erz, and R. Duran. 2014. Population dynamics of reintroduced Gunnison's Prairie dogs in the southern portion of their range. J. Wildl. Manag. 78(3): 429-439.
- Davidson, A. D., E. A. Hunter, J. Eerz, D. C. Lightfoot, A. M. McCarthy, J. K. Mueller, and K. T. Shoemaker. 2018. Reintroducing a keystone burrowing rodent to restore an arid North American grassland: challenges and successes. Restoration Ecol. 26(5): 909-920.
- Davis, J. R. and T. C. Theimer. 2003. Increased lesser earless lizard (*Holbrookia maculata*) abundance on Gunnison's prairie dog colonies and short term responses to artificial prairie dog burrows. Am. Midl. Nat. 150: 282-290.
- Delibes-Mateos, M., A. T. Smith, C. N. Slobodchikoff and J. E. Swenson. 2011. The paradox of keystone species persecuted as pests: A call for the conservation of abundant small mammals in their native range. Biol. Conserv. 144: 1335-1346.
- Desmond, M. 2004. Effects of grazing practices and fossorial rodents on a winter avian community in Chihuahua, Mexico. Biol. Conserv. 116: 235-242.
- Desmond, M. J., J. A. Savidge, and K. M. Eskridge. 2000. Correlations between burrowing owl and black-tailed prairie dog declines: A 7-year analysis. J. Wildl. Manage. 64: 1067-1075.
- Desmond, M. J., J. A. Savidge, and T. F. Seibert. 1995. Spatial patterns of burrowing owls (*Speotyto cunicularia*) nests within black-tailed prairie dog (*Cynomys ludovicianus*) towns. Canadian Journal of Zoology. 73: 1375-1379.
- Detling, J. K. 1998. Mammalian herbivores: Ecosystem-level effects in two grassland national parks. Wildlife Society Bulleting. 26: 438-448.

- Detling, J. K. and A. D. Whicker. 1988. A control of ecosystem processes by prairie dogs and other grassland herbivores. Pp. 23-29, in Proceedings of the eighth Great Plains wildlife damage control workshop, Rapid City SD, U. S. Forest Service.
- Dinsmore, S. J., G. C. White, and F. L. Knopf. 2005. Mountain Plover population responses to black-tailed prairie dogs in Montana. Journal of Wildlife Management 69:1546-1553.
- Duchardt, C. J., L. M. Prensky, D. J. Augustine, and J. L. Beck. 2018. Disturbance shapes avian communities on a grassland–sagebrush ecotone. Ecosphere 9(10): e02483. 10.1002/ecs2.2483.
- Duchardt, C. J., D. J. Augustine, and J. L. Beck. 2019. Threshold responses of grassland and sagebrush birds to patterns of disturbance created by an ecosystem engineer. Landscape Ecol. 34: 895–909.
- Eads, D. A., D. E. Biggins, T. M. Livieri and J. J. Millspaugh. 2014. Space use, resource selection and territoriality of black-footed ferrets: implications for reserve design. Wild. Biol. 20: 27-36.
- Ellison Manning, A. E. and C. M. White. 2001. Nest site selection by mountain plovers (*Charadrius montanus*) in a shrub-steppe habitat. W. N. Amer. Nat. 61: 229-235.
- Facka, A. N., G. W. Roemer, V. L. Mathis, M. Kam, and E. Geffen. 2010. Drought leads to collapse of black-tailed prairie dog populations reintroduced to the Chihuahuan Desert. J. Wildl. Manage. 74(8): 1752-1762.
- Geaumont, B. A, T. J. Hovick, R. F. Limb, W. M. Mack, A. R. Lipinski, and K. K. Sedivec. 2019. Plant and bird community dynamics in mixed-grass prairie grazed by native and domestic herbivores. Rangeland Ecol. Manage. 72(2): 374-384.
- Gervin, C. A., H. H. Bruun, T. Seipel, and N. D. Burgess. 2019. Presence of both active and inactive colonies of prairie dogs contributes to higher vegetation heterogeneity at the landscape scale. Am. Midl. Nat. 181: 183–194.
- Goguen, C. B. 2012. Comparison of bird and mammal communities on black-tailed prairie dog (*Cynomys ludovicianus*) colonies and uncolonized shortgrass prairie in New Mexico. J. Arid Ecol. 80: 27-34.
- Goodrich, J. M. and S. W. Buskirk. 1998. Spacing and ecology of North American badgers (*Taxidae taxus*) in a prairie-dog (*Cynomys leucurus*) complex. Journal of Mammalogy. 79: 171-179.
- Goodrich, J. M., and S. W. Buskirk. 1995. Control of abundant native for conservation of endangered species. Conservation Biology. 9(6): 1357-1364.
- Guenther, D. A. and J. K. Detling. 2003. Observations of cattle use of prairie dog towns. J. Range Manage. 56: 41-417.
- Hansen, R. M. and I. K. Gold. 1977. Black-tailed prairie dogs, desert cottontails and cattle trophic relations on shortgrass range. Journal of Range Management. 30: 210-14.
- Holycross, A. T. and J. D. Fawcett. 2002. Observations on neonatal aggregations and associated behaviors in the prairie rattlesnake, *Crotalus viridis viridis*. Am. Midl. Nat. 148: 181-184.
- Hoogland, J. L. 1995. The black-tailed prairie dog: social life of a burrowing mammal. University of Chicago Press, Chicago, IL.
- Ingham, R. E. and J. K. Detling. 1984. Plant-herbivore interactions in a North American mixedgrass prairie III. Soil nematode populations and root biomass on *Cynomys ludovicianus* colonies and adjacent uncolonized areas. Oecologia. 63: 307-313.

- Jachowski, D. S., R. A. Gitzen, M. B. Grenier, B. Holmes, and J. J. Millspaugh. 2011a. The importance of thinking big: Large-scale prey conservation drives black-footed ferret reintroduction success. Biol. Conserv. 144: 1560–1566.
- Jachowski, D. S., J. J. Millspaugh, D. E. Biggins, T. M. Livieri, M. R. Matchett, and C. D. Rittenhouse. 2011b. Resource selection by black-footed ferrets in South Dakota and Montana. Nat. Areas J. 31: 218–225.
- Kenney, J. J., J. Detling, and R. P. Reading. 2016. Influence of black-tailed prairie dogs (*Cynomys ludovicianus*) on short-horned grasshoppers (Orthoptera: Acrididae) on the shortgrass steppe of Colorado. Journal of Arid Environments 127: 93-99.
- Klatt, L. E. and D. Hein. 1978. Vegetative differences among active and abandoned towns of black-tailed prairie dogs (*Cynomys Ludovicianus*). Journal of Range Management. 31: 315-17.
- Knowles, C. J. and P. R. Knowles. 1994. Additional records of mountain plovers using prairie dog towns in Montana. Prairie Naturalist. 16: 183-186.
- Knowles, Craig J. 1985. Observations on Prairie Dog Dispersal in Montana. Prairie Naturalist. 17(1):33-40.
- Knowles, Craig J. 1988. An evaluation of shooting and habitat alteration for control of blacktailed prairie dogs. Pp. 53-56 in D. W. Uresk, G. L. Schenbeck, and R. Cefkin, eds. Eighth Great Plains wildlife damage control workshop proceedings. 28-30 April 1987, Rapid City, South Dakota. USDA Forest Service General Technical Report RM-154. 231 pp.
- Knowles, Craig J. and Pamela R. Knowles. 1994. A Review of Black-tailed Prairie Dog Literature in Relation to Rangelands Administered by the Custer National Forest. USDA, Forest Service. 61 pp.
- Kolbe, J. J., B. E. Smith, and D. M. Browning. 2002. Burrow use by tiger salamanders (*Ambystoma tigrinum*) at a black-tailed prairie dog (*Cynomys ludovicianus*) town in southwestern South Dakota. Herpt. Review 33: 35-99.
- Kotliar, N. B. 2000. Application of the new keystone-species concept to prairie dogs: How well does it work? Conservation. Biology. 15: 1715-1721
- Kotliar, N. B., B. W. Baker, and A. D. Whicker. 1997. Are prairie dogs a keystone species. Pp. 183, in Seventh International Theriological Congress, Acapulco, Mexico (Abstract).
- Kotliar, N. B., B. W. Baker, A D. Whicker, and G. Plumb. 1999. A critical review of assumptions about the prairie dog as a keystone species. Environ. Manage. 24: 177-192.
- Kretzer, J. E. and J. F. Cully, Jr. 2001. Effects of black-tailed prairie dogs on reptiles and amphibians in Kansas shortgrass prairie. Southwest. Nat. 46: 171-177.
- Krueger, K. 1986. Feeding relationships among bison, pronghorn, and prairie dogs: an experimental analysis. Ecology. 67(3): 760-770.
- Lacher, T. E., Jr., A. D. Davidson, T. H. Fleming, E. P. Gomez-Ruiz, G. F. McCracken, N. Owen-Smith, C. A. Peres, and S. B. Vander Wall. 2019. The functional roles of mammals in ecosystems. J. Mammal. 100(3): 942–964.
- Lantz, S. J. and C. J. Conway. 2009. Factors affecting daily nest survival of burrowing owls within black-tailed prairie fog colonies. J. Wildl. Manage. 73(2): 232-241.
- List, R. 1997. Ecology of kit fox (*Vulpes macrotis*) and coyote (*Canis latrans*) and the conservation of the prairie dog ecosystem in northern Mexico. Ph. D. dissertation, University of Oxford, Oxford, UK.

- List, R. and D. W. MacDonald. 2003. Home range and habitat use of the kit fox (*Vulpes macrotis*) in a prairie dog (*Cynomys ludovicianus*) complex. J. Zool. (Lond.) 259: 1-5.
- Lomolino, M. V. and G. A. Smith. 2003a. Prairie dog towns as islands: Applications of island biogeography and landscape ecology for conserving nonvolant terrestrial vertebrates. Global Ecol. & Biogeo. 12: 275-286.
- Lomolino, M. V. and G. A. Smith. 2003b. Terrestrial vertebrate communities at black-tailed prairie dog (*Cynomys ludovicianus*) towns. Biol. Conserv. 115: 89-100.
- Magle, S. B., K. A. Salamack, K. R. Crooks, and R. P. Reading. 2012. Effects of habitat fragmentation and black-tailed prairie dogs on urban avian diversity. Biodiversity and Conservation 21: 2803-2821.

Manzano-Fischer, P. 1996. Avian communities associated with prairie dog towns in northwestern Mexico. M.S. Thesis, Oxford University, Oxford, U.K.

- Manzano, P. 1996. Avian communities associated with prairie dog towns in northwestern Mexico. M. S. thesis. University of Oxford, Oxford, UK.
- Manzano-Fischer, P., R. List, and G. Ceballos. 1999. Grassland birds in prairie dog towns in northwestern Chihuahua, Mexico. Studies in Avian. Biol. 19: 263-271.
- Marsh, R. E. 1984. Ground squirrels, prairie dogs, and marmots as pests on rangeland. Pp. 195-208, in Proceedings of the conference for organization and practice of vertebrate pest control, 30 August-3 September 1982, Fernherst, UK, ICI Plant Protection Division, Fernherst, UK.
- Martínez-Estévez L., P. Balvanera, J. Pacheco, and G. Ceballos. 2013 Prairie dog Decline reduces the supply of ecosystem services and leads to desertification of semiarid grasslands. PLoS ONE 8(10): e75229. doi:10.1371/journal.pone.0075229.
- Miller, B. and G. Ceballos. 1994. Managing conflict and biotic diversity in the prairie dog ecosystem. Endangered Species Update. 11(6 &7): 1-6.
- Miller, B. and R. P. Reading. 2012. Challenges to black-footed ferret recovery: protecting prairie dogs. W. North Amer. Nat. Vol. 72(2): 228-240.
- Miller, B., D. E. Biggins, and R. Crete. 1993. Workshop Summary. in Proceedings of the Symposium on the Management of Prairie Dog Complexes for the Reintroduction of the Black-footed Ferret. Eds. John L. Oldemeyer, Dean E. Biggins, and Brian J. Miller. Washington, DC: USFWS.
- Miller, B., G. Ceballos and R. P. Reading. 1994. The prairie dog and biotic diversity. Conservation Biology. 8: 677-681.
- Miller, B, R. P. Reading, and S. Forrest. 1996. Prairie night: Recovery of the black-footed ferret and other endangered species. Smithsonian Institution Press, Washington, DC.
- Miller, B. J., R. P. Reading, D. E. Biggins, J. K. Detling, S. C. Forrest, J. L. Hoogland, J. Javersak, S. D. Miller, J. Proctor, J. Truett, and D. W. Uresk. 2007. Prairie dogs: An ecological review and current biopolitics. J. Wildl. Manage. 71(8): 2801-2810
- Miller, B., R. Reading, J. Hoogland, T. Clark, G. Ceballos, R. List, S. Forrest, L. Hanebury, P. Manzano, J. Pacheco, and D. Uresk. 2000. The role of prairie dogs as keystone species: Response to Stapp. Conservation Biology. 14: 318-321.
- Miller, B., R. Reading, J. Strittholt, C. Carroll, R. Noss, M Soulé, O. Sánchez, J. Terborgh, D. Brightsmith, T. Cheeseman, and D. Foreman. 1998/99. Focal species in design of reserve networks. Wild Earth. 8(4): 81-92.

- Miller, B., C. Wemmer, D. Biggins, and R. Reading. 1990. A proposal to conserve black-footed ferrets and the prairie dog ecosystem. Environmental Management. 14: 763-769.
- Miller, J. A. 1991. Research updates. BioScience. 41: 750-753.
- Miller, S. 2002. Norton: Science to seal dogs' fate. Rapid City Journal, August 17, 2002.
- Mills, L. S., M. E. Soulé, and D. F. Doak. 1993. The history and current status of the keystone species concept. BioScience. 43: 219-224.
- Munn, L. C. 1993. Effects of prairie dogs on physical and chemical properties of soils. Pp. 11-17, in Management of prairie dog complexes for the reintroduction of the black-footed ferret. J. L. Oldemeyer, D. E. Biggins, and B. J. Miller (Eds.). U. S, Department of the Interior, Washington, D. C.
- O'Meilia, M. E., F. L. Knopf, and J. C. Lewis. 1982. Some consequences of competition between prairie dogs and beef cattle. Journal of Range Management. 35: 580-585.
- Outwater, A. 1996. Water: A natural history. Basic Books, New York, NY.
- Paine, R. T. 1980. Food webs: Linkage, interaction strength and community infrastructure. Journal of Animal Ecology. 49: 667-685.
- Plumpton, D. L. and D. E. Andersen. 1997. Habitat use and time budgeting by wintering ferruginous hawks. The Condor. 99: 888-893.
- Plumpton, D. L. and D. E. Andersen. 1998. Anthropogenic effects on winter behavior of ferruginous hawks. Journal of Wildlife Management. 62: 340-346.
- Plumpton, D. L. and R. S. Lutz. 1993. Nesting habitat use by burrowing owls in Colorado. Journal of Raptor Research. 27: 175-179.
- Power, M. E., D. Tilman, J. A. Estes, B. A. Menge, W. J. bond, L. S. Mills, G. Daily, J. C. Castilla, J. Lubchenco, and R. T. Paine. 1996. Challenges in the quest for keystones. BioScience. 466: 9-20.
- Price, E. R. 2002. Observation of rock wrens using white-tailed prairie dog burrows. Prairie Nat. 34: 149-150.
- Randall, D. 1976. Shoot the damn prairie dogs! Defenders 51:378-381. Reading, R. Denver Zoological Foundation and University of Denver, pers. comm. 2004
- Reading, R. P. 1993. Toward an endangered species reintroduction paradigm: A case study of the black-footed ferret. Ph. D. Dissertation. Yale University, New Haven, CT. 558 pp.
- Reading, R. P. 2009. Vertebrate species associated with prairie dogs. Pp. 139-146, in Prairie dogs: Communication and community in an animal society. C. Slobodchikoff, B. S. Perla, and J. L. Verdolin. Harvard University Press, Cambridge, MA
- Reading, R. P. and S. R. Kellert. 1993. Attitudes toward a proposed black-footed ferret (*Mustela nigripes*) reintroduction. Conservation Biology. 7: 569-580.
- Reading, R. P., B. J. Miller, and S. R. Kellert. 1999. Values and attitudes toward prairie dogs. Anthrozoös. 12: 43-52.
- Reading, R. P., S. R. Beissinger, J. J. Grensten, and T. W. Clark. 1989. Attributes of black-tailed prairie dog colonies in northcentral Montana, with management recommendation for conservation of biodiversity. Montana BLM Wildlife Technical Bulletin. 2: 13-27.
- Reading, R. P., T. W. Clark, A. Vargas, L. R. Hanebury, B. J. Miller, and D. Biggins. 1996. Recent directions in black-footed ferret (*Mustela nigripes*) recovery. Endangered Species Update 13(10-11): 1-6.

- Reading, R. P., T. W. Clark, A. Vargas, L. R. Hanebury, B. J. Miller, D. Biggins, and P. Marinari. 1997. Black-footed ferret (*Mustela nigripes*) conservation update. Small Carnivore Conservation. 17: 1-6
- Reading, R. P., T. W. Clark, L. A. McCain, and B. J. Miller. 2002. Black-tailed prairie dog conservation: A new approach for a 21st century challenge. Endangered Species Update. 19: 162-170.
- Reading, Richard P., Grensten, John J., Beissinger, Steven R., and Tim W. Clark. 1989.
 Attributes of Black-tailed Prairie Dog Colonies, Associated Species, and Management Implications. In The Prairie Dog Ecosystem: Managing for Biological Diversity. Eds. Tim W. Clark, Dan Hinckley, and Terrell Rich. Billings, MT: BLM.
- Restani, M., L. R. Rau, and D. L. Flath. 2001. Nesting ecology of burrowing owls occupying black-tailed prairie dog towns in southeastern Montana. J. Raptor Res. 35: 296-303.
- Shalaway, S. and C. N. Slobodchikoff. 1988. Seasonal change in the diet of prairie dogs. J. Mammal. 69: 835-841.
- Sharps, J. C. and D. W. Uresk. 1990. Ecological review of black-tailed prairie dogs and associated species in western South Dakota. Great Basin Naturalist. 50:339-345.
- Sheffield, S. R. and M. Howery. 2001. Current status, distribution, and conservation of the burrowing owl in Oklahoma. J. Raptor Res. 35: 351-356.
- Shipley, B. and R. P. Reading. 2006. Comparative abundance of herpetofauna and small mammals between prairie dog colonies and non-colonized grassland. J. Arid Ecol.. 66(1): 27-41.
- Shipley, B., R. P. Reading, and B. Miller. 2008. Comparing capture rates of reptiles and amphibians in black-tailed (*Cynomys ludovicianus*) prairie dog colonies and uncolonized prairie in Colorado, USA. Western North American Naturalist 62(2): 245-248.
- Sierra–Corona, R, A. Davidson, E. L. Fredrickson, H. Luna-Soria, H. Suzan-Azpiri, E. Ponce-Guevara, and G. Ceballos. 2015. Black-tailed prairie dogs, cattle, and the conservation of North America's arid grasslands. PLoS ONE 10(3): e0118602. doi:10.1371/journal.pone.0118602
- Sidle, J. G., M. Ball, T. Byer, J. J. Chynoweth, G. Foli, R. Hodorff, G. Moravek, R. Peterson, and D. J. Svingen. 2001. Occurrence of burrowing owls in black-tailed prairie dog colonies on Great Plains National Grasslands. J. Raptor Res. 35: 316-321.
- Slobodchikoff, C. N., B. Perla and J. L. Verdolin. 2009. Prairie Dogs: Communication and Community in an Animal Society. Harvard University Press, Cambridge, MA.
- Smith, G. A., and M. V. Lomolino. 2004. Black-tailed prairie dogs and the structure of avian communities on the shortgrass prairie. Oecologia 138: 592-602.
- Soulé, M. and R. Noss. 1998. Rewilding and biodiversity: complementary goals for conservation. Wild Earth. 8: 19-28.
- Stapp, P. 2007. Rodent communities in active and inactive colonies of black-tailed prairie dogs in shortgrass steppe. J. Mammal. 88(1): 241-249.
- Stapp, P. 1998. A reevaluation of the role of prairie dogs in the Great Plains grasslands. Conservation Biology. 12: 1253-1259.
- Terborgh, J. 1988. The big things that run the world -- a sequel to E. O. Wilson. Conservation Biology. 2: 402-403.

- Terborgh, J., J. Estes, P. Paquet, K. Ralls, D. Boyd, B. Miller, and R. Noss. 1999. Role of top carnivores in regulating terrestrial ecosystems. Pp. 39-64, in Continental conservation: Design and management principles for long-term regional conservation networks. M. Soulé, and J. Terborgh (Eds.). Island Press, Covelo, CA.
- Treviño-Villarreal, J. 2000. Mexican Prairie Dog. In Endangered Animals: A Reference Guide to Conflicting Issues. R. P. Reading and B. J. Miller (Eds.). Greenwood Press, Westport, CT.
- Treviño-Villarreal, J. and W. E. Grant. 1998. Geographic range of the endangered Mexican prairie dog (*Cynomys mexicanus*). Journal of Mammalogy. 79: 1273-1287.
- Truett, J. C. 2002. Aplomado falcons and grazing: Invoking history to plan restoration. Southwest. Nat. 47: 379-400.
- Truett, J. C., M. Phillips, K. Kunkel and R. Miller. 2001. Managing bison to restore biodiversity. Great Plains Research. 11:123-44. U. S. Bureau of Land Management. 2005. Final Programmatic Environmental Impact Statement on Wind Energy Development on BLM-Administered Lands in the Western United States. June 2005.
- Tyler, J. D. 1968. Distribution and vertebrate associations of the black-tailed prairie dog in Oklahoma. Ph.D. Thesis, University of Oklahoma, Norman, OK. 85 pp.
- Uresk, D. W. and J. C. Sharps. 1986. Denning habitat and diet of the swift fox in western South Dakota. Great Basin Naturalist. 46: 249-253.
- Uresk, D. W. 1985. Effects of controlling black-tailed prairie dogs on plant production. Journal of Range Management. 38(5): 466-8.
- van Nimwegen, R. E., J. Kretzer, and J. F. Cully Jr. 2008. Ecosystem engineering by a communal mammal: How prairie dogs structure rodent communities. Ecology 89(12): 3298-3305.
- Vanderhye, A. V. R. 1985. Interspecific nutritional facilitation: Do bison benefit from feeding on prairie dog towns? M. S. thesis. Colorado State University, Fort Collins, CO.
- VerCauteren, T. L. S. W. Gillihan, and S. W. Hutchings. 2001. Distribution of burrowing owls on public and private lands in Colorado. J. Raptor Res. 35 (4): 357-361.
- Wallace, M. G., H. J. Cortner, M. A. Moote, and S. Burke. 1996. Moving toward ecosystem management: examining a change in philosophy for resource management. Journal of Political Ecology. 3: 1-36.
- Weltzin, J. F., S. L. Dowhower, and R. K. Heitschmidt. 1997a. Prairie dog effects on plant community structure in southern mixed-grass prairie. Southwestern Naturalist. 42: 251-258.
- Weltzin, J. F., S. Archer, and R. K. Heitschmidt. 1997. Small-mammal regulation of vegetation structure in a temperate savanna. Ecology. 78(3): 751-763.
- Whicker, A. and J. K. Detling. 1993. Control of grassland ecosystem processes by prairie dogs. Pp. 18-27, in Management of prairie dog complexes for the reintroduction of the blackfooted ferret. J. L. Oldemeyer, D. E. Biggins, and B. J. Miller (Eds.). U. S, Department of the Interior, Washington, D. C.
- Whicker, A. D. and J. K. Detling. 1988a. Ecological consequences of prairie dog disturbances. BioScience. 38: 778-785.
- Whicker, A. D. and J. K. Detling. 1988b. Modification of vegetation structure and ecosystem processes by North American grassland mammals. Pages 301-316, in Plant form and vegetation structure: Adaptation, plasticity and relation to herbivory. J. J. A. Werger, P. J. M.

van der Aart, H. J. During, and J. T. A. Verhoeven (Eds.). SPB Academic Publishing, the Hague, the Netherlands.

- White, E. M. 1985. Antiquity, original size and location of prairie dog towns in Wind Cave National Park. Final report for National Park Service, contract CX-1200-4-A040.
- Winter, S. L., J. F. Cully, Jr., and J. S. Pontius. 2003. Breeding season avifauna of prairie dog colonies and non-colonized areas in shortgrass prairie. Trans. Kans. Acad. Sci. 106 (3/4): 129-138.
- Wuerthner, G. 1997. Viewpoint: The black-tailed prairie dog—headed for extinction? Journal of Range Management. 50: 459-466.

Sp	ecies	Benefit ^a	Citation ^b
Ob	bligate Associate ($N = 1$)		
	Black-footed ferret, Mustela nigripes	PD, BU	1, 8, 39, 41, 48, 49, 50, 51
Str	rong, Well Documented Benefits of Association (N = 1	7)	
	Tiger Salamander, Ambstoma tigrinum	BU, TP	2, 3, 13, 14, 25, 27, 28
	Lesser earless lizard, Holbrookia maculata	BU, PV	2, 4, 13, 31
	Prairie rattlesnake, Crotalus viridus	BU, PD, OP	13, 14, 27, 28, 33, 34, 41
	Killdeer, Charadrius vociferous	OV	2, 3, 6, 7, 8, 10, 28, 34, 35, 44, 52, 59, 61
	Mountain plover, Charadrius montanus	OV	2, 3, 7, 11, 26, 35, 44, 47, 53, 54, 57, 59, 60
	Ferruginus hawk, Buteo regalis	PD, OP	2, 3, 4, 6, 7, 8, 11, 21, 28, 30, 32, 34, 37, 41. 57, 59
	Golden eagle, Aquila chrysaetos	PD, OP	1, 2, 3, 7, 8, 34, 59, 61
	Burrowing owl, Athene cunicularia	AC, BU, OV	 72, 3, 4, 6, 7, 8, 11, 17, 18, 19, 20, 22, 28, 34, 35, 36, 37, 41, 44, 57, 58, 59
	Horned lark, Eremophila alpestris	OV	2, 3, 4, 6, 7, 8, 10, 11, 28, 34, 35, 37, 41, 44, 59, 60, 61
*	Curve-billed thrasher, Toxostoma curvirostre	PV	4, 10, 57
	American badger, Taxidea taxus	BU, PD, OP	1, 2, 3, 4, 7, 8, 12, 27, 28, 34, 41, 57
	Coyote, Canis latrans	BU, PD, OP	1, 2, 3, 4, 7, 8, 12, 27, 28, 34
	Desert cottontail, Sylvilagus audubonii	BU, PV	1, 2, 3, 4, 8, 9, 12, 14, 34, 38, 57
	Eastern cottontail. Svlvilagus floridanus	BU. PV	8, 27, 28, 34
	N. grasshopper mouse, Onychomys leucogaster	PV, OP	1, 3, 6, 8, 14, 27, 28, 34, 42, 43, 45, 55
	Swift fox, <i>Vulpes velox</i>	PD, OP	2, 8, 27, 28, 34
	Pronghorn, Antilocapra americana	PV	1, 3, 4, 7, 8, 27, 28, 34, 57

Table 1. Animal species cited as associated with prairie dogs and evidence of thisassociation. Updated from Kotliar et al. (1999).

Sp	pecies	Benefit ^a	Citation ^b
So	ome Documented Evidence of a Positive Association (N	N=24)	
*	Plans spadefoot toad, Spea bombifrons	BU, PV	41
*	Greater short-horned lizard, <i>Phrvnosoma hernandesi</i>	PV, OP	14, 42
*	Lesser earless lizard, Holbrookia maculate	PV	41
*	Plans blackhead snake, Tantilla nigriceps	PV, OP	41
*	Aplomado falcon, Falco femoralis	PD, OP, OV	16
	Swainson's hawk, Buteo swainsoni	PD, OP	2, 3, 4, 6, 7, 8, 9, 10, 11, 57
	Red-tailed hawk, Buteo jamaicensis	PD, OP	1, 2, 3, 6, 8, 9, 34, 52, 57, 59
*	Zone-tailed hawk, Buteo albonotatus	OP	41
*	Bald eagle, Haliaeetus leucocephalus	PD, OP	41, 61
	Great horned owl, Bubo virginianus	OP	3,6
	Snowy owl, Nyctea scandiaca	PD, OP	8
*	Short-eared owl, Asio flammeus	PV, OP	11
*	Cactus wren, Campylorhynchus brunneicapillus	PV	11
*	McCown's longspur, Calcarius mccownii	PV	41, 44
	White-tailed jackrabbit, Lepus townsendii	PV	1, 3, 7, 8
*	Northern pygmy mouse, Baiomys taylori	PV	41
*	Southern grasshopper mouse, Onychomys torridus	PV, OP	41
*	Mearn's grasshopper mouse, Onychomys arenicola	PV, OP	41
	Red fox, Vulpes vulpes	PD, OP	3, 7, 8
*	Kit fox, Vulpes macrotis	BU, OP	41
	Bobcat, Lynx rufus	PD, OP	1, 2, 8
	Elk, Cervus elaphus	PV	7, 57
	Mule deer, Odocoileus hemionus	PV	3, 7, 8
	Bison, Bison bison	PV	1, 2, 8, 57, 62, 63
Da	becomented Evidence of a Negative Association ($N = 23$	3)	
*	Plains spadefoot toad, Spea bombifrons	PV	14, 42
*	Bunchgrass lizard, Sceloporus scalaris	PV	41
*	Gopher snake, Pituophis catenifer	PV, OP	14, 42
*	Yellow-bellied racer, Coluber constrictor	BU, OP	13
	Northern bobwhite, Colinus virginianus	PV	2, 34
	Common nighthawk, Chordeiles minor	PV	2, 3, 7, 10, 34, 57, 59
	Sage thrasher, Oreoscoptes montanus	PV	7, 59, 60
	Grasshopper sparrow, Ammodramus savannarum	PV	6, 7, 8, 11, 34, 35, 37, 44, 59, 60, 61
	Cassin's sparrow, Aimophila cassinii	PV	10, 11, 34, 37, 57

Species		Benefit ^a	Citation ^b	
	Brewer's sparrow, Spizella breweri	PV	7, 11, 37, 44, 57, 59, 60	
	Baird's sparrow, Ammodramus bairdii	PV	7, 37, 44	
	Northern mockingbird, Mimus polyglottos	PV	2, 8, 10, 34, 57	
	Bullock's oriole. <i>Icterus bullockii</i>	PV	2, 34, 59	
*	Lark bunting, Calamospiza melanocorvs	PV	57, 59, 60	
*	Black-tailed jackrabbit, Lepus californicus	PV	41, 57	
*	Hispid cotton rat, Sigmodon hispidus	PV	55	
*	Merriam's kangaroo rat, Dipodomys merriami	PV	41, 43	
*	Hispid pocket mouse, Chaetodipus hispidus	PV	41, 43, 45, 55	
	Plains harvest mouse, Reithrodontomys montanus	PV	8, 55	
*	Silky pocket mouse, Pergnathus flavus	PV	41, 45, 55	
	Prairie vole, Microtus ochrogaster	PV	6, 8, 43, 55	
	Raccoon, Procyon lotor	PV, OP	2, 3, 8, 12	
*	Hooded skunk, Mephitis macroura	BU, OP	12	
Mi	nimal, Contradictory, or No Evidence of a Positive of	r Negative Asso	ociation ($N = 81$)	
*	Many-lined skink, Eumeces multivirgata	PV	14, 42, 44	
*	Prairie falcon, Falco mexicanus	PD, OP, OV	57	
	American kestrel, Falco sparverius	OP	2, 3, 4, 6, 7, 8, 9, 10, 34, 52, 57, 59	
*	Northern harrier, Circus hudsonius	OP	44	
*	Turkey vulture, Cathartes aura	PD, OP	57, 59, 61	
	Scaled quail, Callipepla squamata	PV	2, 10, 34, 57	
	Lesser prairie chicken, <i>Tympanuchus pallidicinctur</i>	OV	2	
	Sharp-tailed grouse, <i>Tympanuchus phasianellus</i>	OV	1, 6, 7, 8, 61	
	Sage grouse, Centrocerus urophasianus	OV	3, 7	
	Long-billed curlew, Numenius americanus	SV, TP, OV	2, 6, 7, 8, 10, 11, 34, 41, 57, 59	
	Upland sandpiper, Bartramia longicauda	PV	2, 6, 7, 8, 34, 44, 61	
*	Mourning dove, Zenaida macroura	PV	44, 52, 57, 59, 61	
	Northern flicker, Colaptes auratus	PV	2, 7, 8, 52, 59, 61	
*	Ash-throated flycatcher, Myiarchus cinerascens	PV	11	
	Scissor-tailed flycatcher, Tyrannus forficatus	PV	2, 28, 34	
	Say's phoebe, Sayornis saya	PV	6, 8, 52, 57, 59	
	Cassin's kingbird, Tyrannus vociferans	PV	10	
	Western kingbird, Tyrannus verticalis	PV	2, 3, 6, 7, 8, 10, 52, 57, 59, 61	
	Eastern kingbird, Tyrannus tyrannus	PV	2, 6, 7, 8, 10, 44, 59, 61	

Species		Benefit ^a	Citation ^b	
	Violet-green swallow, Tachycineta thalassina	PV	8. 57	
	N. rough-winged swallow, <i>Stelgidoptervx serripennis</i>	PV	6, 8, 57	
*	Tree swallow, <i>Tachycineta bicolor</i>	PV	11, 41, 52	
*	Bank swallow, <i>Riparia riparia</i>	PV	11, 61	
	Cliff swallow, Hirundo pyrrhonota	PV	2, 3, 7, 8, 10, 44, 57, 59,	
	Barn swallow, <i>Hirundo rustica</i>	PV	2, 3, 6, 7, 8, 10, 44, 52, 57, 61	
	American robin, Turdus migratorius	PV	7, 8, 52	
	Northern shrike, Lanius excubitor	OP	8	
	Water pipit, Anthus spinoletta	PV	8	
	Sprague's pipit, Anthus spragueii	PV	2, 44	
*	Brown thrasher, Toxostoma rufum	PV	61	
	Loggerhead shrike, Lanius ludovicianus	PV, OP	2, 3, 6, 7, 8, 10, 28, 34, 57, 59	
*	Rock wren, Salpinctes obsoletus	BU	24, 59	
*	Lucy's warbler, Vermivora luciae	PV	11	
*	Yellow-rumped warbler, Dendroica coronata	PV	11	
	Lapland longspur, Calcarius lapponicus	PV	2	
	Chestnut-collared longspur, Calcarius ornatus	OV	2, 3, 6, 7, 8, 28, 34, 37, 44	
	Rufous-sided towhee, Pipilo erythrophthalmus	PV	8	
	Savannah sparrow, Passerculus sandwichensis	PV	2, 3, 7	
*	Song sparrow, Melospiza melodia	PV	11, 52	
*	Black-throated sparrow, Amphispiza bilineata	PV	11	
	Dark-eyed junco, Junco hyemalis	PV	8, 52	
*	Field sparrow, Spizella pusilla	PV	61	
*	Clay-colored sparrow, Spizella pallida	PV	61	
	White-crowned sparrow, Zonotrichia leucophrys	PV	8, 52	
	Chipping sparrow, Spizella passerine	PV	8, 44, 59, 61	
	Vesper sparrow, Pooecetes gramineus	PV	2, 3, 7, 8, 10, 11, 44, 52, 57, 59, 61	
	Lark sparrow, Chondestes grammacus	PV	2, 4, 7, 8, 10, 11, 28, 34, 44, 57, 59, 61	
	Bobolink. dolichonyx oryzivorus	PV	8. 61	
	Dickcissel Spiza americana	PV	8, 61	
	Snow buntings, <i>Plectrophenax nimalis</i>	PV	3	
*	Western meadowlark, Sturnella neglecta	PV	41, 44, 52, 57, 59, 60, 61	

Species		Benefit ^a	Citation ^b
	Eastern meadowlark, Sturnella magna	PV	2, 28, 34, 41
	Brown-headed Cowbird, Molothrus ater	PV	2, 3, 7, 8, 28, 34, 44, 57, 59, 61
	Black-billed magpie, <i>Pica pica</i>	PV	3, 7, 8, 52, 59, 61
	Boat-tailed grackle, Quiscalus major	PV	2
*	Common grackle, Quiscalus quiscula	PV	44, 52, 59, 61
	Yellow-headed blackbird, Xanthocephalus xanthoceph	alus	PV 6, 7, 8, 11
	Red-winged blackbird, Agelaius phoeniceus	PV	2, 6, 7, 8, 10, 34, 44, 52, 59, 61
	Brewer's blackbird, Euphagus cyanocephalus	PV	2, 7, 8, 59, 61
	American crow, Corvus brachyrhynchos	OP	2, 6, 8, 10, 44, 52
	Chihuahuan raven, Corvus cryptoleucus	OP	2, 10, 57
	Common raven, Corvus corax	OP	8
	Pine siskin, Carduelis pinus	PV	8
	American goldfinch, Carduelis tristis	PV	8, 52, 61
*	House finch, Carpodacus mexicanus	PV	11, 52
*	Blue grosbeak, Guiraca caerulea	PV	11, 61
	Eastern mole, Scalopus aquaticus	PV	2
	Northern pocket gopher, Thomomys talpoides	PV	8
	Plains pocket gopher, Geomys bursarius	PV	2, 4, 8
*	North American deer mouse, Peromyscus maniculatus	PV	14, 43, 45, 46
*	Western harvest mouse, Reithrodontomys megalotis	PV	14, 45
	Ord's kangaroo rat, <i>Dipodomys ordi</i>	PV	1, 2, 4, 8, 9, 41, 43, 45
*	Banner-tailed kangaroo rat, Dipodomys spectabilis	PV	43
	Southern plains woodrat, Neotoma micropus	PV	2, 4, 43, 45
*	White-footed dormouse, Peromyscus leucopus	PV	43
*	Plains harvest mouse, Reithrodontomys montanus	PV	43
	Least chipmunk, Eutamias minimus	PV	3, 43
*	Thirteen-lined gr. squirrel, Ictidomys tridecemlineatus	PV	14, 43, 46, 55, 57
	Richardson's gr. squirrel, Spermophilus richardsonii	BU, PV	7
*	Spotted ground squirrel, Spermophilus spilosoma	PV	43
*	Ringtail, Bassariscus astutus	OP	12

No Data on Association, but Species Life History Suggests an Accidental Occurrence (N = 54)

*	Western chorus frog, Pseudacris triseriata	14, 42
	Northern leopard frog, Rana pipiens	3
	Western toad, Bufo boreas	3
	Bullfrog, Rana catesbeiana	8

Species		Benefit ^a	Citation ^b	
	Sagebrush lizard Scelonorus graciosus		3.4	
	Chihuahua spotted whiptail, <i>Cnemidophorus exsanguis</i>		4, 12	
	Smooth green snake, <i>Opheodrys vernalis</i>		8	
	Common garter snake, <i>Thamnophis sirtalis</i>		3	
*	Plains garter snake, <i>Thamnophis radix</i>		14, 42	
*	Western terrestrial garter snake. <i>Thamnophis elegans</i>		14, 42	
	Eared grebe, <i>Podiceps nigricollis</i>		7	
	Pied-billed grebe, <i>Podilymbus podiceps</i>		7	
	White pelican. <i>Pelecanus ervthrorhvnchos</i>		7	
	Double-crested cormorant, <i>Phalacrocorax auritus</i>		7	
	Black-crowned night heron, Nycticorax nycticorax		7	
	Great blue heron. Ardea herodias		7.8	
*	Osprey, Pandion haliaetus		11	
	Trumpeter swan, Cvngus buccinator		8	
	Snow goose, Chen caerulescens		2	
	Canada goose, Branta canadensis		7, 8, 52	
	Mallard, Anas platvrhvnchos		6, 7, 8, 52, 59	
	Gadwall, Anas strepera		7, 8	
	Ruddy duck, Oxyura jamaicensis		7	
	Green-winged teal, Anas crecca		3,7	
	American wigeon, Anas americana		7	
	Northern pintail, Anas acuta		6, 7, 8	
	Northern shoveler, Anas clypeata		8	
	Blue-winged teal, Anas discors		6, 7, 8	
	Canvasback, Aythya valisineria		8	
	Redhead, Aythya americana		7	
	Lesser scaup, Aythya affinis		7	
	Sora, Porzuna Carolina		6, 8	
	American coot, Fulicaa americana		7	
	Belted kingfisher, Ceryle alcyon		8	
	Red-headed woodpecker, Melanerpes erythrocephalus		8, 59, 61	
	Downy woodpecker, Picoides pubescens		8	
	Ladder-backed woodpecker, Picoides scalaris		2	
	Blue Jay, Cyanocitta cristata		8, 52	
	Eastern bluebird, Sialia sialis		8	
	Mountain bluebird, Sialia currocoides		7, 8	
	Ring-billed gull, Larus delawarensis		7, 8	
	Herring gull, Larus argentatus		7	

Species		Benefit ^a	Citation ^b
	California gull, Larus californicus		7
	Gray catbird, Dumetella carolinensis		8, 61
	Yellow warbler, Dendroica petechia		8, 61
	Common yellowthroat, Geothlypis trichas		8
*	Spotted towhee, Pipilo maculatus		52, 61
	Yellow-breasted chat, Icteria virens		8
	Western tanager, piranga ludoviciana		8
*	Black-capped chickadee, Poecile atricapillus		52
*	Lesser goldfinch, Carduelis psaltria		52, 57
	Common redpoll, Carduelis flammea		8
	Porcupine, Erethizon dorsatum		8
	Mink, Mustela vison		3, 8
De	omestic or Introduced Species ($N = 10$)		
	Gray partridge, Perdix perdix		7
	Ring-necked pheasant, Phasianus colchicus		8, 34, 61
	Rock dove, Columbia livia		2, 6, 8, 10, 52
	European starling, Sturnus vulgaris		2, 6, 7, 8, 52, 59
	House sparrow, Passer domesticus		2, 6, 8, 52
	Norway rat, Rattus norvegicus		8
	House mouse, Mus musculus		6, 8, 43
	Domestic horse, Equus caballus		3, 7
	Domestic cattle, Bos Taurus		3, 7, 34, 56
	Domestic sheep, Ovis aries		3, 7

* New species associate reported since Kotliar (1999).

^a Adapted from Kotliar et al. (1999): PD = prairie dogs as prey or carrion; OP = other vertebrate prey or carrion found on colonies; BU = burrows for nesting/shelter; OV = open vegetation or bare ground for nesting/foraging; PV = prairie vegetation for nesting/foraging; TP = temporary pools for breeding/forage; AC = alarm calls of prairie dogs.

^b Citations: 1 = Koford 1958; 2 = Tyler 1968; 3 = Campbell and Clark 1981; 4 = Clark et al. 1982; 5 = O'Meilia et al. 1982; 6 = Agnew et al. 1986; 7 = Reading et al. 1989; 8 = Sharps and Uresk 1990; 9 = Mellink and Madrigal 1993; 10 = Barko 1996; 11 = Manzano-Fischer 1996 and Manzano-Fischer et al. 1999; 12 = Ceballos et al. 1999; 13 = Kretzer and Cully 2001; 14 = Shipley and Reading 2006; 15 = List and MacDonald 2003; 16 = Truett, J. C. 2002; 17 = Restani et al. 2001; 18 = Sidle et al. 2001; 19 = Sheffield and Howery 2001; 20 = Arrowood et al. 2001, 21 = Bak et al. 2001; 22 = Desmond et al. 2000; 23 = Davidson et al. 1999 (Gunnison's prairie dogs); 24 = Price 2002 (white-tailed prairie dogs); 25 = Kolbe et al. 2002;

26 = Ellison Manning and White 2001, 27 = Lomolino and Smith 2003a; 28 = Lomolino and Smith 2003b, 30 = Cook et al. 2003, 31 = Davis and Theimer 2003, 32 = Cartron et al. 2004, 33 = Holycross and Fawcett 2002, 34 = Smith and Lomolino 2004, 35 = Winter et al. 2003, 36 = VerCauteren et al. 2001, 37 = Desmond 2004, 38 = Dano 1952, 39 = Miller et al. 1996, 40 = Bryan and Wunder 2014, 41 = Ceballos et al. 2010, 42 = Shipley et al. 2008, 43 = Cully et al. 2010, 44 = Augustine and Baker 2013, 45 = van Nimwegen et al. 2008, 46 = Stapp 2007, 47 = Dinsmore et al. 2005, 48 = Eads et al. 2014, 49 = Jachowski et al. 2011a; 50 = Jachowski et al. 2011b; 51 = Miller and Reading 2012; 52 = Magle et al. 2012; 53 = Augustine 2011; 54 = Augustine and Derner 2012, 55 = Pruett et al. 2010; 56 = Sierra-Corona et al. 2015; 57 = Goguen 2012; 58 = Lantz and Conway 2009; 59 = Duchardt et al. 2018; 60 = Duchardt et al. 2019; 61 = Geaumont et al. 2019; 62 = Copposk et al. 1983b; 63 = Chipault and Detling 2013.