

RISK ASSESSMENT OF LEAD POISONING IN RAPTORS CAUSED BY RECREATIONAL SHOOTING OF PRAIRIE DOGS

ROBERT M. STEPHENS^{1,3}, ARAN S. JOHNSON¹, REGAN E. PLUMB¹, KIMBERLY DICKERSON², MARK C. MCKINSTRY¹, AND STANLEY H. ANDERSON^{1,4}

¹*US Geological Survey, Wyoming Cooperative Fish and Wildlife Research Unit, Department 3166, 1000 East University Avenue, Laramie, WY 82071, USA. E-mail: Kimberly_dickerson@fws.gov*

²*US Fish & Wildlife Service, 5353 Yellowstone Road, Suite 308A, Cheyenne, WY 82009, USA.*

EXTENDED ABSTRACT.—Recreational shooting of Black-tailed Prairie Dogs (*Cynomys ludovicianus*) is a common activity at Thunder Basin National Grassland (TBNG), Wyoming where annual use by prairie dog shooters can be as high as 8,500 shooter-use-days. The prairie dog carcasses left in the area are scavenged by raptors and other animals susceptible to lead (Pb) poisoning if they consume Pb bullet fragments or Pb shot. In 2000, a local rehabilitator noted an increase of Pb poisoning cases in raptors from the area. We collected 22 shooter-killed prairie dog carcasses from TBNG in 2001 to determine if Pb fragments remained embedded in the tissue that potentially would be consumed by raptors. Radiographs of 19 of the 22 prairie dog carcasses showed fragments consistent with Pb.

In 2002, we conducted a more in-depth study to determine if Pb poisoning was occurring in raptors at TBNG by documenting the number of raptors on prairie dogs at colonies where shooting occurred, assaying bullet fragments in shot prairie dogs to determine Pb content, and analyzing blood and feather samples of Ferruginous Hawk (*Buteo regalis*) and Golden Eagle (*Aquila chrysaetos*) nestlings and feathers from Burrowing Owls (*Athene cunicularia*) for clinical signs of Pb poisoning. We observed raptors foraging at prairie dog colonies and collected data on the number of shooters at prairie dog colonies. Shooter intensity did not predict raptor use when compared to a site near Rawlins, Wyoming where shooting did not occur.

We detected metal fragments in 4 of 10 shot prairie dog carcasses collected. The total weight of the fragments found in each carcass ranged from 10 – 146 mg. Copper was the primary metal detected in three of four carcasses; but, significant amounts of Pb were found in the three carcasses. These fragments contained an average of 11.5 mg Pb and weighed an average of 24.7 mg with copper presumably making up the difference. Fragments <25 mg are considered small enough to be ingested and pose a risk of Pb absorption (Pauli and Buskirk 2007). Blood Pb concentrations in Ferruginous Hawk nestlings were below sub-clinical levels at TBNG and the control site. Analysis of red blood cell delta-aminolevulinic acid dehydratase activity, hemoglobin levels, and protoporphyrin levels also did not indicate Pb poisoning in Ferruginous Hawk nestlings. Additionally, blood and feather samples from Golden Eagle nestlings and feather samples from Burrowing Owls (juveniles and adults) at TBNG did not indicate Pb poisoning.

There are several possible reasons why Pb levels we found in Ferruginous Hawks and Golden Eagles (and possibly Burrowing Owls) that scavenge on the carcasses of shot prairie dogs were low. First, a sylvatic plague (*Yersinia pestis*) epizootic drastically reduced prairie dog numbers at many of the colonies in TBNG during 2001–2002. Second, 13% of TBNG was closed in 2001 to prairie dog shooting in an effort to reintroduce Black-footed Ferrets (*Mustela nigripes*). New shooting regulations and a

³Idaho Fish and Game, 3316 16th Street, Lewiston, ID 83501, USA. E-mail: rstephens@idfg.idaho.gov

⁴Deceased.

dwindling prairie dog population reduced the number of shooters that visited TBNG during the course of our study relative to previous years. Finally, long-term surveys in this area indicate that lagomorphs were abundant during our study, reducing the likelihood of raptors scavenging shot prairie

dogs. Further study is needed to determine if the occurrence of Pb poisoning in TBNG relates to prairie dog and raptor abundance, availability of alternate food sources, and regulations on shooting. Received 30 May 2008, accepted 22 October 2008.

STEPHENS, R. M., A. S. JOHNSON, R. E. PLUMB, K. DICKERSON, M. C. MCKINSTRY, AND S. H. ANDERSON. 2009. Risk assessment of lead poisoning in raptors caused by recreational shooting of prairie dogs. Extended abstract in R.T. Watson, M. Fuller, M. Pokras, and W.G. Hunt (Eds.). Ingestion of Lead from Spent Ammunition: Implications for Wildlife and Humans. The Peregrine Fund, Boise, Idaho, USA. DOI 10.4080/ilsa.2009.0212

Key words: Lead, raptors, prairie dogs, recreational shooting, scavenging.

This paper published in full as: STEPHENS, R. M., A. S. JOHNSON, R. E. PLUMB, K. DICKERSON, M. C. MCKINSTRY, AND S. H. ANDERSON. 2008. Risk assessment of lead poisoning in raptors caused by recreational shooting of prairie dogs. Intermountain Journal of Science 13(4):116-123. Also available as a US Fish and Wildlife Report: STEPHENS, R.M.; A.S. JOHNSON; R. PLUMB; K.K. DICKERSON; M.C. MCKINSTRY; AND S.H. ANDERSON. 2006. Secondary lead poisoning in Golden Eagle and Ferruginous Hawk chicks consuming shot Black-tailed Prairie Dogs, Thunder Basin National Grassland, Wyoming. US Fish and Wildlife Service Contamination Report Number R6/720C/05. Cheyenne, Wyoming. http://www.fws.gov/mountain%2Dprairie/contaminants/papers/r6ecpubs_wy.htm

LITERATURE CITED

- BECHARD, M. J., AND J. K. SCHMUTZ. 1995. Ferruginous Hawk (*Buteo regalis*). In A. Poole and F. Gill (Eds.). The Birds of North America, No. 172. The Academy of Natural Sciences, Philadelphia, and The American Ornithologists' Union, Washington, D.C., USA.
- BELLROSE, F. C. 1959. Lead poisoning as a mortality factor in waterfowl populations. Illinois Natural History Bulletin 27:235-288.
- BURCH, H. B., AND A. L. SIEGEL. 1971. Improved method for measurement of delta-aminolevulinic acid dehydratase activity of human erythrocytes. Clinical Chemistry 17:1038-1041.
- BURGER, J., AND M. GOCHFELD. 2000. Metals in Laysan Albatrosses from Midway Atoll. Arch. Environmental Contamination and Toxicology 38:254-259.
- CRAIG, T. H., J. W. CONNELLY, E. H. CRAIG, AND T. L. PARKER. 1990. Lead concentrations in Golden and Bald Eagles. Wilson Bulletin 102:130-133.
- EISLER, R. 1998. Copper hazards to fish, wildlife, and invertebrates: a synoptic review. US Geological Survey, Biological Resources Division, Biological Science Report USGS/BRD/BSR--1998-0002.
- FERNANDEZ, F. J., AND D. HILLIGOSS. 1982. Improved graphite furnace method for determination of lead in blood using matrix modification and the L'vov platform. Journal of Analytical Atomic Spectroscopy. 3:130-131.
- FORBES, R. M., AND G. C. SANDERSON. 1978. Lead toxicity in domestic animals and wildlife. Pages 225-277 in J. O. Nriagu (Ed.). The Biogeochemistry of Lead in the Environment. Part B: Biological Effects. Elsevier/ North Holland Biomedical Press, Amsterdam, New York, and Oxford.
- FRANSON, J. 1996. Interpretation of tissue lead residues in birds other than waterfowl. Pages 265-279 in W. N. Beyer, G. H. Heinz, and A. W. Redmon-Norwood (Eds.). Environmental Contaminants in Wildlife: Interpreting Tissue Concentrations. Lewis Publishers, Boca Raton, Florida, USA.

- FRANSON, J. C., W. L. HOHMAN, J. L. MOORE, AND M. R. SMITH. 1996. Efficacy of protoporphyrin as a predictive marker for lead exposure in Canvasback Ducks: Effect of sample storage time. *Environmental Monitoring Assessment* 43:181–188.
- HARMATA, A. R., AND M. RESTANI. 1995. Environmental contaminants and cholinesterase in blood of vernal migrant Bald and Golden Eagles in Montana. *Intermountain Journal of Sciences* 1:1–15.
- HENNY, C. J., L. J. BLUS, D. J. HOFFMAN, L. SILEO, D. J. AUDET, AND M. R. SNYDER. 2000. Field evaluation of lead effects on Canada Geese and Mallards in the Coeur d'Alene River Basin, Idaho. *Archives of Environmental Contamination and Toxicology* 39:97–112.
- HOFFMAN, D. J., J. C. FRANSON, O. H. PATTEE, C. M. BUNCK, AND H. C. MURRAY. 1985. Biochemical and hematological effects of lead ingestion in nestling American Kestrels (*Falco sparverius*). *Comparative Biochemistry and Physiology* 80:431–439.
- HOFFMAN, D. J., B. A. RATTNER, G. A. BURTON JR., AND J. CAIRNS JR. 1995. *Handbook of Ecotoxicology*. Lewis Publishers, Ann Arbor, Michigan, USA.
- KOCHERT, M. N., K. STEENHOF, C. L. MCINTYRE, AND E. H. CRAIG. 2002. Golden Eagle (*Aquila chrysaetos*). In A. Poole, and F. Gill (Eds.). *The Birds of North America*, No. 684. The Birds of North America, Inc., Philadelphia, Pennsylvania, USA.
- MCANNIS, D. M. 1990. Home range, activity budgets, and habitat use of Ferruginous Hawks (*Buteo regalis*) breeding in southwest Idaho. Master's thesis, Boise State University, Boise, Idaho, USA.
- MILLER, M. J. R., M. E. WAYLAND, E. H. DZUS, AND G. R. BORTOLOTTI. 2000. Availability and ingestion of lead shotshell pellets by migrant Bald Eagles in Saskatchewan. *Journal of Raptor Research* 34:167–174.
- PATTEE, O. H., AND S. K. HENNES. 1983. Bald Eagles and waterfowl: the lead shot connection. *Transactions of the North American Wildlife and Natural Resources Conference* 48:230–237.
- PAULI, J. N., AND S. W. BUSKIRK. 2007. Recreational shooting of Prairie Dogs: A portal for lead entering wildlife food chains. *Journal of Wildlife Management* 71:103–108.
- RAWSON, R. E., G. D. DELGIUDICE, H. E. DZIUK, AND L. D. MECH. 1992. Energy metabolism and hematology of White-tailed Deer fawns. *Journal of Wildlife Diseases* 28:91–94.
- SARI, M., S. DE PEE, E. MARTINI, S. HERMAN, SUGIATMI, W. M. BLOEM, AND R. YIP. 2001. Estimating the prevalence of anemia: a comparison of three methods. *Bulletin of the World Health Organization* 79:506–511.
- WAYLAND, M., AND T. BOLLINGER. 1999. Lead exposure and poisoning in Bald Eagles and Golden Eagles in the Canadian prairie provinces. *Environmental Pollution* 104:341–350.
- WYOMING STATE VETERINARY LAB. 2001. Toxicology Section, Lead Standard Operating Procedure, Laramie, Wyoming, USA.