

Index of Species Information

WILDLIFE SPECIES: Bos bison

- [Introductory](#)
 - [Wildlife Distribution and Occurrence](#)
 - [Biological Data and Habitat Requirements](#)
 - [Fire Effects and Use](#)
 - [Fire Case Study](#)
 - [References](#)
-

Introductory

WILDLIFE SPECIES: Bos bison

AUTHORSHIP AND CITATION :

Tesky, Julie L. 1995. Bos bison. In: Fire Effects Information System, [Online]. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory (Producer). Available: <http://www.fs.fed.us/database/feis/animals/mammal/bobi/all.html> [].

ABBREVIATION :

BOBI

COMMON NAMES :

American bison
bison
plains bison
wood bison
prairie bison
woodland bison
mountain bison

TAXONOMY :

The currently accepted scientific name for the American bison is *Bos bison* Linnaeus [99]. It is in the family Bovidae [34]. American bison taxonomy has been a controversial issue for many years and classification to the subspecies level remains a matter of debate [11,53]. However, most authorities recognize two subspecies, the plains bison (*Bos bison bison*) and the wood bison (*B. bison athabasca* Rhoads) [11,34,53].

ORDER :

Artiodactyla

CLASS :

Mammal

FEDERAL LEGAL STATUS :

Wood bison are listed as Threatened. The Yellowstone population is Under Review for listing as Threatened or Endangered [80].

OTHER STATUS :

In Canada, wood bison are federally listed as endangered in Alberta, British Columbia, Northwest Territories, and Yukon Territory [98]. The Nature Conservancy lists them as critically imperiled in British Columbia [77]. They are also listed in the Red Data Book by the International Union for the Conservation of Nature and Natural Resources (IUCN) and are thereby recognized world-wide as endangered. Wood bison are classified as an Appendix I animal in the Conservation on International Trade in Endangered Species of Wild Flora and Fauna (CITES), which provides regulated protection from international trade [63]. American bison are protected in Idaho [59].

WILDLIFE DISTRIBUTION AND OCCURRENCE

WILDLIFE SPECIES: Bos bison

GENERAL DISTRIBUTION :

Historically, American bison were widespread in North America from Alaska and western California across the United States and into northern New Mexico [11, 53, 66]. Today, American bison occur in geographically isolated populations in parks and preserves (See ADMINISTRATIVE UNITS), other public lands, and on private ranches. The only large herds of American bison in North America are in Yellowstone National Park, Montana and Wyoming, Wood Buffalo National Park, Alberta, and Mackenzie Bison Sanctuary and the Slave River Lowlands, Northwest Territories [11, 27, 53]. Smaller herds occur in Alaska; northeastern British Columbia; near Nahanni Butte, Northwest Territories; northwestern Saskatchewan; Elk Island National Park, Alberta; Grand Teton National Park, Wyoming; National Bison Range and Crow Reservation, Montana; Wind Caves National Park, South Dakota; Wichita Mountains Wildlife Refuge and the Tallgrass Prairie Preserve, Oklahoma; Theodore Roosevelt National Park, North Dakota; Henry Mountains, southern Utah; northeastern Colorado; Nebraska; and Kansas [11, 35, 51, 53, 83, 84, 91, 93, 96].

ECOSYSTEMS :

FRES11 Spruce-fir
 FRES19 Aspen-birch
 FRES21 Ponderosa pine
 FRES23 Fir-spruce
 FRES26 Lodgepole pine
 FRES29 Sagebrush
 FRES35 Pinyon-juniper
 FRES36 Mountain grasslands
 FRES37 Mountain meadows
 FRES38 Plains grasslands
 FRES39 Prairie
 FRES40 Desert grasslands
 FRES44 Alpine

STATES :

AK CO KS MT NE ND OK SD UT WY AB BC NT SK YT

BLM PHYSIOGRAPHIC REGIONS :

8 Northern Rocky Mountains
 9 Middle Rocky Mountains
 12 Colorado Plateau
 13 Rocky Mountain Piedmont
 14 Great Plains
 15 Black Hills Uplift
 16 Upper Missouri Basin and Broken Lands

KUCHLER PLANT ASSOCIATIONS :

K011 Western ponderosa forest
 K016 Eastern ponderosa forest
 K017 Black Hills pine forest
 K018 Pine-Douglas-fir forest
 K023 Juniper-pinyon woodland
 K024 Juniper steppe woodland
 K038 Great Basin sagebrush
 K051 Wheatgrass-bluegrass
 K052 Alpine meadows and barren
 K055 Sagebrush steppe
 K056 Wheatgrass-needlegrass shrubsteppe
 K057 Galleta-three-awn shrubsteppe
 K063 Foothills prairie
 K064 Grama-needlegrass-wheatgrass
 K065 Grama-buffalograss
 K066 Wheatgrass-needlegrass
 K067 Wheatgrass-bluestem-needlegrass
 K068 Wheatgrass-grama-buffalograss
 K069 Bluestem-grama prairie
 K070 Sandsage-bluestem prairie
 K074 Bluestem prairie
 K075 Nebraska Sandhills prairie

K081 Oak savanna
K085 Mesquite-buffalograss
K098 Northern floodplain forest

SAF COVER TYPES :

12 Black spruce
13 Black spruce-tamarack
16 Aspen
204 Black spruce
217 Aspen
218 Lodgepole pine
220 Rocky Mountain juniper
235 Cottonwood-willow
237 Interior ponderosa pine
239 Pinyon-juniper
253 Black spruce-white spruce
254 Black spruce-paper birch

SRM (RANGELAND) COVER TYPES :

101 Bluebunch wheatgrass
102 Idaho fescue
104 Antelope bitterbrush-bluebunch wheatgrass
105 Antelope bitterbrush-Idaho fescue
107 Western juniper/big sagebrush/bluebunch wheatgrass
110 Ponderosa pine-grassland
301 Bluebunch wheatgrass-blue grama
302 Bluebunch wheatgrass-Sandberg bluegrass
303 Bluebunch wheatgrass-western wheatgrass
304 Idaho fescue-bluebunch wheatgrass
305 Idaho fescue-Richardson needlegrass
306 Idaho fescue-slender wheatgrass
307 Idaho fescue-threadleaf sedge
309 Idaho fescue-western wheatgrass
310 Needle-and-thread-blue grama
311 Rough fescue-bluebunch wheatgrass
313 Tufted hairgrass-sedge
314 Big sagebrush-bluebunch wheatgrass
317 Bitterbrush-bluebunch wheatgrass
321 Black sagebrush-Idaho fescue
401 Basin big sagebrush
402 Mountain big sagebrush
403 Wyoming big sagebrush
409 Tall forb
410 Alpine rangeland
411 Aspen woodland
412 Juniper-pinyon woodland
422 Riparian
601 Bluestem prairie
602 Bluestem-prairie sandreed
603 Prairie sandreed-needlegrass
604 Bluestem-grama prairie
605 Sandsage prairie
606 Wheatgrass-bluestem-needlegrass
607 Wheatgrass-needlegrass
608 Wheatgrass-grama-needlegrass
609 Wheatgrass-grama
610 Wheatgrass
611 Blue grama-buffalograss
612 Sagebrush-grass
614 Crested wheatgrass
615 Wheatgrass-saltgrass-grama
704 Blue grama-western wheatgrass
708 Bluestem-dropseed
709 Bluestem-grama
710 Bluestem prairie
715 Grama-buffalograss
717 Little bluestem-Indiangrass-Texas wintergrass
718 Mesquite-grama
722 Sand sagebrush-mixed prairie

PLANT COMMUNITIES :

Before European settlement American bison occurred primarily on the central grasslands and northern parklands of North America, but habitats ranging from semidesert to boreal forest were also used [53]. In Montana the Lewis and Clark expedition observed vast numbers of American bison in areas floristically dominated by shortgrass species [11].

Today, American bison occupy shortgrass and tallgrass prairies, boreal parklands, montane meadows, desert grasslands, and shrub-grass habitats. In Canada, coniferous forest and quaking aspen (*Populus tremuloides*) parklands with interspersed meadows and prairies form the main habitat for wood bison [11]. In the Northwest Territories, bison habitat along the Slave River Lowlands is within the boreal forest region of Canada, where white spruce (*Picea glauca*) forests separate vast open meadows supporting sedge (*Carex* spp.) and grass communities [11]. On the Beaver Hills near Edmonton, Alberta, Hudson and Frank [40] found that American bison foraged most often on grassy upland meadows dominated by Kentucky bluegrass (*Poa pratense*) and smooth brome (*Bromus inermis*) and least in forests dominated by balsam poplar (*Populus balsamifera*) and quaking aspen [40].

American bison in Wind Caves National Park commonly occur on grasslands dominated by little bluestem (*Schizachyrium scoparium*), big bluestem (*Andropogon gerardii* var. *gerardii*), Kentucky bluegrass, and western wheatgrass (*Pascopyrum smithii*) [18].

In Yellowstone National Park, American bison inhabit meadows with sedge and grass interspersed with lodgepole pine (*Pinus contorta*) forests [11]. In southern Utah, American bison occupy pinyon-juniper (*Pinus* spp.-*Juniperus* spp.) habitat [29,72,91].

BIOLOGICAL DATA AND HABITAT REQUIREMENTS

WILDLIFE SPECIES: *Bos bison*

TIMING OF MAJOR LIFE HISTORY EVENTS :

Breeding season - Female American bison are seasonally polyestrous, with a cycle of approximately 3 weeks' duration [2,11]. However, unseasonal estrus and mating sometimes occur [2,11,51]. The breeding season for American bison generally occurs between July and October [2,11,52]. However, it varies in length depending on herd location. The breeding season has been observed to last from June 15 to September 30 at Hayden Valley, Yellowstone National Park [51]; from mid-July to mid-August in other areas of Yellowstone National Park [52]; and from June 1 to July 30 at the Wichita Mountains Wildlife Refuge [35].

Age at sexual maturity - Sexual maturity most commonly occurs at 2 to 4 years of age. However, some females breed as yearlings [11,53]. In Wood Buffalo National Park, a small proportion of the yearling bulls, approximately one-third of the 2-year-olds, and virtually all American bison 3 years of age and older were sexually mature [11]. Bulls attain sexual maturity well in advance of becoming part of the active breeding population [52]. In the Hayden Valley American bison herd, Yellowstone National Park, bulls 8 years old and older were the most active sexually [51].

Gestation and calving season - The gestation period for American bison is 9 to 9.5 months [11,51,53]. In most regions, the calving season is generally from mid-April to May with births concentrated from the end of April through the first 2 weeks of May [53]. However, conception and, therefore, parturition can occur at any time of year [11]. McHugh [51] noted that a few calves were born from June through October in herds at Yellowstone National Park, the Crow Reservation, Montana, the National Bison Range, and Wind Caves National Park.

Number of young - Females generally give birth to one calf; twins are rare [11,53]. American bison generally produce two calves every 3 years [11].

Development of young - Calves are able to stand and nurse within 30 minutes of birth. They may try to graze by 5 days of age and will drink water after the first week. The cow nurses her calf for at least 7 to 8 months; most calves are weaned by the end of the first year [53].

Longevity - In the wild few American bison survive more than 20 years, although there are records of cows surviving at least 40 years [53]. In wild populations, by the time a American bison has reached age 15 it can be considered old. In captivity, lifespan increases [11].

PREFERRED HABITAT :

The primary habitat requirements for American bison are adequate forage, water, and space [8]. American bison thrive in open grasslands, meadows, and parklands.

When available, American bison use forested areas for shade and to escape insects. These areas also furnish forage when open areas are covered by "deep" snow [8]. However, American bison can survive in open valleys covered with several feet of snow [94]. During severe weather, forested areas and some topographical features furnish cover [8]. Scattered thermal sites (particularly warm ground with less snow than the surrounding area) provide favorable sites for wintering American bison in Yellowstone National Park [94]. In northern Canada forested and shrub covered areas are often used as daily and seasonal travel corridors [11].

Most American bison are seasonally migratory; movements are both directional and altitudinal in some regions [11,52,70]. During historical times, large herds of American bison commonly moved southward 200 miles (322 km) or more to winter range [2]. Directional and altitudinal movements between summer and winter ranges still occur annually at Wood Buffalo National Park and Yellowstone National Park [2,11,52,94]. During November and May, American bison at Wood Buffalo National Park migrate from wooded hills to the Peace River Valley, a distance of as much as 150 miles (241 km) [2]. Factors that may influence seasonal migrations include tradition, supply and accessibility of forage, open water, shelter, insect harassment, spring weather conditions and temperatures, and fall snowstorms at higher elevations [53,58].

In mountainous areas, altitudinal movements to lowland winter range in fall and to higher summer range in spring are quite common. Snipe flies (*Symphoromyia* spp.) may be responsible for some altitudinal movements by Yellowstone National Park American bison herds during the summer [52]. Large, windswept prairies may also be chosen in summer for relief from insects. American bison, particularly cows, show strong affinity to traditional winter range [52,70]. Shaw and Carter [70] found that older females appear to be more prone than younger ones to seek new winter range and return to the new range in subsequent winters.

COVER REQUIREMENTS :

American bison thrive in open areas. When available, shrub or forest cover may be used for shade, escape from insects, or shelter from severe weather [8].

FOOD HABITS :

Adult American bison require approximately 30 pounds of forage every day [25]. Where grasses and sedges are available in the habitat, they are selectively grazed by American bison, and where they are sparse, browse may be substituted [8,11,20,53,88].

Dietary shifts from grasses to sedges and back again within a habitat type are usually associated with plant phenology [11]. In Yellowstone National Park, sedges comprised the highest proportion of American bison diets in all seasons, while grasses were second in importance. Minor quantities of forbs (6%) and browse (2%) were consumed, mainly in summer [52]. In northeastern Colorado, sedges were important to American bison only during spring. American bison herds located at Wood Buffalo National Park and Elk Island National Park were observed feeding on grasses in summer and sedges in winter [11].

In the Slave River Lowlands, American bison diets contained 29 different plant species. Slough sedge (*Carex atherodes*) was the most abundant plant in the diet, varying from 42 percent in winter to 77 percent in spring. The second most common food was reedgrass (*Calamagrostis* spp.), which varied from 15 percent of the diet in spring to 35 percent in winter. Together, these two forages comprised more than 70 percent of the American bison diet at all seasons [64].

Diet and habitat selection of wood bison were studied in the Mackenzie Bison Sanctuary between February 1986 and April 1988. Wood bison showed pronounced seasonal changes in their diet. Sedges constituted 96.1 to 98.8 percent of the winter diet. During the summer, the diet became a more diverse mix of sedge (*Carex* spp.), grasses (*Poaceae*), and willow (*Salix* spp.). Reindeer lichen (*Cladonia* spp.) became a major dietary component in fall. Summer browsing on willows increased when sedge standing crops were reduced [45].

In some areas forbs are seasonally important to foraging American bison [11,89]. In semidesert range in southwestern Colorado, forbs were common food items during all seasons but never exceeded 17 percent in any one season [89]. In Yellowstone National Park and in northern Canada, forbs appeared to be important to American bison only during summer; in northeastern Colorado, forbs were important during fall and winter [11].

On the shortgrass plains in Colorado, where blue grama (*Bouteloua*

gracilis) is the dominant species, American bison consumed 36 different plant species. However, only 11 contributed significantly to the total. Blue grama and buffalo grass (*Buchloe dactyloides*) were the most abundant plants in the habitat and also in the diet. Where available, western wheatgrass was preferred over blue grama. Other commonly consumed species were red threeawn (*Aristida purpurea*), sun sedge (*Carex heliophila*), scarlet globemallow (*Sphaeralcea coccinea*), sand dropseed (*Sporobolus cryptandrus*), needle-and-thread grass (*Stipa comata*), fringed sagebrush (*Artemisia frigida*), buckwheat (*Eriogonum* spp.), and sixweeks fescue (*Vulpia octoflora*) [61].

On semidesert range at Colorado National Monument, the most common plant species in the American bison diet during most seasons was fourwing saltbush (*Atriplex canescens*), followed by needle-and-thread grass, which was important during cooler months. Sand dropseed and galleta (*Hilaria jamesii*) were prominent in the diet in warmer seasons. The only forbs significantly utilized during all seasons except winter were mallows (*Sphaeralcea* spp.). Prickly pear (*Opuntia* spp.) was among the ten top forage species during all seasons except summer. Some of the most common plants in the habitat, cheatgrass (*Bromus tectorum*), Utah juniper (*Juniperus osteosperma*), and big sagebrush (*Artemisia tridentata*), were the least preferred forages [89].

PREDATORS :

The main predators of American bison are gray wolf (*Canis lupus*), [10,11,38,65] grizzly bear (*Ursus arctos horribilis*), and coyote (*Canis latrans*) [11]. On the Slave River Lowlands, American bison were the most important of six major prey types eaten by gray wolves and represented 88 percent of prey weight during the period of snow cover (Nov. 8, 1976-April 15, 1977). Gray wolf predation in 1976 and 1977 accounted for about 31 percent of adult and subadult American bison mortality and approximately 27 percent of calf mortality [65]. Winter-killed American bison may be important food sources to grizzly bear in early spring after they emerge from their dens [11].

MANAGEMENT CONSIDERATIONS :

American bison are attracted to grassland sites altered by black-tailed prairie dog (*Cynomys ludovicianus*). American bison often feed selectively near the perimeters of colonies. These areas are constantly clipped by black-tailed prairie dog and, therefore, have more readily digestible perennial grasses, with higher nitrogen concentration and greater accessibility of green tissues, than vegetation from uncolonized areas. Consequently, black-tailed prairie dog colonies may receive a disproportionately high amount of American bison use. Prolonged grazing pressure on colonies may result in changes in plant composition [18,19,12,87]. On a mixed-grass prairie, selective use of black-tailed prairie dog colonies by American bison resulted in considerably more biomass removed than by prairie dog activity alone. Additionally, selective use of plant species (i.e., grasses) by American bison may contribute to an increase in forb:graminoid ratios [95].

The thick hair on the head and forequarters of American bison is ideally suited for dispersal of awned, barbed, or sticky seed-bearing structures. For example, the seeds of buffalo grass, cocklebur (*Xanthium perforatum*), and St. Johnswort (*Hypericum perforatum*) readily adhere to American bison hair. The dissemination of the latter throughout the National Bison Range is thought to have been accomplished by American bison [51]. American bison also aid in dispersal by ingesting seeds. Samples from four buffalo chips at Wichita Mountains Wildlife Refuge contained 219 seeds that germinated. Of these, 195 (89%) were monocots, mostly Japanese brome (*Bromus japonicus*). American bison may accelerate seed dispersal to burned sites because American bison are attracted to recently burned areas [17].

Localized stands of timber may be considerably affected by American bison horning and thrashing during the rut and at other times. McHugh [51] estimated that 51 percent of lodgepole pine in some areas of Yellowstone National Park has been horned by American bison. Such activity may inhibit succession of grassland to forest [11].

Where American bison trails or wallows (concave disturbances formed as American bison paw the ground and roll in the exposed soil) are cut into steep hillsides, considerable water and wind erosion can occur. Hillside trails can serve as drainage channels, effectively lowering the water table in upland areas and causing a change in the vegetation. Where trails cut near the top of steep, sandy hills, erosion and slippage may produce barren areas. However, by creating trails through different habitats, American bison help provide access corridors for many species of mammals, including humans [11].

American bison wallows can serve as water catchments on flat terrain. In Oklahoma American bison wallows have been observed to hold water for prolonged periods during the spring rainy season. Such small ponds become available to both vertebrates and invertebrates. These water-holding wallows may also enhance growth of specific vegetation such as ruderal species and species adapted to wet habitats [11,81]. On the Wichita Mountains Wildlife Refuge, ruderal species such as Japanese brome and false-pennyroyal (*Hedeoma hispida*) had highest cover values within American bison wallows. Other common taxa within the wallows were Torrey rush (*Juncus torreyi*), purple ammania (*Ammannia coccinea*), lythrum (*Lythrum* spp.), and taperleaf flatsedge (*Cyperus acuminatus*), all of which are species adapted to wet habitats [17].

Diseases - Anthrax (an infectious disease caused by the bacteria *Bacillus anthracis*) outbreaks cause sporadic mortality in northern American bison herds. In Wood Buffalo National Park, 50 percent of American bison may be infected by tuberculosis, a chronic infectious disease [53]. Tuberculosis in a herd of American bison for more than 26 years did not appear to interfere with herd productivity. However, the importance of tuberculosis as a mortality factor is difficult to determine for large American bison herds [11]. Brucellosis is an infectious disease caused by the bacteria *Brucella abortus*. Abortion caused by brucellosis has been reported in American bison. It is assumed that infected American bison shed brucella organisms, thereby contaminating feed and water. Dissemination of the disease is enhanced due to the gregarious nature of American bison [11]. The role of brucellosis and its affect on reproductive activity in American bison is difficult to determine due to the lack of data on the incidence of abortion in American bison [11,52].

FIRE EFFECTS AND USE

WILDLIFE SPECIES: Bos bison

DIRECT FIRE EFFECTS ON ANIMALS :

Fires commonly occur on American bison ranges without causing appreciable American bison mortality [11]. In the past, when large herds of American bison roamed the prairies, some prairie fires killed hundreds of American bison [11,13,57]. One report in 1850 stated that as many as 300 American bison were seen lying together on the ground with their hair burned off by a prairie fire [11]. There were only nine known direct American bison mortalities due to the 1988 fires in Yellowstone National Park [56].

HABITAT RELATED FIRE EFFECTS :

Fire is important in creating and maintaining American bison habitat. Fire regenerates grasslands and enhances production, availability, and palatability of many American bison forage species [9,11,67,68]. Fire frequency has been estimated to occur once every 3 to 5 years on some prairies [97]. During presettlement times American bison habitats were to a large extent created and maintained by lightning-caused fires or fires set by Native Americans [44,47,49,50]. The results of intense grazing by large American bison herds on recently burned areas may have reduced fuel loads, making the grazed areas less likely to burn and even allowing them to function as firebreaks [49,75]. In contrast, unburned areas would have been little grazed, thereby increasing fuel loads and the probability of burning. The slaughter of American bison in the late 1800's may have shortened fire return intervals and increased fire severity during the early settlement period. Steuter [75] suggested that integrating a regional fire behavior model with estimates of presettlement American bison patterns could provide a valuable tool for natural area management in the northern mixed-grass prairie.

Several studies have shown that American bison prefer to forage on recently burned areas [5,7,22,28,70,87]. In tallgrass prairie on the Konza Prairie Research Natural Area, northeastern Kansas, 45 American bison range over an array of watersheds with different fire regimes [87]. The watersheds are burned in April annually or at 2-, 4-, or 20-year intervals. In the spring of 1988 and 1989, Vinton and others [87] studied American bison grazing and use patterns among these watersheds as influenced by fire regime. American bison used some watersheds preferentially and the pattern of watershed use changed seasonally. During the spring of both years (April-June 30), American bison selected only watersheds that had been recently burned (annually or biennially), and were observed up to three times more frequently than expected on these watersheds. In 1988, preferential grazing of recently burned watersheds persisted through the summer

months. During autumn and winter of both years, American bison preferred the annual and 20-year burn watersheds to watersheds that were burned every 2 or 4 years.

On the same study site as above, little bluestem was sampled to determine how fire influences its use by American bison and its responses to grazing. Plants were marked at the beginning of the 1992 growing season. Little bluestem was sampled in an annually burned watershed and a watershed burned at 4-year intervals (referred to as "unburned") that had been grazed by American bison since 1987, and nearby annually burned and 4-year burn interval watershed that were ungrazed. The 4-year burn interval watersheds had last burned 2 years before sampling. On unburned prairie, American bison grazed only 5 percent of the available little bluestem, selecting it only 30 percent as frequently as big bluestem, the codominant species. On burned prairie, grazing frequency of little bluestem was more than 3 times as great as on unburned sites and equal to that of big bluestem. The increased grazing frequency on little bluestem in recently burned prairie is most likely the result of the removal of its persistent standing dead tillers by burning. Burning did not affect grazing on big bluestem, a plant lacking persistent standing dead tillers. With longer intervals between fires, American bison might display even greater avoidance of little bluestem in favor of other grasses [93].

A combination of fire and American bison grazing may increase the standing crop of rhizomatous grasses at the expense of bunchgrasses. Pfeiffer and Steuter [96] conducted a study on Nebraska sandhills during the 1991 and 1992 growing seasons to determine the response of sandhills prairie vegetation to spring and summer prescribed burns and subsequent American bison grazing. Approximately 1,235 acres (500 ha) were burned in early May, and another 247 acres (100 ha) were burned in late July, 1991. During the 1992 growing season, American bison grazing on burned areas reduced bunchgrass standing crop by 56 percent, while reducing rhizomatous grass standing crop by only 18 percent. Forbs generally appeared unaffected by American bison grazing. The increased grazing pressure by American bison lasted only one season. Rhizomatous grasses of the Great Plains are better adapted to large herbivore grazing than are bunchgrasses. Burning and grazing would increase the amount of forage available since, in unburned prairie, standing dead tillers deter use of bunchgrasses.

Several studies concerning American bison response to prescribed fire have been conducted at Wind Cave National Park [5,28,32]. Two prescribed fires in ponderosa pine (*Pinus ponderosa*)-grassland habitat were conducted on October 16, 1974 and May 9, 1975. The spring fire was conducted on a site adjacent to the fall fire. American bison were noted in the area of the burns during the course of burning. They utilized regrowth vegetation on the burned areas throughout the summer of 1975 [32]. A prescribed fire conducted on April 1, 1981, burned 110 acres (44.5 ha) of mixed-grass prairie and 134 (54.4 ha) of forest land. American bison fed within the burn in 1981 and 1982, moving in 1983 to an area burned by wildfire [28].

On the Wichita Mountains Wildlife Refuge, Shaw and Carter [70] studied seasonal range use by American bison before and after spring prescribed fires on a mixed-grass prairie interspersed with post oak (*Quercus stellata*)-blackjack oak (*Q. marilandica*) woodlands. American bison increased use of the burned portion of their summer range. They showed no apparent response to prescribed burning of an area of new winter range, but they delayed their spring departure to traditional summer range.

Some studies have shown that cow-calf herds graze burned areas more often than bulls [5,18]. The first postfire years following a fall prescribed fire in grassland habitat at Wind Cave National Park, bulls were found less than cow-calf herds on burned sites. Both cow-calf herds and bull groups tended to use the burn more in June of the first postfire season than at any other time. However, only cow-calf herds consistently grazed the burn during the rest of the summer [18].

Wallows enhance species diversity in American bison habitat. In Wichita Mountains National Wildlife Refuge, ruderal species (e.g., Japanese brome and false pennyroyal) and mesic species (e.g., purple ammania, pepperwort [*Marsilea mucronata*], and seacoast sumpweed) had higher cover values within wallow than outside them. Wallows may be especially abundant and heavily used on burned sites because American bison are attracted to graze in such areas [16]. Collins and Uno [16] examined the effects of February, 1982, prescribed fire on wallow vegetation in Wichita Mountains Wildlife Refuge. Vegetation samples taken during June and early July, 1982, from the edge and interior of unburned wallows were more similar to each other than were edge and interior samples from burned wallows. Species diversity and richness were significantly lower in burned than in unburned wallows. Winter annuals were more abundant

in unburned wallows, perhaps because they were burned during their growing season. The authors suggested that spring fires may reduce cover of winter annuals in wallows, but summer and fall fires could increase their importance [16].

Sedge-grasslands, which are important winter habitat for American bison, often increase in area after fire removes surrounding shrubs or trees [9]. Fires in open black spruce (*Picea mariana*) forests and shrublands may result in expansion of sedge-grasslands. In 1977, the Bear Creek wildfire near Farewell, Alaska, moderately to severely burned a closed spruce-hardwood forest and an open black spruce forest with an understory of willow, shrubs, and sedges. The fire converted 100 square miles (260 km sq.) of predominantly open black spruce forest to sedge-grassland. Most of the American bison in this area winter on sites with extensive sedge cover. By postfire year 4 the sedge-grassland habitat had more than doubled in area. Fire-related snowpack changes also may have stimulated American bison winter range expansion. Before the fire, the disjunct and widely scattered sedge-grasslands were separated by extensive open black spruce forest and shrublands. This habitat generally has a greater snowpack than sedge-grasslands and, therefore, is likely to discourage American bison movements. After the 1977 fire, sedge-grasslands showed less snow cover than adjacent unburned open black spruce forests and shrublands [9].

FIRE USE :

Prescribed fire has been used to manage free-roaming American bison herds [6, 18, 46, 70]. Strategic placement of burns should integrate knowledge of American bison foraging behavior and preferences, American bison travel routes, and distributions of mineral licks and water [6, 18]. Using prescribed fire to improve grasses and sedges may reduce the need for expensive supplemental feeding of American bison in some areas [68].

Prescribed fire is effective in mitigating American bison impacts on black-tailed prairie dog colonies. American bison use of a black-tailed prairie dog colony was compared before and after a prescribed fire on adjacent, uncolonized grassland at Wind Cave National Park, in 1979 and 1980. Cow-calf herds increased their use (measured as hours of feeding time) of the burned grassland by a factor of 12 and decreased their use of the colony by 30 to 63 percent following the burn. Bulls were less attracted to the burned site than cow-calf herds. To decrease American bison impacts on black-tailed prairie dog colonies, burns should be located a "considerable" distance from colonies [18].

FIRE REGIMES :

Find fire regime information for the plant communities in which this species may occur by entering the species name in the [FEIS home page](#) under "Find Fire Regimes".

FIRE CASE STUDY

SPECIES: Bos bison

FIRE CASE STUDY CITATION :

Tesky, Julie L., compiler. 1995. Bos bison. Improving bison forage with fall prescribed fire in Wind Cave National Park, South Dakota

FIRE CASE STUDY REFERENCE:

Bock, Jane H.; Bock, Carl E. 1981. Some effects of fire on vegetation and wildlife in ponderosa pine forests of the southern Black Hills. Final Report. Contracts CX-1200-9-B034, CX-1200-0-B018, CX-1200-1-B022; Grant No. RM-80-105 GR. Unpublished report on file with: U.S. Department of Agriculture, Forest Service, Intermountain Research Station, Fire Sciences Lab, Missoula, MT. 58 p. [5].

SEASON/SEVERITY CLASSIFICATION :

Fall/low-moderate

STUDY LOCATION :

The study was located on a small knob in the northwest corner of Wind Cave National Park, Custer County, South Dakota.

PREFIRE HABITAT :

This study was located in a ponderosa pine (*Pinus ponderosa*) forest and ponderosa pine-grassland ecotone. The dominant species in the canopy was ponderosa pine. Common understory species included sedges (*Carex* spp.), big bluestem (*Andropogon gerardii* var. *gerardii*), little bluestem (*Schizachyrium scoparium*), Sandberg bluegrass (*Poa secunda*), sideoats grama (*Bouteloua curtipendula*), blue grama (*B. gracilis*), hairy grama (*B. hirsuta*), needlegrass (*Stipa* spp.), wheatgrass, leadplant (*Amorpha canescens*), pulsatilla (*Pulsatilla patens*), Louisiana sagewort (*Artemisia ludoviciana*), beautiful shootingstar (*Dodecatheon pulchellum*), and western snowberry (*Symphoricarpos occidentalis*).

SITE DESCRIPTION :

A detailed description of the site was not given.

FIRE DESCRIPTION :

Conditions for the prescribed fire (the "Cone Burn") were met at midday 17 October 1979, when the wind was about 10 miles per hour (16 km/h), temperature was 58 degrees Fahrenheit (14.4 deg C), and relative humidity was 45 percent. The fire passed over a 156-acre (63 ha) area in about 4 hours. A few standing dead snags on the north and west faces of the area were still burning at midmorning the following day, when they were extinguished by a rain shower. Fire varied from discontinuous, low-severity (leaving some patches unburned) to crowning behavior. Combustion of surface fuels was nearly complete, except in unburned patches.

FIRE EFFECTS ON ANIMAL SPECIES AND HABITAT :

Effect on vegetation - Vegetation was measured 1 year before the fire and for 2 years after (May-June of 1980 and 1981) the fire. Plants that increased as a result of the fire included sedges, needlegrasses, and beautiful shootingstar; bare ground also increased. Plants reduced by the fire included bluegrasses (*Poa* spp.), little bluestem (1980 only), ponderosa pine (both canopy and immature trees), and shrubs (1980 only). The fire caused only a modest decline in overall similarity between experimental and control vegetation plots, and this decline persisted through two postfire growing seasons.

Effect on American bison - American bison preferred feeding on burned sites during the first postfire growing season. Before the fire, American bison cow-calf herds preferred the control areas. After burning, they were attracted to the recently burned area for feeding. However, this grazing preference disappeared in the second postfire year. Before the fire, bulls preferred the control site. During the first postfire year the bulls used both sites, while in the following year, they showed a preference for the unburned site. Small bull groups tended to defer in habitat use to cow-calf herds.

FIRE MANAGEMENT IMPLICATIONS :

This study showed that prescribed fires can be used in ponderosa pine-grassland ecotones of the Black Hills to temporarily improve forage for American bison.

REFERENCES

WILDLIFE SPECIES: Bos bison

REFERENCES :

1. Alexander, Martin E.; Dube, Dennis E. 1983. Fire management in wilderness areas, parks, and other nature reserves. In: Wein, Ross W.; MacLean, David A., eds. The role of fire in northern circumpolar ecosystems. Scope 18. New York: John Wiley & Sons: 273-297. [18711]
2. Banfield, A. W. F. 1974. The mammals of Canada. Toronto, ON: University of Toronto Press. 438 p. [21084]
3. Bernard, Stephen R.; Brown, Kenneth F. 1977. Distribution of mammals, reptiles, and amphibians by BLM physiographic regions and A.W. Kuchler's associations for the eleven western states. Tech. Note 301. Denver, CO: U.S. Department of the Interior, Bureau of Land Management. 169 p. [434]

4. Bock, J. H. 1987. Fire as an evolutionary force in North American grasslands. *American Journal of Botany*. 74(5): 661. [Abstract]. [24524]
5. Bock, Jane H.; Bock, Carl E. 1981. Some effects of fire on vegetation and wildlife in ponderosa pine forests of the southern Black Hills. Final Report. Contracts CX-1200-9-B034, CX-1200-0-B018, CX-1200-1-B022; Grant No. RM-80-105 GR. Unpublished report on file with: U.S. Department of Agriculture, Forest Service, Intermountain Research Station, Fire Sciences Lab, Missoula, MT. 58 p. [479]
6. Bone, Steven D.; Klukas, Richard W. 1990. Prescribed fire in Wind Cave National Park. In: Alexander, M. E.; Bisgrove, G. F., technical coordinators. The art and science of fire management: Proceedings, 1st Interior West Fire Council annual meeting and workshop; 1988 October 24-27; Kananaskis Village, AB. Inf. Rep. NOR-X-309. Edmonton, AB: Forestry Canada, Northwest Region, Northern Forestry Centre: 297-302. [14145]
7. Boyce, Mark S.; Merrill, Evelyn H. 1991. Effects of the 1988 fires on ungulates in Yellowstone National Park. In: Proceedings, 17th Tall Timbers fire ecology conference; 1989 May 18-21; Tallahassee, FL. Tallahassee, FL: Tall Timbers Research Station: 121-132. [17604]
8. Boyd, Raymond J.; Cooperrider, Allen Y.; Lent, Peter C.; Bailey, James A. 1986. Ungulates. In: Cooperrider, Allen Y.; Boyd, Raymond J.; Stuart, Hanson R., eds. Inventory and monitoring of wildlife habitat. Denver, CO: U.S. Department of the Interior, Bureau of Land Management, Service Center: 519-564. [10856]
9. Campbell, Bruce H.; Hinkes, Mike. 1983. Winter diets and habitat use of Alaska bison after wildfire. *Wildlife Society Bulletin*. 11(1): 16-21. [8389]
10. Carbyn, L. N. 1987. Responses of bison on their calving grounds to predation by wolves in Wood Buffalo National Park. *Canadian Journal of Zoology*. 65: 2072-2078. [24527]
11. Chapman, Joseph A.; Feldhamer, George A., eds. 1982. Wild mammals of North America. Baltimore, MD: The Johns Hopkins University Press. 1147 p. [21085]
12. Cid, M. Silvia; Detling, James K.; Whicker, April D.; Brizuela, Miguel A. 1991. Vegetation responses of a mixed-grass prairie site following exclusion of prairie dogs and bison. *Journal of Range Management*. 44(2): 100-104. [14133]
13. Cole, James E. 1954. Buffalo (Bison bison) killed by fire. *Journal of Mammalogy*. 35(3): 453-454. [16152]
14. Collins, Scott L.; Barber, Susan C. 1985. Effects of disturbance on diversity in mixed-grass prairie. *Vegetatio*. 64: 87-94. [2707]
15. Collins, Scott L.; Gibson, David J. 1990. Effects of fire on community structure in tallgrass and mixed-grass prairie. In: Collins, Scott L.; Wallace, Linda L., eds. Fire in North American tallgrass prairies. Norman, OK: University of Oklahoma Press: 81-98. [14196]
16. Collins, Scott L.; Uno, Gordon E. 1983. The effect of early spring burning on vegetation in buffalo wallows. *Bulletin of the Torrey Botanical Club*. 110(4): 474-481. [4352]
17. Collins, Scott L.; Uno, Gordon E. 1985. Seed predation, seed dispersal, and disturbance in grasslands: a comment. *American Naturalist*. 125(6): 866-872. [665]
18. Coppock, D. Layne; Detling, James K. 1986. Alteration of bison and black-tailed prairie dog grazing interaction by prescribed burning. *Journal of Wildlife Management*. 50(3): 452-455. [689]
19. Coppock, D. L.; Ellis, J. E.; Detling, J. K.; Dyer, M. I. 1983. Plant-herbivore interactions in a North American mixed-grass prairie. II. Responses of bison to modification of vegetation by prairie dogs. *Oecologia*. 56: 10-15. [688]
20. Detling, J. K.; Dyer, M. I.; Procter-Gregg, C.; Winn, D. T. 1980. Plant-herbivore interactions: examination of potential effects of bison saliva on regrowth of *Bouteloua gracilis* (H.B.K.) Lag. *Oecologia*. 45: 26-31. [5000]
21. Eaheart, David. 1992. Mystery grass turns into business. *Rangelands*.

14(2): 103-104. [19395]

22. Easterly, Thomas G.; Jenkins, Kurt J. 1991. Forage production and use on bighorn sheep winter range following spring burning in grassland and ponderosa pine habitats. *Prairie Naturalist*. 23(4): 193-200. [19277]
23. Edwards, Tom. 1978. Buffalo and prairie ecology. In: Glenn-Lewin, David C.; Landers, Roger Q., Jr., eds. *Proceedings, 5th Midwest prairie conference; 1976 August 22-24; Ames, IA. Ames, IA: Iowa State University: 110-112. [3359]*
24. England, Raymond E.; DeVos, Antoon. 1969. Influence of animals on pristine conditions on the Canadian grasslands. *Journal of Range Management*. 22: 87-94. [865]
25. Evans, Keith E.; Probasco, George E. 1977. *Wildlife of the prairies and plains. Gen. Tech. Rep. NC-29. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station. 18 p. [14118]*
26. Eyre, F. H., ed. 1980. *Forest cover types of the United States and Canada. Washington, DC: Society of American Foresters. 148 p. [905]*
27. Finch, Deborah M. 1992. *Threatened, endangered, and vulnerable species of terrestrial vertebrates in the Rocky Mountain Region. Gen. Tech. Rep. RM-215. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station. 38 p. [18440]*
28. Forde, Jon D.; Sloan, Norman F.; Shown, Douglas A. 1984. *Grassland habitat management using prescribed burning in Wind Cave National Park, South Dakota. Prairie Naturalist. 16(3): 97-110. [938]*
29. Frischknecht, Neil C. 1975. *Native faunal relationships within the pinyon-juniper ecosystem. In: The pinyon-juniper ecosystem: a symposium: Proceedings; 1975 May; Logan, UT. Logan, UT: Utah State University, College of Natural Resources, Utah Agricultural Experiment Station: 55-56. [974]*
30. Gainer, Bob. 1985. *Free-roaming bison in northern Alberta. Alberta Naturalist. 15(3): 86-87. [24523]*
31. Garrison, George A.; Bjugstad, Ardell J.; Duncan, Don A.; [and others]. 1977. *Vegetation and environmental features of forest and range ecosystems. Agric. Handb. 475. Washington, DC: U.S. Department of Agriculture, Forest Service. 68 p. [998]*
32. Gartner, F. Robert. 1977. *Ecological changes on pine grassland burned in fall and spring. Final Report Contract No. PX 120051027, U.S. Department of the Interior National Park Service, Rocky Mountain Regional Office. Rapid City, SD: South Dakota State University, Agricultural Research and Extension Center. 35 p. [1001]*
33. Feldhake, C. M.; Boyer, D. G. 1995. *Soil water depletion by C3 and C4 pasture grasses in central Appalachia. Journal of Soil and Water Conservation. 50(1): 106-109. [24529]*
34. Hall, E. Raymond. 1981. *The mammals of North America. 2nd ed. Vol. 2. New York: John Wiley and Sons. 1271 p. [14765]*
35. Halloran, Arthur F.; Glass, Bryan P. 1959. *The carnivores and ungulates of the Wichita Mountains Wildlife Refuge, Oklahoma. Journal of Mammalogy. 40(3): 360-370. [24597]*
36. Hanson, William A. 1979. *Preliminary results of the Bear Creek fire effects studies. Proposed open file report. Anchorage, AK: U.S. Department of the Interior, Bureau of Land Management, Anchorage District Office. 83 p. [6400]*
37. Haugen, Arnold O.; Shult, Milo J. 1973. *Approximating pre-white-man animal influences and relationships in prairie natural areas. In: Hulbert, Lloyd C., ed. Thrid Midwest prairie conference proceedings; 1972 September 22-23; Manhattan, KS. Manhattan, KS: Kansas State University, Division of Biology: 17-19. [3326]*
38. Herman, Margaret, Willard, E. Earl. 1978. *Rocky Mountain wolf and its habitat. Missoula, MT: U.S. Department of Agriculture, Forest Service, National Forest System Cooperative Forestry, Forestry Research, Region 1. 17 p. [16522]*
39. Houston, Douglas B. 1971. *Ecosystems of national parks. Science. 172:*

648-651. [16074]

40. Hudson, R. J.; Frank, S. 1987. Foraging ecology of bison in aspen boreal habitats. *Journal of Range Management*. 40(1): 71-75. [1201]
41. Knapp, A. K.; Seastedt, T. R. 1986. Detritus accumulation limits productivity of tallgrass prairie. *BioScience*. 36(10): 662-668. [2880]
42. Kruse, Arnold D.; Higgins, Kenneth F. 1990. Effects of prescribed fire upon wildlife habitat in northern mixed-grass prairie. In: Alexander, M. E.; Bisgrove, G. F., technical coordinators. The art and science of fire management: Proceedings, 1st Interior West Fire Council annual meeting and workshop; 1988 October 24-27; Kananaskis Village, AB. Inf. Rep. NOR-X-309. Edmonton, AB: Forestry Canada, Northwest Region, Northern Forestry Centre: 182-193. [14146]
43. Kuchler, A. W. 1964. Manual to accompany the map of potential vegetation of the conterminous United States. Special Publication No. 36. New York: American Geographical Society. 77 p. [1384]
44. Higgins, Kenneth F.; Kruse, Arnold D.; Piehl, James L. 1989. Effects of fire in the Northern Great Plains. Ext. Circ. EC-761. Brookings, SD: South Dakota State University, Cooperative Extension Service, South Dakota Cooperative Fish and Wildlife Research Unit. 47 p. [14749]
45. Larter, Nicholas C.; Gates, Cormack C. 1991. Diet and habitat selection of wood bison in relation to seasonal changes in forage quantity and quality. *Canadian Journal of Zoology*. 69: 2677-2685. [24528]
46. Larson, Lorence; Murdock, G. K. 1989. Small bison herd utilization of tallgrass prairie. In: Bragg, Thomas B.; Stubbendieck, James, eds. *Prairie pioneers: ecology, history and culture: Proceedings, 11th North American prairie conference; 1988 August 7-11; Lincoln, NE.* Lincoln, NE: University of Nebraska: 243-245. [14055]
47. Little, Silas. 1974. Effects of fire on temperate forests: northeastern United States. In: Kozlowski, T. T.; Ahlgren, C. E., eds. *Fire and ecosystems*. New York: Academic Press: 225-250. [9859]
48. Lowe, David W.; Matthews, John R.; Moseley, Charles J. 1990. The official World Wildlife Fund guide to endangered species of North America. Volume 1. Plants: Mammals. Washington, DC: Beacham Publishing, Inc. 556 p. [+ appendices]. [23209]
49. Madson, John. 1990. On the Osage. *Nature Conservancy Magazine*. 40(3): 7-15. [11799]
50. McCormack, Patricia A. 1992. The political economy of bison management in Wood Buffalo National Park. *Arctic*. 45(4): 367-380. [21080]
51. McHugh, Tom. 1958. Social behavior of the American buffalo (Bison bison). *Zoologica*. 43(1): 1-40. [3981]
52. Meagher, Mary M. 1973. The bison of Yellowstone National Park. *Scientific Monograph Series 1*. [Denver, CO]: U.S. Department of the Interior, National Park Service. 161 p. [24554]
53. Meagher, Mary. 1986. Bison bison. *Mammalian Species*. 266: 1-8. [24519]
54. Meagher, Mary. 1989. Evaluation of boundry control for bison of Yellowstone National Park. *Wildlife Society Bulletin*. 17: 15-19. [6887]
55. Merrill, Evelyn H.; Wilson, Cathy; Marrs, Ronald W. 1990. Remote sensing of vegetation recovery in grasslands after the 1988 Yellowstone fires in Yellowstone National Park. In: Boyce, Mark S.; Plumb, Glenn E., eds. *National Park Service Research Center, 14th annual report*. Laramie, WY: University of Wyoming, National Park Service Research Center: 151-153. [15435]
56. Mills, Susan M., editor. 1989. The Greater Yellowstone postfire assessment. [Denver, CO]: U.S. Department of Agriculture, Forest Service, Northern Region. [Pages unknown]. In cooperation with: U.S. Department of the Interior, National Park Service, Grand Teton and Yellowstone National Parks. [24521]
57. Moore, Conrad Taylor. 1972. Man and fire in the central North American grassland 1535-1890: a documentary historical geography. Los Angeles, CA: University of California. 155 p. Dissertation. [5732]
58. Morgan, R. Grace. 1980. Bison movement patterns on the Canadian plains: an ecological analysis. *Plains Anthropology*. 25: 142-160. [1694]

59. Moseley, Robert; Groves, Craig, compilers. 1990. Rare, threatened and endangered plants and animals of Idaho. Boise, ID: Idaho Department of Fish and Game, Nongame and Endangered Wildlife Program, Natural Heritage Section. 33 p. [19328]
60. Norland, Jack E.; Marlow, Clayton B. 1984. Use of wooded draws by free-roaming bison. In: Noble, Daniel L.; Winokur, Robert P., eds. Wooded draws: Characteristics and values for the Northern Great Plains: Symposium proceedings; 1984 June 12-13; Rapid City, SD. Great Plains Agricultural Council Publication No. 111. Rapid City, SD: South Dakota School of Mines and Technology, Biology Department: 40-44. [1777]
61. Peden, Donald G. 1976. Botanical composition of bison diets on shortgrass plains. *American Midland Naturalist*. 96(1): 225-229. [24596]
62. Peden, D. G.; Van Dyne, G. M.; Rice, R. W.; Hansen, R. M. 1974. The trophic ecology of *Bison bison* L. on shortgrass plains. *Journal of Applied Ecology*. 11: 489-497. [1861]
63. Reynolds, Hal W. 1986. The Canadian Wildlife Service program to restore wood bison. In: Workshop on endangered species in the Prairie Provinces; [Date unknown]; [Place unknown]. Occasional Paper No. 9. [Edmonton, AB]: Alberta Provincial Natural History Museum: 323-328. [24525]
64. Reynolds, H. W.; Hansen, R. M.; Peden, D. G. 1978. Diets of the Slave River lowland bison herd, Northwest Territories, Canada. *Journal of Wildlife Management*. 42(3): 581-590. [25134]
65. Reynolds, H. W.; Hawley, A. W. L. 1987. Bison ecology in relation to agricultural development in the Slave River lowlands, Northwest Territories. Occasional Paper No. 63. Ottawa, ON: Canadian Wildlife Service. 74 p. [24553]
66. Reynolds, H. W.; Gates, C. C. 1991. Managing wood bison: a once endangered species. In: Renecker, Lyle A.; Hudson, Robert J., eds. Wildlife production: conservation and sustainable development. Misc. Publ. 91-6. Fairbanks, AK: Alaska University, Agricultural and Forestry Experiment Station: 363-371. [24522]
67. Risser, Paul G. 1990. Landscape processes and the vegetation of the North American grassland. In: Collins, Scott L.; Wallace, Linda L., eds. Fire in North American tallgrass prairies. Norman, OK: University of Oklahoma Press: 133-146. [14199]
68. Samoil, J. 1992. Fire on the range: burning for bison habitat. *Timberlines*. 1: 4. [18185]
69. Sharps, Jon C.; Uresk, Daniel W. 1990. Ecological review of black-tailed prairie dogs and associated species in western South Dakota. *Great Basin Naturalist*. 50(4): 339-344. [14895]
70. Shaw, James H.; Carter, Tracy S. 1990. Bison movements in relation to fire and seasonality. *Wildlife Society Bulletin*. 18(4): 426-430. [14777]
71. Shiflet, Thomas N., ed. 1994. Rangeland cover types of the United States. Denver, CO: Society for Range Management. 152 p. [23362]
72. Short, Henry L.; McCulloch, Clay Y. 1977. Managing pinyon-juniper ranges for wildlife. Gen. Tech. Rep. RM-47. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station. 10 p. [2137]
73. Singer, Francis. 1991. Ungulate populations models and carrying capacity: Predicting effects of fires, new winter range, and proposed wolf reintroduction. *Park Science*. 11(1): 8-9. [13928]
74. Stelfox, John G.; Taber, Richard D. 1969. Big game in the northern Rocky Mountain coniferous forest. In: Taber, Richard D, ed. Coniferous forests of the northern Rocky Mountains: Proceedings of the 1968 symposium; 1968 September 17-20; Missoula, MT. Missoula, MT: University of Montana Foundation, Center for Natural Resources: 197-222. [7546]
75. Steuter, Allen A. 1986. Fire behavior and standing crop characteristics on repeated seasonal burns--northern mixed prairie. In: Koonce, Andrea L., ed. Prescribed burning in the Midwest: state-of-the-art: Proceedings of a symposium; 1986 March 3-6; Stevens Point, WI. Stevens Point, WI: University of Wisconsin, College of Natural Resources, Fire Science Center: 54-59. [16269]

76. Steuter, Allen A. 1988. Restoring a mixed prairie process: the fire-bison grazing interaction. In: Bulletin of the Ecological Society of America. [Supplement]. 69(2): 308. [Abstract]. [24520]
77. The Network of Natural Heritage Programs and Conservation Data Centers and The Nature Conservancy. 1994. Element distribution - North America, vertebrates. Arlington, VA: The Nature Conservancy, Central Conservation Databases. 31 p. [23374]
78. Thomas, Gregory. 1977. Fire and the fur trade: The Saskatchewan District: 1790-1840. The Beaver. Autumn: 32-39. [14773]
79. Tieszen, Larry L.; Ode, David J.; Barnes, Paul W.; Bultsma, Paul M. 1983. Seasonal variation in C3 and C4 biomass at the Ordway Prairie and selectivity by bison and cattle. In: Kucera, Clair L., ed. Proceedings, 7th North American prairie conference; 1980 August 4-6; Springfield, MO. Columbia, MO: University of Missouri: 165-174. [3218]
80. U.S. Department of the Interior, Fish and Wildlife Service. 2016. Endangered Species Program, [Online]. Available: <http://www.fws.gov/endangered/>. [86564]
81. Uno, Gordon E. 1987. Buffalo wallows: ephemeral pools in the Great Plains. In: American Journal of Botany. 74(5): 663. [Abstract]. [24526]
82. Urness, Philip J. 1986. Value of crested wheatgrass for big game. In: Johnson, Kendall L., ed. Crested wheatgrass: Its values, problems and myths: Symposium proceedings; 1983 October 3-7; Logan, UT. Logan, UT: Utah State University: 147-153. [2406]
83. Van Gelden, Richard George. 1982. Mammals of the National Parks. Baltimore, MD: Johns Hopkins University Press. 310 p. [20893]
84. Van Vuren, Dirk. 1984. Summer diets of bison and cattle in southern Utah. Journal of Range Management. 37(3): 260-261. [24531]
85. Van Vuren, Dirk. 1987. Bison west of the Rocky Mountains: an alternative explanation. Northwest Science. 61(2): 65-69. [52]
86. Vinton, M. A.; Hartnett, D. C. 1992. Effects of bison grazing on *Andropogon gerardii* and *Panicum virgatum* in burned and unburned tallgrass prairie. Oecologia. 90(3): 374-382. [19436]
87. Vinton, Mary Ann; Harnett, David C.; Finck, Elmer J.; Briggs, John M. 1993. Interactive effects of fire, bison (Bison bison) grazing and plant community composition in tallgrass prairie. American Midland Naturalist. 129: 10-18. [20182]
88. Waggoner, Van; Hinkes, Mike. 1986. Summer and fall browse utilization by an Alaskan bison herd. Journal of Wildlife Management. 50(2): 322-324. [24530]
89. Wasser, C. H. 1977. Bison induced stresses in Colorado National Monument. In: Final Report. National Park Service Contract PX 120060617. 120 p. [24555]
90. Wydeven, Adrian P.; Dahlgren, Robert B. 1985. Ungulate habitat relationships in Wind Cave National Park. Journal of Wildlife Management. 49(3): 805-813. [57]
91. Zarn, Mark. 1977. Ecological characteristics of pinyon-juniper woodlands on the Colorado Plateau: A literature survey. Tech. Note T/N 310. Denver, CO: U.S. Department of the Interior, Bureau of Land Management, Denver Service Center. 183 p. [2689]
92. Chan, Susan. 1993. Bison reintroduced to Oklahoma tall grass prairie preserve. Animal Keepers' Forum. 20(11): 379-380. [24532]
93. Pfeiffer, Kent E.; Hartnett, David C. 1995. Bison selectivity and grazing response of little bluestem in tallgrass prairie. Journal of Range Management. 48(1): 26-31. [24533]
94. Meagher, Mary M. 1978. Bison. In: Schmidt, J. L.; Gilbert, D. L., eds. Big game of North America: ecology and management. Harrisburg, PA: Stackpole Books: 123-133. [24556]
95. Koehler, John T. 1992. Prescribed burning: a wildfire prevention tool?. Fire Management Notes. 53-54(4): 9-13. [24598]
96. Pfeiffer, Kent E.; Steuter, Allen A. 1994. Preliminary response of Sandhills prairie to fire and bison grazing. Journal of Range Management. 47(5): 395-397. [23954]

97. Wright, Henry A.; Bailey, Arthur W. 1982. Fire ecology: United States and southern Canada. New York: John Wiley & Sons. 501 p. [2620]
 98. Committee on the Status of Endangered Wildlife in Canada. 1992. Canadian species at risk. Ottawa, ON. 10 p. [26183]
 99. Baker, Robert J.; Bradley, Lisa C.; Bradley, Robert D.; Dragoo, Jerry W.; Engstrom, Mark D.; Hoffmann, Robert S.; Jones, Cheri A.; Reid, Fiona; Rice, Dale W.; Jones, Clyde. 2003. Revised checklist of North American mammals north of Mexico, 2003. Occasional Papers No. 229. Lubbock, TX: Museum of Texas Tech University. 23 p. [50946]
-