

Projected Impacts of Climate Change on Forests of the Dolores Watershed

A Practical Management Strategy

Jim Worrall

Suzanne Marchetti

US Forest Service
Rocky Mountain Region
Gunnison, Colorado

Jerry Rehfeldt

Nick Crookston

US Forest Service
Rocky Mtn. Res. Station
Moscow, Idaho



Outline

1. Impacts of turn-of-the-century drought
2. Bioclimate models and projections
3. Strategy for applying projections to forest management



Impacts to Forests


TURN-OF-THE-CENTURY DROUGHT




Southern Rockies

Drying trend
since mid '80s

Record drought
2001-2003



The turn-of-the-century drought in the context of the past




Conclusion:

Most recent drought that severe
West-wide occurred >800 yr ago.

Piñon mortality SW CO

Dominique Bachelet, Data Basin



2.5 million acres in 4-corner states

Mountain pine beetle



3.4 million acres in Colorado

2002

927,000 acres burned
in Colorado

SAD



1.2 million acres in Colorado

Spruce beetle



1.7 million acres in Colorado

Future Droughts


- By the 2030's, moisture conditions during the turn-of-the-century drought "... will become the new norm in western North America." (Schwalm et al. 2012, Nature Geosciences)
- Later in the century, conditions may be worse than during TOC.
- How can we anticipate forest impacts so that we can manage appropriately?




Characterize climate suitable for a species


BIOCLIMATE MODELS







Full model window, southwestern Colorado



Telluride

Rico

Dolores

Cortez

Mancos

Dolores Watershed

Bioclimate Models

- Train on real point locations
 - Known presence/absence of species
 - Known historic/reference climate
 - Known slope and aspect
- Learn the combinations and interactions of variables associated with presence
- Given a topoclimate, can then predict the likelihood of presence



How they were built


- Training data
 - 854,000 points of known presence/absence
 - Mostly spatial veg data like FSVeg and FIA plots
 - Long-term historical climate at each point
 - Use reference climate 1961-1990
 - Also slope and aspect
- “Random forests” algorithm in R
 - “forests” of many decision “trees”, using random subsets of training data



Output

- Proportion of decision trees that 'vote' for the topoclimate being suitable.
- If we run each pixel on a map through the model, using each pixel's climate, slope and aspect . . .






PIPO
Proportion
of votes




0-10




20-30



40-50



60-70



80-90



10-20



30-40




50-60



70-80




90-100




ABLA occurrence in shapefiles (black) vs. reference period votes ≥ 0.5 (blue overlay)

Subalpine fir



PIEN occurrence in shapefiles (black) vs. reference period votes ≥ 0.5 (blue overlay)


Engelmann spruce




QUGA occurrence in shapefiles (black) vs. reference period votes ≥ 0.5 (blue overlay)


Gambel oak

Process for projecting into the future





Ponderosa pine 2060



Applying bioclimate models


A FOREST MANAGEMENT STRATEGY



Project Change

1. Make grids of suitability based on reference and future climates.
2. Use difference in model output between reference and future periods to classify change zones.
 - a) Model output (“votes”) are 0 (no chance) to 1 (very likely)

Change classes	Reference period votes	2056-2065 votes
LOST	≥ 0.5	< 0.3
THREATENED	≥ 0.5	0.3-0.5
PERSISTENT	≥ 0.5	≥ 0.5
EMERGENT	< 0.5	≥ 0.5



CGH Change
Zones for ABCO



Lost



Threatened




Persistent



Emergent

White fir



CGH Change
Zones for ABLA



Lost



Threatened




Persistent



Emergent

Subalpine fir



CGH Change
Zones for JUOS



Lost



Threatened




Persistent



Emergent

Utah juniper



CGH Change
Zones for JUSC2


Lost

Threatened

Persistent

Emergent

Rocky Mountain juniper



CGH Change
Zones for PIED


Lost

Threatened

Persistent

Emergent

Piñon



CGH Change
Zones for PIEN



Lost



Threatened




Persistent



Emergent


Engelmann spruce



CGH Change Zones for PIPO

	Lost		Threatened		Persistent		Emergent
---	------	---	------------	---	------------	---	----------

Ponderosa pine



CGH Change
Zones for PIPU


Lost

Threatened

Persistent

Emergent

Blue spruce



CGH Change
Zones for POTR5


Lost

Threatened

Persistent

Emergent


Aspen



CGH Change Zones for PSME

 Lost	 Threatened	 Persistent	 Emergent
--	--	--	--

Douglas-fir



CGH Change
Zones for QUGA



Lost



Threatened




Persistent



Emergent


Gambel oak



CCSM4_rcp45 Change Zones for ABLA

■ Lost	■ Threatened	■ Persistent	■ Emergent
---	--	---	--

**Favorable
climate**







HadGEM2ES_rcp85 Change Zones for ABLA

■ Lost	■ Threatened	■ Persistent	■ Emergent
---	--	---	--

**Subalpine fir
Engelmann spruce**

**Unfavorable
climate**





CCSM4_rcp45 Change Zones for POTR5

■ Lost	■ Threatened	■ Persistent	■ Emergent
---	--	--	--



HadGEM2ES_rcp85 Change Zones for POTR5

■ Lost	■ Threatened	■ Persistent	■ Emergent
---	--	--	--

**Favorable
climate**

**Aspen
Douglas-fir**

**Unfavorable
climate**



Potential Applications


- Having a good estimate of where species will and will not be suited in the future, we can:
 1. On a given site, apply treatments most appropriate for the future.
 2. Focus efforts where they have the greatest likelihood of long-term success.
 3. Identify sites and strategies to protect special species.
- Planning
 - Forest
 - Landscape assessment
 - Projects



Project area pre-determined

APPLY TREATMENTS APPROPRIATE FOR PROJECTED FUTURE





Telluride


Rico

Dolores

Cortez

Mancos

Dolores Watershed



CGH Change
Zones for PIPO


Lost

Threatened

Persistent

Emergent

Ponderosa pine



CGH Change
Zones for PIED


Lost

Threatened

Persistent

Emergent

Piñon



CGH Change
Zones for JUOS



Lost



Threatened



Persistent




Emergent

Utah juniper

Where should we manage spruce for timber?

**IDENTIFY AREAS ON THE LANDSCAPE
WHERE TREATMENTS HAVE THE
GREATEST CHANCE OF LONG-TERM
SUCCESS**





CGH Change
Zones for PIEN

Lost

Threatened

Persistent

Emergent

Englemann spruce

Where to manage spruce?

PERSISTENT zone

- Manage as normal.

THREATENED zone

- Manage as 2nd priority, but
- Focus more on resilience.

LOST zone

- Avoid investing in the future of spruce.
- Treat for short-term benefits and to
- Begin pushing cover-type to the future.




What about the EMERGENT zone?

ASSISTED AND FACILITATED MIGRATION



- Assisted migration:
 - Movement of species and populations to facilitate range expansion in direct management response to climate change
- Facilitated migration
 - Enhancing opportunities for self-migration by favoring seed production and dispersal in current habitat and receptive seedbeds in nearby emergent habitat.






Douglas-fir
emergent habitat

CGH Change Zones for PSME

 Lost	 Threatened	 Persistent	 Emergent
--	--	--	--



Aspen emergent habitat

CGH Change Zones for POTR5

Lost


Threatened

Persistent

Emergent

Facilitating Migration

PERSISTENT OR
THREATENED



EMERGENT




Promote existing
(female) aspen near
Emergent areas to
enhance seed production

Favor disturbances
(esp. fire) in newly
suitable areas to
facilitate aspen
establishment

Climate refugia

PROTECTING SPECIES AT RISK






Persistent - potential refugia for spruce.
Enhance regeneration, protect from severe disturbance

CGH Change Zones for PIEN

 Lost	 Threatened	 Persistent	 Emergent
---	---	---	---


Engelmann spruce




CGH Change Zones for PIPO



 Lost	 Threatened	 Persistent	 Emergent
--	--	--	--

Ponderosa pine



Ponderosa pine 2060





Conclusion

- Bioclimate models can be used to:
 - Design treatments most appropriate for the future.
 - Focus resources where there is the greatest likelihood of long-term success.
 - Identify best approaches and sites to manage species at risk.



“If you don’t plan for the future,
you’re planning for the past”