

Post-Wildfire Logging Hinders Regeneration and Increases Fire Risk

D. C. Donato,^{1*} J. B. Fontaine,² J. L. Campbell,¹ W. D. Robinson,² J. B. Kauffman,³ B. E. Law¹

¹Department of Forest Science, Oregon State University, Corvallis, OR 97331, USA. ²Department of Fisheries and Wildlife, Oregon State University, Corvallis, OR 97331, USA. ³Institute of Pacific Islands Forestry, USDA Forest Service, Pacific Southwest Research Station, 60 Nowelo Street, Hilo, HI 96720, USA.

*To whom correspondence should be addressed. E-mail: dan.donato@oregonstate.edu

Recent increases in wildfire activity in the United States have intensified controversies surrounding the management of public forests after large fires (1). The view that postfire (salvage) logging diminishes fire risk via fuel reduction, and that forests will not adequately regenerate without intervention that includes logging and planting, is widely held and commonly cited (2, 3). An alternative view maintains that postfire logging is detrimental to long-term forest development, wildlife habitat and other ecosystem functions (1). Scientific data directly informing this debate are lacking.

Here we present data from a study of early conifer regeneration and fuel loads following the 2002 Biscuit Fire, Oregon, USA, with and without postfire logging. Because of the fire's size (~200,000 hectares), historic reforestation difficulties in the region (4), and an ambitious postfire logging proposal, the Biscuit Fire has become a national icon of postfire management issues. We used a spatially nested design of logged and unlogged plots replicated across the fire area and sampled before (2004) and after (2005) logging (5).

Natural conifer regeneration on sites that experienced high-severity fire was variable but generally abundant, with a median stocking density of 767 seedlings per hectare, primarily of Douglas-fir (*Pseudotsuga menziesii*) (Fig. 1A). Such density exceeds regional standards for fully stocked sites, suggesting that active reforestation efforts may be unnecessary. Postfire logging subsequently reduced regeneration by 71%, to 224 seedlings per hectare (Fig. 1A), due to soil disturbance and physical burial by woody material during logging operations. Thus, if postfire logging is conducted in part to facilitate reforestation, replanting could result in no net gain in early conifer establishment.

Postfire logging significantly increased both fine and coarse downed woody fuel loads (Fig. 1B). This pulse was comprised of unmerchantable material (e.g., branches), and far exceeded expectations for postfire logging-generated fuel loads (5, 6). In terms of short-term fire risk, a reburn in logged stands would likely exhibit elevated rates of fire spread, fireline intensity and soil heating impacts (7).

Postfire logging alone was notably incongruent with fuel reduction goals. Fuel reduction treatments (prescribed burning or mechanical removal) are frequently intended following postfire logging, including in the Biscuit plan, but resources are often not allocated to complete them (8). Our study underscores that, after logging, mitigation of short-term fire risk is not possible without subsequent fuel reduction treatments. However, implementing these treatments is also problematic. Mechanical removal is generally precluded by its expense, leaving prescribed burning as the most feasible method. This will result in additional seedling mortality and potentially severe soil impacts due to long duration combustion of logging-generated fuel loads. Therefore, the lowest fire risk strategy may be to leave dead trees standing as long as possible (where they are less available to surface flames), allowing for aerial decay and slow, episodic input to surface fuel loads over decades.

Our data show that postfire logging, by removing naturally seeded conifers and increasing surface fuel loads, can be counterproductive to goals of forest regeneration and fuel reduction. In addition, forest regeneration is not necessarily in crisis across all burned forest landscapes. The results presented here suggest that postfire logging may conflict with ecosystem recovery goals.

References and Notes

1. D. B. Lindenmayer *et al.*, *Science* 303, 1303 (2004).
2. U.S. House Committee on Resources, *Forest Recovery Bill Hearing Press Release* (November 9, 2005).
3. J. Sessions, P. Bettinger, R. Buckman, M. Newton, J. Hamann, *J. For.* 102, 38 (2004).
4. S. D. Tesch, S. D. Hobbs, *W. J. Appl. For.* 4, 89 (1989).
5. Materials and methods are available as supporting material on *Science Online*.
6. Timber decay associated with delays in postfire logging was anticipated to exacerbate the observed pulse of surface fuel. However, our data indicate that delay was responsible for ~10% of woody fuel present after logging.

7. J. K. Agee, *Fire Ecology of Pacific Northwest Forests* (Island Press, Washington, DC, 1993).
8. R. W. Gorte, "Forest Fires and Forest Health" *Congressional Research Service* (Publication 95-511, 1996).
9. This work was supported by the Joint Fire Science Program and DOE grant DE-FG02-04ER63917. We thank field technicians and the Siskiyou National Forest.

Supporting Online Material

www.sciencemag.org/cgi/content/full/1122855/DC1

Materials and Methods

SOM text


References and Notes

21 November 2005; accepted 21 December 2005

Published online 5 January 2006; 10.1126/science.1122855

Include this information when citing this paper.

Fig. 1. Natural conifer regeneration (**A**) and surface woody fuel loads (**B**) before and after postfire logging of the Biscuit Fire, Oregon, USA. (A) Regeneration was abundant following fire. Postfire logging significantly reduced seedling densities ($P < 0.01$, Wilcoxon signed rank test) from 767 seedlings ha^{-1} to 224 seedlings ha^{-1} . (B) Postfire logging significantly increased downed fine ($P < 0.01$) and coarse ($P < 0.05$) woody fuel loads (Mg ha^{-1}) relative to burn-only by Wilcoxon signed rank test. To provide context, fuel data from unburned stands are shown as reference for pre-fire conditions (fuel loads in burn-logged stands were at or well above pre-fire levels). Graphs of seedling densities and fine (≤ 7.62) and coarse (> 7.62) surface woody fuels are medians \pm SE; $n = 8$ stands for no burn, $n = 9$ for burn-only and burn-logged (5).

A**B**