

Wildfire-Driven Forest Conversion in Western North American Landscapes

JONATHAN D. COOP, SEAN A. PARKS, CAMILLE S. STEVENS-RUMANN, SHELLEY D. CRAUSBAY, PHILIP E. HIGUERA, MATTHEW D. HURTEAU, ALAN TEPLEY, ELLEN WHITMAN, TIMOTHY ASSAL, BRANDON M. COLLINS, KIMBERLEY T. DAVIS, SOLOMON DOBROWSKI, DONALD A. FALK, PAULA J. FORNWALT, PETER Z. FULÉ, BRIAN J. HARVEY, VAN R. KANE, CAITLIN E. LITTLEFIELD, ELLIS Q. MARGOLIS, MALCOLM NORTH, MARC-ANDRÉ PARISIEN, SUSAN PRICHARD, AND KYLE C. RODMAN

Changing disturbance regimes and climate can overcome forest ecosystem resilience. Following high-severity fire, forest recovery may be compromised by lack of tree seed sources, warmer and drier postfire climate, or short-interval reburning. A potential outcome of the loss of resilience is the conversion of the prefire forest to a different forest type or nonforest vegetation. Conversion implies major, extensive, and enduring changes in dominant species, life forms, or functions, with impacts on ecosystem services. In the present article, we synthesize a growing body of evidence of fire-driven conversion and our understanding of its causes across western North America. We assess our capacity to predict conversion and highlight important uncertainties. Increasing forest vulnerability to changing fire activity and climate compels shifts in management approaches, and we propose key themes for applied research coproduced by scientists and managers to support decision-making in an era when the prefire forest may not return.

Keywords: climate change, ecological transformation, high-severity fire, tree regeneration, tree seedlings, stand-replacing fire, wildfire, vegetation type conversion.

When a forest burns in a wildfire, should we expect it to return as it was before? If the fire burns at low severity—such as a surface fire that does little damage to large, thick-barked trees—the forest character may remain essentially unchanged. However, following a high-severity fire that kills most trees, the near-term postfire environment may include some combination of dead snags, non-forest vegetation (e.g., grasses, resprouting shrubs) or tree seedlings. Given similar climate and disturbance regimes, well-understood successional processes are expected to lead, over time, to the recovery of prefire forest composition and structure. But under changing disturbance regimes and climate, can we still count on the return of the forest as it was before fire?

Across western North America—as in many other locations—the last several decades have been marked by increasing forest fire activity (figure 1; Westerling 2016, Hanes et al. 2018). These changes are coupled with the direct effects of rising temperature and evaporative demand (figure 1) on postfire vegetation dynamics. Western North

American forests have long been shaped by wildfire (box 1), and most tree species exhibit fire-adaptive traits (Rowe and Scotter 1973, Baker 2009, Pausas and Keeley 2014). These include survival mechanisms that confer individual-level resistance to fire-caused mortality, and population-level mechanisms that promote postfire regeneration; collectively, these processes confer ecological *resilience*—the capacity to absorb disturbance and recover toward prior composition, structure, and function (Gunderson 2000, Reyer et al. 2015, Ghazoul et al. 2015, Falk et al. 2019). However, specific traits are adaptive only within particular fire regimes, and altered fire regimes can render formerly well-adapted species vulnerable (Keeley et al. 1999, Brown and Johnstone 2012). Fire size, frequency, or intensity outside the ranges to which dominant species are adapted can overcome both resistance and recovery mechanisms of forest ecosystems (Johnstone et al. 2016). Climate warming is also creating local post-fire environmental conditions outside of the regeneration niche of dominant tree species (Stevens-Rumann et al. 2018, Davis et al. 2019a). As such, heightened vulnerability

BioScience XX: 1–15. © The Author(s) 2020. Published by Oxford University Press on behalf of the American Institute of Biological Sciences. This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com.
doi:10.1093/biosci/biaa061




Figure 1. Trends in annual log-scale area burned and mean annual Z-scores of climatic moisture deficit (CMD), and mean July temperature (Tmax) in forested western North America (1984–2017). Z-scores are based on a 1981–2010 reference period. The annual area burned and the number of fires have significantly increased in all regions except the Mediterranean and Western Boreal. The burn area data are from the Monitoring Trends in Burn Severity for American fires and Canadian National Fire Database for Canadian fires (with a threshold at the same minimum fire size of at least 400 hectares), the forested area is derived from 2001 and 2018 MODIS Land Cover product, the climate data are from TerraClimate, and the ecoregions are adapted from the Commission for Environmental Cooperation North American Terrestrial Ecoregions.

to the combined impacts of changing fire regimes and climate raises concerns that, for some forest types, resilience to fire may be increasingly compromised (for a review, see Johnstone et al. 2016, Davis et al. 2018, Falk et al. 2019, Hessburg et al. 2019).

A direct outcome of declining forest resilience to fire is an increase in the proportion of a burned landscape with post-fire vegetation that diverges considerably from its prefire state. Where altered postfire vegetation is spatially extensive and temporally enduring (i.e., longer than the known historical postfire recovery time), such changes may be referred to as *conversion*. This terminology (e.g., forest conversion, site conversion, and type conversion), although lacking a formal definition, is increasingly used by natural resource managers and researchers alike. Conversion implies major, extensive, and enduring changes in dominant species, life forms, or functions (e.g., shifts from one forest type to another, or from a forest to nonforest vegetation). Conversion may be considered inclusive of a suite of other terms used to portray major and abrupt changes to ecological systems, including reorganization (Falk 2017), regime shift (Anderson et al.

2009), state shift (Barnosky et al. 2012), state transition (Stringham et al. 2003), critical transition (Scheffer et al. 2009), and transformation (Folke et al. 2010).

In the present article, our focus is *fire-driven forest conversion*, which can be viewed as a two-step process. First, major vegetation shifts are initiated by high-severity fire that removes large areas of mature forest from the landscape. Second, recovery mechanisms are inhibited by the absence of seed sources, short-interval reburning, or post-fire climate and other environmental conditions unfavorable to seedling recruitment. Return of the prefire forest is protracted or prevented, and fire-initiated changes persist, although subject to the influences of subsequent fire–vegetation feedbacks. The direction of conversion is dependent on the specific factors that lead to loss of system resilience. In some cases, for example, fire may catalyze vegetation change expected under future climate (e.g., conversion to grassland at trailing edge lower forest ecotones; Donato et al. 2016); in other cases, change may be linked to shifts in subsequent fire regime (e.g., conversion to highly flammable chaparral; Tepley et al. 2017).

Box 1. Forests and fire regimes in western North America.

Temperate and boreal forests in western North America are dominated mostly by conifer species (including fir, *Abies*; juniper, *Juniperus*; spruce, *Picea*; pine, *Pinus*; Douglas fir, *Pseudotsuga*; and hemlock, *Tsuga* spp.), but with some important components of broadleaf species including trembling aspen (*Populus tremuloides*), birch (*Betula* spp.), maple (*Acer* spp.), and oak (*Quercus* spp.). Forest types vary predictably along moisture and temperature gradients, which interact with vegetation to shape fire regimes—the characteristic frequency and severity of wildfire over decades to centuries—via differing constraints on fire activity (Schoennagel et al. 2004, Baker 2009, Littell et al. 2018). Generally, forests in warm and dry settings (e.g., ponderosa pine (*Pinus ponderosa*) and dry mixed-conifer forests) experience climatic conditions suitable for burning on subdecadal to decadal frequencies. Productivity in these systems is moderate to low, and historically, fuel quantity and continuity were reduced by frequent, low-intensity fires. Historical fire regimes in these systems are considered to be more fuel limited than climate limited. Consequently, although fire may have been frequent, severity was typically low to moderate. Systems characterized by such a regime may be termed *frequent-fire* forest types. Conversely, in cooler and wetter settings (e.g., boreal, subalpine, and mesic Pacific Northwest forests), fire activity is considered climate limited: Biomass is usually sufficient to support high-intensity fire, but conditions required to dry out fuels occur less often. Conditions suitable for burning occur on multidecadal or multicentury frequencies. However, under warm, dry, and windy conditions, these forests can support crown fires, resulting in large patches of tree mortality (e.g., Turner and Romme 1994). Such systems may be termed *infrequent-fire* forest types. Frequent- and infrequent-fire forests represent the extremes of a continuum, between which many forests share some characteristics of both.




Figure 2. Hypothetical ecological outcomes of fire. The x-axis represents time since the initial fire; the y-axis represents ecological (e.g., compositional) dissimilarity relative to the prefire state. A low value indicates little change, and a high value indicates substantial change, such as a shift from dominance by one species or functional group to another. A forest stand with high resistance may be essentially unchanged following a fire—for example, when a surface fire burns through a ponderosa pine (*Pinus ponderosa*) forest. When resistance is overcome, a resilient system may change substantially, but still return toward its predisturbance state over time—for example, tree regeneration from seed following high-severity fire in a high-elevation lodgepole pine (*Pinus contorta*) forest. When resilience processes are overwhelmed, recovery toward a predisturbance state may be severely protracted or entirely precluded. Conversion refers to this condition. However, studies of recent postfire dynamics are limited to only a few decades postfire, represented by the solid lines; the time scales of these processes are highly uncertain. The dashed lines represent possible but uncertain future trajectories that cannot be known because of changes in climate, fire regimes, and society.

Fire-driven conversion represents one potential outcome on a continuum of postfire vegetation dynamics (figure 2). Climate variability and disturbance have interacted over millennia to drive ecosystem change (Nolan et al. 2018). In fact, some contemporary vegetation patterns still

bear the imprint of long-past conversion events, where succession toward the prefire state has been prevented (e.g., meadows generated by ancient fire and climate change; Lynch 1998) or drawn out over centuries (e.g., protracted aspen to conifer succession following high-severity fire; Margolis et al. 2007). Given the relatively short periods (years to decades) of recent postfire observations, the ultimate duration of contemporary changes cannot be known—a fundamental uncertainty in our understanding. However, accumulating evidence suggests that long-term conversions are increasingly taking place across a range of western North American forest types, and that we are likely to face substantially altered ecosystems on timescales exceeding management planning horizons and human lifespans.

A growing body of observations, empirical work, and understanding of causal processes calls for a synthesis of fire-driven forest conversion. In the present article, we begin by cataloguing the mechanisms that can generate and maintain conversions of forested landscapes across western North America. In compiling these mechanisms within the framework of conversion, we build

on earlier reviews exploring and emphasizing key components of forest resilience and vulnerability (Johnstone et al. 2016, Davis et al. 2018, Falk et al. 2019, Hessburg et al. 2019, Stevens-Rumann and Morgan 2019), illustrating recent observations of their operation, considering interactions




Figure 3. Processes that may give rise to fire-driven forest conversion. (a) Conversion is initiated by processes that result in extensive areas of adult tree mortality (the solid arrows; red represents fire, and yellow represents climate). (b) Conversion is maintained by processes that impede regeneration of prefire tree species (dashed red and yellow arrows) and protract vegetation change temporally. (c) The duration of forest conversion may be further influenced by positive and negative fire-vegetation feedbacks (dashed purple arrows), which respectively promote or inhibit additional burning.

and feedback processes, and highlighting their consequences and implications on the ground. Next, we draw from this mechanistic understanding to assess our capacity to predict future fire-driven forest conversions. Throughout, we identify key uncertainties. Finally, we outline a management framework for navigating conversion, in which we propose four themes for applied and coproduced research to better inform decision-making in a time of certain change, but of uncertain rate, magnitude, and extent.

Mechanisms of forest conversion

Fire-driven conversion results when forest resilience is overcome, inhibited, or otherwise absent. In what follows, we consider how increasing fire activity and warming and drought can break down resistance and recovery processes through increased tree mortality (prior to and as a result of fire; figure 3a) and reduced postfire tree regeneration (figure 3b). Subsequently, the duration and extent of conversion may be modulated by fire-vegetation feedbacks (figure 3c).

Changing fire regimes. Fire that kills most or all trees is a requisite first step toward fire-driven conversion (figure 3a). Fire regimes across western North America have undergone profound changes in the modern era. In most of this region, recent trends of increasing annual area burned, the number of fires, and the average fire size have been observed (figure 1; Dennison et al. 2014, Westerling 2016, Hanes et al. 2018). In addition, the proportion of area burned at high-severity has also increased in some ecoregions (Miller et al. 2009, Harvey et al. 2016b, Singleton et al. 2019). These increases are driven wholly or in part by anthropogenic climate change (Abatzoglou and Williams 2016), increasing human ignitions (Balch et al. 2017), and in formerly frequent-fire forest types, fuel accumulation due to fire exclusion (Steel et al. 2015).

Expanding annual area burned at high severity increases the landscape fraction of early seral postfire vegetation, but long-term conversion is contingent on impeded forest recovery processes, which may also be imparted by increasing fire activity. In some cases a single large and severe fire can effectively lead to conversion through the elimination of biological legacies vital to recovery (Turner et al. 1998, Johnstone et al. 2016). Where forests are composed of obligate seeding, nonserotinous tree species (e.g., ponderosa pine), large high-severity patches can limit postfire establishment because distances to live tree seed sources exceed characteristic seed dispersal distances (figures 3b and 4). These constraints may be further enhanced by the loss of climate buffering by the forest canopy (Davis et al. 2019b) and the development of competing vegetation (Stevens-Rumann et al. 2018). Where high-severity patches are exceptionally large, recovery may require multiple generations of tree colonization, maturation, and dispersal (Haire and McGarigal 2010, Chambers et al. 2016, Harvey et al. 2016c). Although the prefire forest type might eventually return within large patches created by high-severity fire, delays increase the likelihood of persistent change as seedling establishment becomes increasingly untenable under a warming climate (Liang et al. 2017).

As annual area burned increases, so too does the probability that a fire burns over a recently burned area (i.e., short-interval fires or early seral reburning; Prichard et al. 2017). Where frequent-fire forests have departed from historic fire regimes, intense fires occurring in short succession may surmount fire resistance and postfire recovery. The first fire shifts vegetation from obligate-seeding conifers to resprouting species while also producing abundant dead and down woody debris; this fuels the second fire, which eliminates conifer seedlings and any remaining seed sources, and further expands resilient resprouting




Figure 4. Exceptionally large high-severity patches in a frequent-fire forest type. (a) The postfire landscape of the Hayman fire in Colorado; (b) distribution of distances from high-severity patches to surviving tree seed sources within the burn perimeter. Fifteen years after a fire, Chambers and colleagues (2016) found that sites less than 50 meters from tree seed sources were not recovering toward prefire forest densities, and most of this landscape is now dominated by shrubs and herbs. Photograph: O. Rhoades. The data are from Jonathan D. Coop.




Figure 5. Shortening fire-free intervals lead to a shift from conifer to broadleaf boreal forests. These fire regime changes also lower the stem densities of stands, creating a more open forest type, and, in extreme cases, result in regeneration failure. (a) The relationship between the fire-free interval and the proportion of conifer seedlings (versus broadleaf tree regeneration) in the postfire cohort from burns in the Northwest Territories, Canada. (b) A short-interval reburn (10 years between severe fires) in Wood Buffalo National Park, Canada, showing exposed mineral soil and charred logs. There are no residual *in situ* seed or sucker sources for either conifers or broadleaves at the center of the reburn. Photograph: Ellen Whitman. The data are from Whitman and colleagues (2019).

species (figure 3b; Coop et al. 2016, Coppoletta et al. 2016). Short-interval fires can also undermine mechanisms conferring resilience to high-severity fire (figures 3a and 5). Serotinous or semiserotinous cones allow species such as lodgepole pine (*Pinus contorta*) and black spruce (*Picea mariana*) to maintain canopy seedbanks and disperse seeds

locally following intense fire. With an adequate fire-free interval, forest composition and structure recover (Buma et al. 2013). However, when reburning occurs before tree maturation, postfire seed sources are absent (Keeley et al. 1999, Brown and Johnstone 2012, Turner et al. 2019). In these cases, infrequent-fire forest types are vulnerable to




Figure 6. Shrinking windows of climatic opportunity for postfire regeneration. After 2000, conditions conducive for postfire regeneration in ponderosa pine and Douglas fir (*Pseudotsuga menziesii*) forests became less prevalent (Davis et al. 2019a). (a) Photo of Canyon Ferry burn in Montana, 17 years postfire, where postfire surveys suggest that there has been little to no recruitment at lower elevations. (b) Modeled probability of recruitment (above the 25th percentile of the annual rate) at this site (Davis et al. 2019a). Photograph: Kimberly T. Davis.

transitions from one forest type to another (e.g., from pine to aspen; Hart et al. 2019, Whitman et al. 2019), or to conversion from forest to grassland or shrubland (Brown and Johnstone 2012).

Direct effects of a changing climate. In addition to shaping fire regimes, climate change also contributes directly to forest conversion through effects on pre- and postfire tree population dynamics. Warmer and drier conditions can stress trees and cause mortality in the absence of fire, or predispose trees to fire-induced mortality (figure 3a; van Mantgem et al. 2013, 2018). Severe drought associated with climate change has triggered major tree die-offs via hydraulic failure or carbon starvation, or mediated through insects and pathogens (Anderegg et al. 2015). Such die-offs may hasten conversion by removing potential seed sources and increasing dead fuels.

Where prefire tree species cannot regenerate under contemporary climate, conversion is maintained (figure 3b). Even where sufficient seed sources are available, warmer and drier postfire conditions can lead to tree recruitment failures (Stevens-Rumann et al. 2018, Hansen and Turner 2019), upholding shifts to nonforest vegetation or tree species with different physiological tolerances or regeneration strategies (Hansen et al. 2016). Postfire tree regeneration is highly sensitive to climate, and directional change, as well as intra- and interannual fluctuations, shape the likelihood of regeneration success (Davis et al. 2018). Postfire recruitment pulses in dry forest types are most common in wet years that are more favorable for seedling establishment (Brown and Wu 2005, O'Connor et al. 2017) but are projected to become less frequent (figure 6; Davis et al. 2019a).

Postfire tree regeneration failures are most likely to occur, and conversion most likely to endure, at the lower elevation treeline or trailing edge ecotone (Rother et al. 2015, Donato et al. 2016, Parks et al. 2019). Low-elevation sites with high incoming solar radiation, minimal upslope water subsidies, and low moisture availability experience the greatest stress and the highest tree seedling mortality rates (Simeone et al. 2019). In such settings, high-severity fire is expected to catalyze conversions from forest to nonforest that would be anticipated to occur eventually, but more slowly, under directional climate change.

Though recent studies have documented postfire declines in regeneration when fires are followed by warm and dry conditions, the ultimate duration of these changes cannot be known. Constraints on postfire regeneration associated with warmth and drought are documented across elevations (Harvey et al. 2016c, Stevens-Rumann et al. 2018, Whitman et al. 2019) and in some locations there has been a lack of postfire regeneration many decades postfire (Savage and Mast 2005, Donato et al. 2016, Stevens-Rumann et al. 2018). However, although much attention is paid to evidence of recent tree regeneration failure, less is known about episodic postfire recruitment during periods of high moisture availability (but see Brown and Wu 2005), challenging our ability to confidently predict that the prefire tree assemblages will not return.

Fire-vegetation feedbacks. The potential for changing fire regimes and climate to sustain forest conversion is modulated by feedbacks between fire and vegetation (McKenzie and Littell 2017). Patterns of fire-induced tree mortality and postfire vegetation development influence the probability of




Figure 7. High-severity reburns perpetuate and expand flammable shrub cover at the expense of forest. (a) Recent fire history in the Klamath Mountains, with the inset showing reburns (1984–2018) within and adjacent to the 2002 Biscuit Fire (200,000 hectares; the thick outline). The Biscuit Fire completely reburned a 1987 fire, and much of the Biscuit Fire was burned again by smaller fires in 2013 and 2015, and large fires in 2017 and 2018. (b) The photo depicts an area that burned in 1994 and 2008. The 1994 fire burned at high severity on the left side of the photo, killing nearly all trees and initiating shrub-dominated vegetation. The 2008 fire burned both sides of the photo at high severity, perpetuating the shrub cover on the left, and expanding it farther into the forest on the right (note the difference between the blackened snags with few branches on the left, which were dead before the 2008 fire, compared to the gray snags with intact fine branches on the right, which survived the 1994 fire and were killed in 2008). Photograph: Alan Tepley.

tree regeneration, growth, and survival (Davis et al. 2018). These changes in turn influence future fire probability and effects (Archibald et al. 2018).

A positive feedback is one in which a fire-initiated vegetation shift is reinforced or amplified through ensuing positive effects on fire activity (figure 3c). High-severity fire can produce abundant dead and down fuels and dense vegetation regrowth (Nelson et al. 2016) prone to reburning and perpetuating nonforest cover. Furthermore, high-severity fire may drive shifts from forests toward more flammable, shrub-dominated vegetation types (Tepley et al. 2018). These scenarios represent self-reinforcing feedbacks in which severe fire begets severe fire (figure 7). In low-elevation forests in California, large forested areas were converted to shrubland when they burned in high-severity fires between 2000 and 2010 (Coppoletta et al. 2016). When these patches reburned in 2012, much of the area reburned at high severity, perpetuating the shrub-dominated vegetation (Coppoletta et al. 2016). This pattern of high-severity fire begetting high-severity fire has been seen in multiple areas across the western United States (e.g., Collins et al. 2009, Harvey et al. 2016a). In some cases, the second fire may burn at even higher severity than the first (Turner et al. 2019).

A negative feedback occurs when fire creates conditions that limit future burning, which may foster the return of the prefire forest (figure 3c). Reduced burn potential may arise because of slow fuel buildup after fire (Parks et al. 2018b), or

a shift from relatively flammable conifers to less flammable deciduous species (e.g., shifts from conifer forests to early seral vegetation dominated by aspen or birch in boreal landscapes; Héon et al. 2014, Whitman et al. 2019). Low burn probabilities following fire represent a dampening feedback that may permit tree seedling establishment, growth, and survival. These effects, however, are generally temporary. Furthermore, negative feedbacks may be overridden by severe fire weather (Parks et al. 2018b, Harvey et al. 2016a) and climate change-driven increases in wildfire activity (Hart et al. 2019).

Spatial and temporal variation in fire–vegetation feedbacks has important implications for understanding vulnerability to fire-catalyzed conversion as the climate warms and becomes increasingly conducive to wildfire. Where the feedbacks are positive, relatively small increases in wildfire activity could lead to abrupt and extensive shifts to persistent nonforest cover. In systems with negative feedbacks, the period of low flammability following high-severity fire provides a degree of resistance to conversion by increasing the fire-free period during which forests can recover. Feedbacks may also change over a sequence of disturbances. For example, abundant dead and down fuels produced by a high-severity fire may promote a second high-severity fire (Lydersen et al. 2019). However, the consumption of such fuels during the second fire can lead to major reductions in fuels, reducing the probability of a third high-severity fire.

What does the future hold for western North American forests?

In western North America, fire activity is expected to continue to increase in association with climate change throughout this century (Kitzberger et al. 2017, Abatzoglou et al. 2019). Furthermore, in many locations, postfire climate conditions are likely to become increasingly unfavorable to tree regeneration, even if seed sources are nearby (Kemp et al. 2019, Liang et al. 2017). Given these projected changes in fire and climate, we anticipate that many forest ecosystems will face increasing risk of fire-catalyzed change, although the nature of change will depend on forest type and fire-vegetation feedbacks.

Forecasts of forested area susceptible to fire-driven conversion project varying degrees and rates of change across forest types by mid- to late-twenty-first century. In the Sierra Nevada in California, for example, Liang and colleagues (2017) project that fire and climate change will reduce forest extent by 5.8% (averaged over GCMs and transects) by the year 2100. Within the intermountain western United States, 1.6% to 15.1% (depending on ecoregion) of forest area has been modeled to be at risk of fire-catalyzed conversion to nonforest by mid-twenty-first century (Parks et al. 2019). In the southwestern United States, where extreme fire weather was incorporated into fire severity estimates, a more substantial 30% of forested area may be vulnerable to fire-driven conversion (Parks et al. 2019). Similarly, on the Kaibab Plateau of northern Arizona, 3% to 49% of the landscape (depending on forest type and climate scenario) was predicted to be nonforest by 2090 when fire was included in simulations, compared to only 0% to 0.3% when fire was excluded (Flatley and Fulé 2016). In the Klamath region of northern California and southern Oregon, approximately one third of conifer-dominated forest could transition to shrub- or hardwood-dominated ecosystems by the late-twenty-first century (Serra-Diaz et al. 2018). In the mountains of central Idaho, climate change and increased fire activity are expected to substantially reduce the prevalence of four common conifer species (Campbell and Shinneman 2017). In Alberta, Canada, wildfire could catalyze conversion of about 50% of upland mixed-wood and conifer forests to more climatically suited mosaics of grassland, shrubland, and deciduous woodland by 2100 (Stralberg et al. 2018). As a very broad generalization across western North America, bioclimatic models suggest that forested areas will have climate and fire regimes more suited to drier forest types and nonforest vegetation (Parks et al. 2018a).

Because forest recovery—or lack thereof—following high-severity burning is predicated on regeneration, studies focusing on seedling establishment and survival under future climate also inform estimations of vegetation change. In New Mexico, for example, a substantial reduction in successful ponderosa pine regeneration is expected along the dry, lower-elevation boundary of its range (i.e., the trailing edge; Allen and Breshears 1998, Feddema et al. 2013). Decreases in postfire ponderosa pine and Douglas fir

(*Pseudotsuga menziesii*) seedling densities are also predicted by mid-twenty-first century at sites in Idaho and Montana, with effects being most pronounced at lower elevations (Kemp et al. 2019). If these trailing-edge forests experience high-severity fire, conversion to nonforest is probable. Similarly, more than 50% of the area currently suitable for montane forest in the Klamath region could have minimal postfire conifer regeneration by the late-twenty-first century, even if seed sources are available (Tepley et al. 2017). Concurrence between process-based (Serra-Diaz et al. 2018) and statistical models (Tepley et al. 2017) provides more confidence in the prediction that conversions are highly likely in this system.

We cannot ignore, however, uncertainties that currently hinder our ability to predict where, when, and how widespread conversions may be in coming decades. For example, whereas there is near-universal agreement among global climate models that temperatures will continue to rise this century, projected changes in precipitation at regional to global scales are variable (Knutti and Sedláček 2013), and this may impart cascading uncertainties in predictions of future fire and regeneration. However, any potential increases in precipitation may be insufficient to offset the effect of rising temperatures on fire activity (Flannigan et al. 2016) and declines in snowpack (with implications for soil moisture; Harpold and Molotch 2015). In one field experiment, ponderosa pine and Douglas fir seedlings that received a combination of warming and supplemental watering demonstrated lower rates of survival than did untreated controls (Rother et al. 2015). Elevated water use efficiency by plants resulting from carbon fertilization may also partially buffer seedlings against warming temperatures (Keenan et al. 2013), although the net effect remains uncertain and may vary with species and age (Peñuelas et al. 2011, Anderson-Teixeira et al. 2013). Continued research (e.g., Battipaglia et al. 2013) is needed to clarify the influence of carbon fertilization on tree establishment under projected future climate.

No-analog or novel climatic conditions challenge existing models built on observed interactions and feedbacks among climate, vegetation, and fire, potentially limiting the ability of retrospective studies to accurately project future dynamics. When informed by empirical research, process-based (i.e., mechanistic) simulation models have the potential to overcome some of the limitations imposed by no-analog conditions (Gustafson 2013, Loehman et al. 2020). However, process-based models may be constrained by an incomplete understanding of underlying mechanisms such as propagule production and dispersal, inter- and intraspecific interactions within a postfire community, as well as genetic variation and phenotypic plasticity. Additional research on these topics is also needed to improve projections of future changes in western North American forests and bolster existing demographic frameworks (e.g., Enright et al. 2015, Davis et al. 2018).

Properly incorporating fire-vegetation feedbacks into predictions of future conversion is also challenging. For example,

Box 2. Fire-driven forest conversion in the paleological record.

Paleoecological records offer unique insights into past wildfire-driven vegetation conversions that ground our understanding of contemporary and future change. Numerous paleological records feature vegetation shifts associated with climate change (Nolan et al. 2018), but most paleological records that address fire highlight ecological resilience to wildfires (e.g., Minckley et al. 2012). However, during periods of rapid climate change, the paleological record illustrates how wildfire can catalyze ecological changes that either would have taken centuries to unfold or may not have occurred at all.

In a lowland forest of the Pacific Northwest, for example, high-resolution pollen and charcoal records indicate two major vegetation conversions over the Holocene (i.e., the past 11,700 years) that were catalyzed by individual high-severity wildfires at the local scale (Crausbay et al. 2017). While the regional expansion of these new vegetation types was ultimately driven by millennium-scale climate change, the timing of conversion was determined by fire. More recently during the Medieval Climate Anomaly (c. 1000 years ago), a change toward a century-long period of elevated wildfire activity caused a continuous subalpine forest landscape to shift abruptly to a ribbon forest (i.e., alternating bands of meadow and forests) that persists today (Calder et al. 2019).

Conversion in these examples occurred via the interaction of two processes: high-severity wildfire, which killed adult trees that could have otherwise persisted for decades or centuries even under a changing climate, and rapid, directional climate change, which created unsuitable conditions for regeneration of the dominant tree species. Both processes are currently interacting across western North America. The paleological record offers another line of evidence that enduring forest conversion is a potential outcome of contemporary high-severity fires under a changing climate.

the long-term impact of repeated fires on vegetation and fuels is not well understood across biophysical gradients, and will be conditional on factors such as tree mortality following the initial or second fire, exact time interval between fires, post-fire climate, and dominant species (McKenzie and Littell 2017, Hurteau et al. 2019a). Fortunately, quantifying ecosystem responses to short-interval fires is an extremely active area of research that is filling knowledge gaps (Coop et al. 2016, Coppelletta et al. 2016, Harvey et al. 2016a, Tepley et al. 2017, Collins et al. 2018, Parks et al. 2018b, Lydersen et al. 2019, Turner et al. 2019, Whitman et al. 2019, Buma et al. 2020).

Human activities also complicate our ability to project forest conversion, through both direct and indirect influences on land use and land cover, fire regimes, and postfire vegetation change. Human land use practices, acting in concert with a warming climate, have led to a disequilibrium (Svenning and Sandel 2013) between the existing distribution of forests and current climatic conditions in parts of western North America, setting the stage for rapid fire-catalyzed forest conversions (Serra-Diaz et al. 2018). As one important example, ongoing fire suppression in some regions has resulted in a fire deficit, whereas in other regions, the introduction of nonnative invasive grasses and increased human ignitions have expanded the spatial and temporal fire niche and resulted in a fire surplus (Parks et al. 2015, Balch et al. 2017). Pre- and postfire management can also influence the likelihood of conversion. Large-scale fuel reduction is predicted to reduce fire-induced mortality under future climatic conditions (McCauley et al. 2019), and widespread tree planting could also forestall conversion. Consequently, future patterns in human development (e.g., the expanding wildland–urban interface) and human actions (e.g., pre- and postfire management actions, fire suppression and ignition) are additional factors that could be considered when predicting fire-catalyzed

forest conversions. Though some process-based models already incorporate these dynamics at a coarse spatial scale (Lawrence et al. 2016), their influences at finer scales is an important area for future study.

Resolving these uncertainties and identifying where there is convergence across the growing body of research can improve our confidence in predictions of where and when fire is most likely to drive forest conversion. Nevertheless, with the preponderance of evidence—from the paleoecological record (box 2), present-day observations, *in situ* experiments, and future projections—we can state with confidence that fire-driven conversions will unfold across many forested landscapes as climate change proceeds. However, perhaps the most important question is the most elusive: How should society respond to these conversions?

A framework for supporting management decisions around forest conversion

Given our developing understanding of wildfire-driven forest conversion and a wide range of inherent uncertainties, how might science and policy best support management decisions? Western North American forests support a wide range of ecological and social values and services, ranging from utilitarian and economic (e.g., timber production) to aesthetic and spiritual. Many of these services will be changed by the forest losses and shifts we describe in this article (box 3). Sustaining these values and services has guided management policy in Canada and the United States for over a century. However, in a time of pervasive and intensifying change, the implicit assumption that the future will reflect the past is a questionable basis for land management (Falk 2017). Increasing forest vulnerability to changing fire regimes and climate compels revised management paradigms, strategies, and tactics, with a robust scientific foundation.

Box 3. Consequences of fire-catalyzed forest conversion.

The direct and indirect effects of fire-driven forest conversion are numerous and wide ranging, but will depend largely on the characteristics and dynamics of postconversion vegetation assemblages, an area requiring further research. In conifer-dominated systems of western North America, most recent studies have examined fire-driven conversion toward vegetation dominated by genera of resprouting broadleaf trees (in particular, aspen and birch), shrubs (such as oak and ceanothus, *Ceanothus*), and herbaceous communities (Savage and Mast 2005, Abella and Fornwalt 2015, Stevens et al. 2015, Airey Lauvaux et al. 2016, Coop et al. 2016, Guiterman et al. 2018, Barton and Poulos 2018). Given recent historic conifer increases in some systems, contemporary fire-catalyzed conversion to nonconifer dominated systems may, in some cases, represent a return to conditions similar to the early twentieth century (Hessburg et al. 2019). However, the spatial and temporal scale of patchiness of fire-catalyzed conversion may not mirror the opposing pattern of recent conifer densification or encroachment. In other cases, conversion could be viewed as an adaptive change that creates a new system better suited for warmer climate with more fire activity. Severe fire serves as a filter with warm- and fire-adapted species (e.g., resprouters, annuals, some invasives) succeeding at the expense of fire-sensitive species (Abella and Fornwalt 2015, Stevens et al. 2015). However, with implications for ecosystem function, as well as the provision of ecosystem services, particularly carbon sequestration, any shifts will necessarily have a range of local, regional, and global impacts.

Fire-driven forest conversion can lead to reduced carbon storage, altered hydrologic dynamics, plant- and animal-community turnover, and impacts on a wide range of human social and economic values. Forests are a substantial contributor to climate regulation through the uptake and storage of carbon (Pan et al. 2011), and conversions, particularly to nonforested vegetation types, are generally expected to result in reduced productivity and carbon storage. However, in temperate regions, the effect of forests on Earth's energy balance is also a function of local climatic conditions. In semiarid regions, for example, forest cover decreases albedo and low water availability limits the latent heat flux from evapotranspiration, suggesting that a wildfire-induced state change may yield a net cooling effect (Jackson et al. 2008). Widespread forest loss and the associated changes in albedo and land-surface energy balance scale up to affect the entire climate system, with global consequences. For example, models suggest that loss of forests in western North America could lead to drying and reduced net primary productivity in other parts of the world (Stark et al. 2016). Forest conversions may also affect erosion rates and water quality and quantity by decreasing transpiration and increasing overland flow (Wine et al. 2018). Forest conversion will also necessarily drive complex changes to biotic community composition and diversity. High-severity fire can generate habitat for some species dependent on postfire attributes such as snags (e.g., some woodpeckers; Hutto et al. 2015), but these may ultimately be diminished if long-term forest recovery is compromised. For example, the capacity for landscapes to harbor species that rely on shrubs or meadows may increase, but forest- or old-growth obligate species will be increasingly susceptible (e.g., lichens; Miller et al. 2018). Nonnative species may also benefit from fire-driven conversion (Abella and Fornwalt 2015, Stevens et al. 2015).

Over the past decade, consensus has built around a three-part concept of the universe of potential management responses (Aplet and Cole 2010) expressed in terms of resisting, accepting, or directing change. Resisting wildfire-driven forest conversion means attempting to sustain existing forests by supporting prefire resistance or postfire recovery. Accepting conversion concedes the replacement of extant forests by other vegetation types after fire without intervening, accommodating modified communities and altered ecosystem services. Directing conversion uses management interventions to favor particular postfire outcomes aligned with human values (e.g., Aplet and Cole 2010, McWethy et al. 2019).

Contemporary forest management policies, mandates, and science generally fall within the paradigm of resisting conversion, through on-the-ground tactics such as fuel reduction or tree planting. Given anticipated disturbance trajectories and climate change, science syntheses and critical evaluations of such resistance approaches are needed because of their increasing relevance in mitigating future wildfire severity (Stephens et al. 2013, Prichard et al. 2017) and managing for carbon storage (Hurteau et al. 2019b). Managers seeking to wisely invest resources and strategically resist change need to understand the efficacy and durability

of these resistance strategies in a changing climate. Managers also require new scientific knowledge to inform alternative approaches including accepting or directing conversion, developing a portfolio of new approaches and conducting experimental adaptation, and to even allow and learn from adaptation failures.

Science to support decisions around resisting, accepting, or directing forest conversion is best formed within coproduction models between scientists and managers, where both parties meaningfully engage (e.g., Meadow et al. 2015) and target decision-making processes. Decision-making processes such as the Climate-Smart Conservation Cycle (Stein et al. 2014) provide a framework highlighting how science can support decisions to resist, accept, or direct ecological change. In the present article, we propose four central themes toward an array of coproduced science to support decisions around wildfire-driven forest conversion. These include (1) *characterizing vulnerability to fire-driven conversion*, (2) *providing plausible scenarios of post-fire ecological futures* under shifting climate and fire regimes, (3) *assessing the feasibility of directing or resisting conversion*, and (4) *understanding the social and ecological consequences* of the choice to resist, accept, or direct change.

The first theme, *characterizing vulnerability to fire-driven conversion*, offers crucial support to the initial steps in a decision-making process. There are many opportunities to further develop and synthesize knowledge about the likelihood of wildfire-driven conversion, including mapping and modeling the locations of fire refugia (Krawchuk et al. 2016), trailing-edge forests (Parks et al. 2019), climate futures for fire weather (Wang et al. 2017), and postfire recruitment (Davis et al. 2019a). As was described previously, however, there are inherent uncertainties associated with each of the mechanisms that can lead to conversion. These are compounded by interactions among processes (Temperton et al. 2004), and potentially exacerbated by expected no-analog climates of the twenty-first century and nonstationarity of ecological processes. However, uncertainty need not be a limitation for forward-looking managers to engage in proactive thinking about the general vulnerability of the forests they manage to fire-driven conversion in the near future.

The second theme, *providing plausible postfire ecological futures* under shifting climate and fire regimes, will allow managers to consider the consequences of accepting ecological reorganization. Research on interactions between disturbance and climate-driven species, habitat, and biome range shifts will provide more plausible postfire ecological scenarios under climate change. As with the first theme, there is currently high uncertainty around the characteristics of the ecological communities most likely to replace any particular forest. In addition, the field lacks a commonly accepted means of forecasting ecological scenarios. Before decisions can be made about resisting, accepting, or directing conversion, land managers will require some degree of clarity, or ability to incorporate scenarios, about the likelihood and probable character of forest conversion.

The third theme, *assessing the feasibility of resisting or directing conversion*, is a call to expand our paradigm outside the traditional narrow focus on ensuring resilience of existing forest communities (Falk et al. 2019). Currently, a large body of work supports tactics to resist conversion, although these pertain primarily to frequent-fire forest types. Well-established fuel reduction techniques emphasize the retention of larger-diameter trees with thick bark and other adaptations to fire, the removal of understory and ladder fuels that promote the transition from surface to crown fire, and maintenance burning (Stephens et al. 2013). Such interventions have been demonstrated to reduce tree mortality during subsequent wildfire (Prichard et al. 2020). Recent work also highlights support for treatments that promote landscape heterogeneity through creation of clumps and gaps (Churchill et al. 2013); stand- and landscape-level heterogeneity have also been shown to increase forest resilience to wildfire (Koontz et al. 2020) and other disturbances such as beetle outbreaks (Seidl et al. 2016). At broader spatial scales, vegetation management projects and strategic fuel breaks can be used to restore more resilient patch mosaics and limit future fire spread into communities or vulnerable late-successional habitat (Hessburg et al. 2016, 2019).

Heterogeneity may be achieved by direct management intervention (mechanical thinning and prescribed fire), but strategically allowing wildfires to burn at low-to-moderate severity under tolerable fire weather conditions also reduces fuels and creates heterogeneity. In particular, where frequent-fire ecosystems are not substantially departed from historic norms, repeated low-to-moderate-intensity burning may confer resilience to forests by maintaining a reduced fuel load and perpetuating a low-severity fire regime (Larson et al. 2013, Walker et al. 2018, Kane et al. 2019). Furthermore, fires burning under benign to moderate conditions may interact with topography to support fire refugia (Krawchuk et al. 2016) that promote forest recovery (Coop et al. 2019). Following high-severity wildfires, strategic tree planting and forest management can also generate heterogeneous forest structure and composition (North et al. 2019). A synthesis of these many existing strategies to resist conversion could provide insight into whether and how long these tactics will be viable under climate change, and also inspire the development of new approaches to mitigate conversion in infrequent-fire forest types, where fewer management interventions are in use.

Although directing forest conversion is within the spectrum of management choices, it currently lacks adequate scientific underpinnings and is therefore poised to become an increasingly important research field. For example, ecological and ethical questions associated with managed relocation or assisted migration of genotypes, species, and vegetation types cover very broad terrain. Topics for applied research include the role of dispersal limitations, habitat connectivity, multispecies interactions, native and nonnative species interactions, no-analog climates, and probability of long-term establishment (Schwartz et al. 2012). In addition, a general framework is needed for conducting experimental adaptation for directing change, testing the efficacy of various tactics, and assessing how different approaches might interact and be sustained across larger spatial scales.

The fourth theme, to better *understand the ecological and social consequences of the choice to resist, accept, or direct conversion*, is key to creating operational models for adapting to change. Ecological and social values will be strongly affected by how the postfire assemblage of species that replaces a particular forest will translate to biodiversity and habitat availability, ecosystem processes and functions (e.g., hydrology), and ecosystem services, economic health, and cultural identity. Connecting science on these ecosystem functions and services to a diverse array of plausible vegetation types for each option to resist, accept, or direct conversion will be key to supporting experimental adaptation and a portfolio of informed management approaches.

Even with strong scientific support, managers may be constrained by agency practices and public expectations. Although there are risks, ultimately managers will require broader social license and support to operate outside of traditional models. Social science is needed to inform and support decisions about forest conversion (e.g., McWethy

et al. 2019), with a better understanding of how society values particular forests, and how those values, social acceptability, and agency mandates constrain a manager's decision space (Higuera et al. 2019). These topics each merit their own assessment of management-focused research needs. Furthermore, an era of profound and global ecological change may demand a strengthened ethical framework within which to consider decisions likely to have wide-reaching and lasting consequences.

Conclusions

Wildfire-driven forest conversion occurs when ecological resilience of forests to wildfire is overcome, leading to extensive and enduring areas of altered vegetation. Conversion is initiated by high-severity fire that removes areas of mature trees, and is maintained by a range of processes that impede tree regeneration, including distant tree seed sources, short-interval fires, or unfavorable postfire climate, further shaped by fire-vegetation feedbacks. An emerging body of research from across western North America highlights the strong potential for anthropogenic climate change and other human-induced changes to create conditions leading to fire-driven forest conversion. Numerous key uncertainties currently limit our capacity to project future changes, but also present research opportunities. However, the prospect of directional climate change beyond historical ranges of variability, and increased frequency and magnitude of extreme disturbance, compels us to consider the possibility of profound and persistent ecological change across forested ecosystems. As such, management and conservation efforts should align with expectations of increasing forest vulnerability to conversion. In an era of change, the forest that was there before the fire may not return.

Acknowledgments

This work arose from an oral session organized by JDC, CSR, and SAP at the US-International Association for Landscape Ecology annual meeting in 2019. The authors thank the many funding organizations, institutions, and individuals that have supported the research discussed herein. Helpful comments and suggestions were provided by Craig D. Allen, the editor, and three anonymous reviewers. In addition, support for this synthesis was provided by the National Fire Plan through agreement no. 15-CR-11223639-118 between the USFS Aldo Leopold Wilderness Institute and Western Colorado University.

References cited

Abatzoglou JT, Williams AP, Barbero R. 2019. Global Emergence of Anthropogenic Climate Change in Fire Weather Indices. *Geophysical Research Letters* 46: 326–336.

Abatzoglou JT, Williams AP. 2016. Impact of anthropogenic climate change on wildfire across western US forests. *Proceedings of the National Academy of Science* 113: 11770–11775.

Abella SR, Fornwalt PJ. 2015. Ten years of vegetation assembly after a North American mega fire. *Global Change Biology* 21: 789–802.

Airey Lauvaux C, Skinner CN, Taylor AH. 2016. High severity fire and mixed conifer forest-chaparral dynamics in the southern Cascade Range, USA. *Forest Ecology and Management* 363: 74–85.

Allen CD, Breshears DD. 1998. Drought-induced shift of a forest-woodland ecotone: Rapid landscape response to climate variation. *Proceedings of the National Academy of Sciences* 95: 14839–14842.

Anderegg WRL, Hicke JA, Fisher RA, Allen CD, Aukema J, Bentz B, Hood S, Lichstein JW, Macalady AK, McDowell N, et al. 2015. Tree mortality from drought, insects, and their interactions in a changing climate. *New Phytologist* 208: 674–683.

Andersen T, Carstensen J, Hernandez-Garcia E, Duarte CM. 2009. Ecological thresholds and regime shifts: Approaches to identification. *Trends in Ecology and Evolution* 24: 49–57.

Anderson-Teixeira KJ, Miller AD, Mohan JE, Hudiburg TW, Duval BD, DeLucia EH. 2013. Altered dynamics of forest recovery under a changing climate. *Global Change Biology* 19: 2001–2021.

Aplet GH, Cole DN. 2010. The trouble with naturalness: Rethinking park and wilderness goals. *BeNaturalness: Rethinking Park Wilderness Stewardship in an Era of Rapid Change* 12: 21–22.

Archibald S, Lehmann CER, Belcher CM. 2018. Biological and geophysical feedbacks with fire in the Earth system. *Environmental Research Letters* 13(33003).

Baker WL. 2009. *Fire ecology in Rocky Mountain landscapes*. Island Press.

Balch JK, Bradley BA, Abatzoglou JT, Nagy RC, Fusco EJ, Mahood AL. 2017. Human-started wildfires expand the fire niche across the United States. *Proceedings of the National Academy of Sciences* 114: 2946–2951.

Barnosky AD, Hadly EA, Bascompte J, Berlow EL, Brown JH, Fortelius M, Getz WM, Harte J, Hastings A, Marquet PA, Martinez ND. 2012. Approaching a state shift in Earth's biosphere. *Nature*. 486: 52–58.

Barton AM, Poulos HM. 2018. Pine versus oaks revisited: Conversion of Madran pine-oak forest to oak shrubland after high-severity wildfire in the Sky Islands of Arizona. *Forest Ecology and Management* 414: 28–40.

Battipaglia G, Saurer M, Cherubini P, Calfapietra C, McCarthy HR, Norby RJ, Francesca Cotrufo M. 2013. Elevated CO₂ increases tree-level intrinsic water use efficiency: Insights from carbon and oxygen isotope analyses in tree rings across three forest FACE sites. *New Phytologist* 197: 544–554.

Brown CD, Johnstone JF. 2012. Once burned, twice shy: Repeat fires reduce seed availability and alter substrate constraints on *Picea mariana* regeneration. *Forest Ecology and Management* 266: 34–41.

Brown PM, Wu R. 2005. Climate and disturbance forcing of episodic tree recruitment in a southwestern ponderosa pine landscape. *Ecology* 86: 3030–3038.

Buma B, Brown CD, Donato DC, Fontaine JB, Johnstone JF. 2013. The impacts of changing disturbance regimes on serotinous plant populations and communities. *BioScience* 63: 866–876.

Buma B, Weiss S, Hayes K, Lucash M. 2020. Wildland fire reburning trends across the US West suggest only short-term negative feedback and differing climatic effects. *Environmental Research Letters* 15 (art. 034026).

Calder WJ, Stefanova I, Shuman B. 2019. Climate–fire–vegetation interactions and the rise of novel landscape patterns in subalpine ecosystems, Colorado. *Journal of Ecology* 107: 1689–1703.

Campbell JL, Shinneman DJ. 2017. Potential influence of wildfire in modulating climate-induced forest redistribution in a central Rocky Mountain landscape. *Ecological Processes* 6: 7.

Chambers ME, Fornwalt PJ, Malone SL, Battaglia MA. 2016. Patterns of conifer regeneration following high severity wildfire in ponderosa pine: Dominated forests of the Colorado Front Range. *Forest Ecology and Management* 378: 57–67.

Churchill DJ, Larson AJ, Dahlgreen MC, Franklin JF, Hessburg PF, Lutz JA. 2013. Restoring forest resilience: From reference spatial patterns to silvicultural prescriptions and monitoring. *Forest Ecology and Management* 291: 442–457.

- Collins BM, Lydersen JM, Everett RG, Stephens SL. 2018. How does forest recovery following moderate-severity fire influence effects of subsequent wildfire in mixed-conifer forests?. *Fire Ecology* 14: 3.
- Collins BM, Miller JD, Thode AE, Kelly M, Van Wagtenonk JW, Stephens SL. 2009. Interactions among wildland fires in a long-established Sierra Nevada natural fire area. *Ecosystems* 12: 114–128.
- Coop JD, Parks SA, McClernan SR, Holsinger LM. 2016. Influences of prior wildfires on vegetation response to subsequent fire in a reburned Southwestern landscape. *Ecological Applications* 26: 346–354.
- Coop JD, DeLory TJ, Downing WM, Haire SL, Krawchuk MA, Miller C, Parisien MA, Walker RB. 2019. Contributions of fire refugia to resilient ponderosa pine and dry mixed-conifer forest landscapes. *Ecosphere* 10: e02809.
- Coppoletta M, Merriam KE, Collins BM. 2016. Post-fire vegetation and fuel development influences fire severity patterns in reburns. *Ecological Applications* 26: 686–699.
- Crausbay SD, Higuera PE, Sprugel DG, Brubaker LB. 2017. Fire catalyzed rapid ecological change in lowland coniferous forests of the Pacific Northwest over the past 14,000 years. *Ecology* 98: 2356–2369.
- Davis KT, Higuera PE, Sala A. 2018. Anticipating fire-mediated impacts of climate change using a demographic framework. *Functional Ecology* 32: 1729–1745.
- Davis KT, Dobrowski SZ, Higuera PE, Holden ZA, Veblen TT, Rother MT, Parks SA, Sala A, Maneta M. 2019a. Wildfires and climate change push low-elevation forests across a critical climate threshold for tree regeneration. *Proceedings of the National Academy of Sciences* 116: 6193–6198.
- Davis KT, Dobrowski SZ, Holden ZA, Higuera PE, Abatzoglou JT. 2019b. Microclimatic buffering in forests of the future: The role of local water balance. *Ecography* 42: 1–11.
- Dennison PE, Brewer SC, Arnold JD, Moritz MA. 2014. Large wildfire trends in the western United States, 1984–2011. *Geophysical Research Letters* 41: 2928–2933.
- Donato DC, Harvey BJ, Turner G. 2016. Regeneration of montane forests 24 years after the 1988 Yellowstone fires: A fire-catalyzed shift in lower treelines? *Ecosphere* 7: e01410.
- Enright NJ, Fontaine JB, Bowman DM, Bradstock RA, Williams RJ. 2015. Interval squeeze: Altered fire regimes and demographic responses interact to threaten woody species persistence as climate changes. *Frontiers in Ecology and the Environment* 13: 265–272.
- Falk DA, Watts AC, Thode AE. 2019. Scaling ecological resilience. *Frontiers in Ecology and Evolution* 7: 275.
- Falk DA. 2017. Restoration ecology, resilience, and the axes of change. *Annals of the Missouri Botanical Garden* 102: 201–216.
- Feddema JJ, Mast JN, Savage M. 2013. Modeling high-severity fire, drought and climate change impacts on ponderosa pine regeneration. *Ecological Modelling* 253: 56–69.
- Flannigan MD, Wotton BM, Marshall GA, De Groot WJ, Johnston J, Jurko N, Cantin AS. 2016. Fuel moisture sensitivity to temperature and precipitation: Climate change implications. *Climatic Change* 134: 59–71.
- Flatley WT, Fulé PZ. 2016. Are historical fire regimes compatible with future climate? Implications for forest restoration. *Ecosphere* 7: e01471.
- Folke C, Carpenter SR, Walker B, Scheffer M, Chapin T, Rockström J. 2010. Resilience thinking: Integrating resilience, adaptability and transformability. *Ecology and Society* 15 (4, art. 20).
- Ghazoul J, Burivalova Z, Garcia-Ulloa J, King LA. 2015. Conceptualizing forest degradation. *Trends in Ecology and Evolution* 30: 622–632.
- Guiterman CH, Margolis EQ, Allen CD, Falk DA, Swetnam TW. 2018. Long-term persistence and fire resilience of oak shrubfields in dry conifer forests of northern New Mexico. *Ecosystems* 21: 943–959.
- Gunderson, LH. 2000. Ecological Resilience: In theory and application. *Annual Review of Ecology and Systematics* 31: 425–439.
- Gustafson EJ. 2013. When relationships estimated in the past cannot be used to predict the future: Using mechanistic models to predict landscape ecological dynamics in a changing world. *Landscape Ecology* 28: 1429–1437.
- Haire SL, McGarigal K. 2010. Effects of landscape patterns of fire severity on regenerating ponderosa pine forests in New Mexico and Arizona, USA. *Landscape Ecology* 25: 1055–1069.
- Hanes CC, Wang X, Jain P, Parisien MA, Little JM, Flannigan MD. 2018. Fire-regime changes in Canada over the last half century. *Canadian Journal of Forest Research* 49: 256–269.
- Hansen WD, Turner MG. 2019. Origins of abrupt change? 2019. Postfire subalpine conifer regeneration declines nonlinearly with warming and drying. *Ecological Monographs* 89: e01340.
- Hansen WD, Romme WH, Ba A, Turner MG. 2016. Shifting ecological filters mediate postfire expansion of seedling aspen (*Populus tremuloides*) in Yellowstone. *Forest Ecology and Management* 362: 218–230.
- Harpold AA, Molotch NP. 2015. Sensitivity of soil water availability to changing snowmelt timing in the western US. *Geophysical Research Letters* 42: 8011–8020.
- Hart SJ, Henkelman J, McLoughlin PD, Nielsen SE, Truchon-Savard A, Johnstone JF. 2019. Examining forest resilience to changing fire frequency in a fire-prone region of boreal forest. *Global Change Biology* 25: 869–884.
- Harvey BJ, Donato DC, Turner MG. 2016a. Burn me twice, shame on who? Interactions between successive forest fires across a temperate mountain region. *Ecology* 97: 2272–2282.
- Harvey BJ, Donato DC, Turner MG. 2016b. Drivers and trends in landscape patterns of stand-replacing fire in forests of the US Northern Rocky Mountains. *Landscape Ecology* 31: 2367–2383.
- Harvey BJ, Donato DC, Turner MG. 2016c. High and dry: Post-fire tree seedling establishment in subalpine forests decreases with post-fire drought and large stand-replacing burn patches. *Global Ecology and Biogeography* 25: 655–669.
- Héon J, Arseneault D, Parisien M-A. 2014. Resistance of the boreal forest to high burn rates. *Proceedings of the National Academy of Sciences* 111: 13888–13893.
- Hessburg PF, Miller CL, Povak NA, Taylor AH, Higuera PE, Prichard SJ, North MP, Collins BM, Hurteau MD, Larson AJ, Allen CD. 2019. Climate, environment, and disturbance history govern resilience of western North American forests. *Frontiers in Ecology and Evolution* 7: 239.
- Hessburg PF, Spies TA, Perry DA, Skinner CN, Taylor AH, Brown PM, Stephens SL, Larson AJ, Churchill DJ, Povak NA, Singleton PH. 2016. Tamm review: Management of mixed-severity fire regime forests in Oregon, Washington, and Northern California. *Forest Ecology and Management* 366: 221–250.
- Higuera PE, Metcalf AL, Miller C, Buma B, McWethy DB, Metcalf EC, Ratajczak Z, Nelson CR, Chaffin BC, Stedman RC, McCaffrey S. 2019. Integrating subjective and objective dimensions of resilience in fire-prone landscapes. *BioScience* 69: 379–388.
- Hurteau MD, Liang S, Westerling AL, Wiedinmyer C. 2019a. Vegetation-fire feedback reduces projected area burned under climate change. *Scientific Reports* 9: 2838.
- Hurteau MD, North MP, Koch GW, Hungate BA. 2019b. Managing for disturbance stabilizes forest carbon. *Proceedings of the National Academy of Sciences* 116: 10193–10195.
- Hutto RL, Bond ML, DellaSala DA. 2015. Using bird ecology to learn about the benefits of severe fire. Pages 55–88 in DellaSala DA, Hanson CT, eds. *The Ecological Importance of Mixed-Severity Fires*. Elsevier.
- Jackson RB, Randerson JT, Canadell JG, Anderson RG, Avissar R, Baldocchi DD, Bonan GB, Caldeira K, Duffenbaugh NS, Field CB, et al. 2008. Protecting climate with forests. *Environmental Research Letters* 3: 44006.
- Johnstone JF, Allen CD, Franklin JF, Frelich LE, Harvey BJ, Higuera PE, Mack MC, Meentemeyer RK, Metz MR, Perry GL, et al. 2016. Changing disturbance regimes, ecological memory, and forest resilience. *Frontiers in Ecology and the Environment* 14: 369–378.
- Kane VR, Bartl-Geller BN, Kane JT, Jeronimo SMA, North MP, Collins B, Lydersen J. 2019. First-entry fires can create forest tree clump and opening patterns characteristic of historic resilient forests. *Forest Ecology and Management* 454: 117659.

- Keeley JE, Nelman G, Fotheringham CJ. 1999. Immaturity risk in a fire-dependent pine. *Journal of Mediterranean Ecology* 1: 41–48.
- Keenan TF, Hollinger DY, Bohrer G, Dragoni D, Munger JW, Schmid HP, Richardson AD. 2013. Increase in forest water-use efficiency as atmospheric carbon dioxide concentrations rise. *Nature* 499: 324–328.
- Kemp KB, Higuera PE, Morgan P, Abatzoglou JT. 2019. Climate will increasingly determine post-fire tree regeneration success in low-elevation forests, Northern Rockies, USA. *Ecosphere* 10: e02568.
- Kitzberger T, DA Falk, AL Westerling, TW Swetnam. 2017. Direct and indirect climate controls predict heterogeneous early mid 21st century wildfire burned area across western and boreal North America. *PLOS ONE* 12 (art. e0188486).
- Knutti R, Sedláček J. 2013. Robustness and uncertainties in the new CMIP5 climate model projections. *Nature Climate Change* 3: 369–373.
- Koontz MJ, North MP, Werner CM, Fick SE, Latimer AM. 2020. Local forest structure variability increases resilience to wildfire in dry western US coniferous forests. *Ecology Letters* 23: 483–494.
- Krawchuk MA, Haire SL, Coop J, Parisien MA, Whitman E, Chong G, Miller C. 2016. Topographic and fire weather controls of fire refugia in forested ecosystems of northwestern North America. *Ecosphere* 7: e01632.
- Larson AJ, Belote RT, Cansler CA, Parks SA, Dietz MS. 2013. Latent resilience in ponderosa pine forest: Effects of resumed frequent fire. *Ecological Applications* 23: 1243–1249.
- Lawrence DM, Hurtt GC, Arneeth A, Brovkin V, Calvin KV, Jones AD, Jones CD, Lawrence PJ, de Noblet-Ducoudré N, Pongratz J, Seneviratne SI. 2016. The Land Use Model Intercomparison Project (LUMIP) contribution to CMIP6: Rationale and experimental design. *Geoscientific Model Development* 9: 2973–2998.
- Liang S, Hurteau MD, Westerling AL. 2017. Response of Sierra Nevada forests to projected climate-wildfire interactions. *Global Change Biology* 23: 2016–2030.
- Littell JS, McKenzie D, Wan HY, Cushman SA. 2018. Climate change and future wildfire in the Western United States: An ecological approach to nonstationarity. *Earth's Future* 6: 1097–1111.
- Loehman RA, Keane RE, Holsinger LM. 2020. Simulation modeling of complex climate, wildfire, and vegetation dynamics to address wicked problems in land management. *Frontiers in Forests and Global Change* (29 January 2020). <https://doi.org/10.3389/ffgc.2020.00003>.
- Lydersen JM, Collins BM, Coppoletta M, Jaffe MR, Northrop H, Stephens SL. 2019. Fuel dynamics and reburn severity following high severity fire in a Sierra Nevada mixed-conifer forest. *Fire Ecology* 15: 43.
- Lynch EA. 1998. Origin of a park–forest vegetation mosaic in the Wind River Range, Wyoming. *Ecology* 79: 1320–1338.
- Margolis EQ, Swetnam TW, Allen CD. 2007. A stand-replacing fire history in upper montane forests of the southern Rocky Mountains. *Canadian Journal of Forest Research* 37: 2227–2241.
- McCaughey LA, Robles MD, Woolley T, Marshall RM, Kretchun A, Gori DF. 2019. Large-scale forest restoration stabilizes carbon under climate change in Southwest United States. *Ecological Applications* 29: e01979.
- McKenzie D, Littell JS. 2017. Climate change and the eco-hydrology of fire: Will area burned increase in a warming western USA? *Ecological Applications* 27: 26–36.
- McWethy DB, Schoennagel T, Higuera PE, Krawchuk M, Harvey BJ, Metcalf EC, Schultz C, Miller C, Metcalf AL, Buma B, et al. 2019. Rethinking resilience to wildfire. *Nature Sustainability* 2: 797–804.
- Meadow AM, Ferguson DB, Guido Z, Horangic A, Owen G, Wall T. 2015. Moving toward the deliberate coproduction of climate science knowledge. *Weather Climate and Society* 7: 179–191.
- Miller JD, Safford HD, Crimmins M, Thode AE. 2009. Quantitative Evidence for Increasing Forest Fire Severity in the Sierra Nevada and Southern Cascade Mountains, California and Nevada, USA. *Ecosystems* 12: 16–32.
- Miller JE, Root HT, Safford HD. 2018. Altered fire regimes cause long-term lichen diversity losses. *Global Change Biology* 24: 4909–4918.
- Minckley TA, Shriver RK, Shuman B. 2012. Resilience and regime change in a southern Rocky Mountain ecosystem during the past 17 000 years. *Ecological Monographs* 82: 49–68.
- Nelson KN, Turner MG, Romme WH, Tinker DB. 2016. Landscape variation in tree regeneration and snag fall drive fuel loads in 24-year old post-fire lodgepole pine forests. *Ecological Applications* 26: 2424–2438.
- Nolan C, Overpeck JT, Allen JR, Anderson PM, Betancourt JL, Binney HA, Brewer S, Bush MB, Chase BM, Cheddadi R, Djarnali M. 2018. Past and future global transformation of terrestrial ecosystems under climate change. *Science* 361: 920–923.
- North MP, Stevens JT, Greene DF, Coppoletta M, Knapp EE, Latimer AM, Restaino CM, Tompkins RE, Welch KR, York RA, Young DJ. 2019. Tamm review: Reforestation for resilience in dry western US forests. *Forest Ecology and Management* 432: 209–224.
- O'Connor CD, Falk DA, Lynch AM, Swetnam TW, Wilcox C. 2017. Disturbance and productivity interactions mediate stability of forest composition and structure. *Ecological Applications* 27: 900–915.
- Pan Y, Birdsey RA, Fang J, Houghton R, Kauppi PE, Kurz WA, Phillips OL, Shvidenko A, Lewis SL, Canadell JG, et al. 2011. A large and persistent carbon sink in the world's forests. *Science* 333: 988–993.
- Parks SA, Miller C, Parisien MA, Holsinger LM, Dobrowski SZ, Abatzoglou J. 2015. Wildland fire deficit and surplus in the western United States. *Ecosphere* 6: 1–13.
- Parks SA, Holsinger LM, Miller C, Parisien MA. 2018a. Analog-based fire regime and vegetation shifts in mountainous regions of the western US. *Ecography* 41: 910–921.
- Parks SA, Parisien MA, Miller C, Holsinger LM, Baggett LS. 2018b. Fine-scale spatial climate variation and drought mediate the likelihood of reburning. *Ecological Applications* 28: 573–586.
- Parks SA, Dobrowski SZ, Shaw JD, Miller C. 2019. Living on the edge: Trailing edge forests at risk of fire-facilitated conversion to non-forest. *Ecosphere* 10: e02651.
- Pausas JG, Keeley JE. 2014. Evolutionary ecology of resprouting and seeding in fire-prone ecosystems. *New Phytologist* 204: 55–65.
- Peñuelas J, Canadell JG, Ogaya R. 2011. Increased water-use efficiency during the 20th century did not translate into enhanced tree growth. *Global Ecology and Biogeography* 20: 597–608.
- Prichard SJ, Stevens-Rumann CS, Hessburg PF. 2017. Tamm Review: Shifting global fire regimes: Lessons from reburns and research needs. *Forest Ecology and Management* 396: 217–233.
- Prichard SJ, Povak N, Kennedy MC, Peterson DW. 2020. Fuel treatment effectiveness following the 2014 Carlton Complex Fire in semi-arid forests of north-central Washington State. *Ecological Applications* (22 February 2020, art. e02104).
- Reyer CB, Brouwers N, Rammig A, Brook BW, Epila J, Grant RF, Holmgren M, Langerwisch F, Leuzinger S, Lucht W, Medlyn B. 2015. Forest resilience and tipping points at different spatio-temporal scales: Approaches and challenges. *Journal of Ecology* 103: 5–15.
- Rother MT, Veblen TT, Furman LG. 2015. A field experiment informs expected patterns of conifer regeneration after disturbances under changing climate conditions. *Canadian Journal of Forest Research* 45: 1607–1616.
- Rowe JS, Scotter GW. 1973. Fire in the boreal forest. *Quaternary Research* 3: 444–464.
- Savage M, Mast JN. 2005. How resilient are southwestern ponderosa pine forests after crown fires? *Canadian Journal of Forest Research* 35: 967–977.
- Scheffer M, Bascompte J, Brock WA, Brovkin V, Carpenter SR, Dakos V, Held H, Van Nes EH, Rietkerk M, Sugihara G. 2009. Early warning signals for critical transitions. *Nature* 461: 53–59.
- Schoennagel T, Veblen TT, Romme WH. 2004. The interaction of fire, fuels, and climate across Rocky Mountain forests. *BioScience* 54: 661–676.
- Schwartz MW, Hellmann JJ, McLachlan JM, Sax DF, Borevitz JO, Brennan J, Camacho AE, Ceballos G, Clark JR, Doremus H, Early R. 2012. Managed relocation: Integrating the scientific, regulatory, and ethical challenges. *BioScience* 62: 732–743.

- Seidl R, Donato DC, Raffa KF, Turner MG. 2016. Spatial variability in tree regeneration after wildfire delays and dampens future bark beetle outbreaks. *Proceedings of the National Academy of Sciences* 113: 13075–13080.
- Serra-Diaz JM, Maxwell C, Lucash MS, Scheller RM, Laflouer DM, Miller AD, Tepley AJ, Epstein HE, Anderson-Teixeira KJ, Thompson JR. 2018. Disequilibrium of fire-prone forests sets the stage for a rapid decline in conifer dominance during the 21st century. *Scientific Reports* 8: 6749.
- Someone C, Maneta MP, Holden ZA, Sapes G, Sala A, Dobrowski SZ. 2019. Coupled ecohydrology and plant hydraulics modeling predicts ponderosa pine seedling mortality and lower treeline in the US Northern Rocky Mountains. *New Phytologist* 221: 1814–1830.
- Singleton MP, Thode AE, Sánchez Meador AJ, Iniguez JM. 2019. Increasing trends in high-severity fire in the southwestern USA from 1984 to 2015. *Forest Ecology and Management* 433: 709–719.
- Stark SC, Breshears DD, Garcia ES, Law DJ, Minor DM, Saleska SR, Swann ALS, Villegas JC, Aragão LEOC, Bella EM, et al. 2016. Toward accounting for ecoclimate teleconnections: Intra- and inter-continental consequences of altered energy balance after vegetation change. *Landscape Ecology* 31: 181–194.
- Steel ZL, Safford HD, Viers JH. 2015. The fire frequency-severity relationship and the legacy of fire suppression in California forests. *Ecosphere* 6: 1–23.
- Stein BA, Glick P, Edelson N, Staudt A. 2014. *Climate-Smart Conservation: Putting Adaptation Principles into Practice*. National Wildlife Federation.
- Stephens SL, Agee JK, Fule PZ, North MP, Romme WH, Swetnam TW, Turner MG. 2013. Managing forests and fire in changing climates. *Science* 6154: 41–42.
- Stevens JT, Safford HD, Harrison S, Latimer AM. 2015. Forest disturbance accelerates thermophilization of understory plant communities. *Journal of Ecology* 103: 1253–1263.
- Stevens-Rumann CS, Kemp KB, Higuera PE, Harvey BJ, Rother MT, Donato DC, Morgan P, Veblen TT. 2018. Evidence for declining forest resilience to wildfires under climate change. *Ecology Letters* 21: 243–252.
- Stevens-Rumann CS, Morgan P. 2019. Tree regeneration following wildfires in the western US: A review. *Fire Ecology* 15: 15.
- Stralberg D, Wang X, Parisien M-A, Robinne F-N, Sóllymos P, Mahon CL, Nielsen SE, Bayne EM. 2018. Wildfire-mediated vegetation change in boreal forests of Alberta, Canada. *Ecosphere* 9: e02156.
- Stringham TK, Krueger WC, Shaver PL. 2003. State and transition modeling: An ecological process approach. *Rangeland Ecology and Management* 56: 106–113.
- Svenning JC, Sandel B. 2013. Disequilibrium vegetation dynamics under future climate change. *American Journal of Botany* 100: 1266–1286.
- Temperton VM, Hobbs RJ, Nuttle T, Halle S. 2004. *Assembly Rules and Restoration Ecology: Science and Practice of Restoration Ecology*. Island Press.
- Tepley AJ, Thomann E, Veblen TT, Perry GL, Holz A, Paritsis J, Kitzberger T, Anderson-Teixeira KJ. 2018. Influences of fire–vegetation feedbacks and post-fire recovery rates on forest landscape vulnerability to altered fire regimes. *Journal of Ecology* 106: 1925–1940.
- Tepley AJ, Thompson JR, Epstein HE, Anderson-Teixeira KJ. 2017. Vulnerability to forest loss through altered postfire recovery dynamics in a warming climate in the Klamath Mountains. *Global Change Biology* 23: 4117–4132.
- Turner MG, Baker WL, Peterson CJ, Peet RK. 1998. Factors influencing succession: Lessons from large, infrequent natural disturbances. *Ecosystems* 1: 511–523.
- Turner MG, Brazianus KH, Hansen WD, Harvey BJ. 2019. Short-interval severe fire erodes the resilience of subalpine lodgepole pine forests. *Proceedings of the National Academy of Sciences* 116: 11319–11328.
- Turner MG, Romme WH. 1994. Landscape dynamics in crown fire ecosystems. *Landscape Ecology* 9: 59–77.
- van Mantgem PJ, Nesmith JC, Keifer M, Knapp EE, Flint A, Flint L. 2013. Climate stress increases forest fire severity across the western United States. *Ecology Letters* 16: 1151–1156.
- van Mantgem PJ, Falk DA, Williams E, Das AJ, Stephenson, NL. 2018. Pre-fire drought and competition mediate post-fire conifer mortality in western U.S. National Parks. *Ecological Applications* 28: 1730–1739.
- Walker RB, Coop JD, Parks SA, Trader L. 2018. Fire regimes approaching historic norms reduce wildfire-facilitated conversion from forest to non-forest. *Ecosphere* 9: e02182.
- Wang X, Parisien MA, Taylor SW, Candau JN, Stralberg D, Marshall GA, Little JM, Flannigan MD. 2017. Projected changes in daily fire spread across Canada over the next century. *Environmental Research Letters* 12: 025005.
- Westerling AL. 2016. Increasing western US forest wildfire activity: Sensitivity to changes in the timing of spring. *Philosophical Transactions of the Royal Society B* 371: 20150178.
- Whitman E, Parisien MA, Thompson DK, Flannigan MD. 2019. Short-interval wildfire and drought overwhelm boreal forest resilience. *Scientific Reports* 9: 1–2.
- Wine ML, Makhnin O, Cadol D. 2018. nonlinear long-term large watershed hydrologic response to wildfire and climatic dynamics locally increases water yields. *Earth's Future* 6: 997–1006.

Jonathan D. Coop is an associate professor in the School of Environment and Sustainability at Western Colorado University, in Gunnison. Sean A. Parks is a research ecologist with the Aldo Leopold Wilderness Research Institute, Rocky Mountain Research Station, US Forest Service, in Missoula, Montana. Camille S. Stevens-Rumann is an assistant professor with the Forest and Rangeland Stewardship Department at Colorado State University, in Fort Collins. Shelley Causbay is a senior scientist with Conservation Science Partners, in Fort Collins, Colorado. Philip E. Higuera is an associate professor with the Department of Ecosystem and Conservation Sciences at the University of Montana, in Missoula, Montana. Matthew D. Hurteau is an associate professor in the Department of Biology at the University of New Mexico, in Albuquerque. Alan Tepley and Ellen Whitman are research scientists with Natural Resources Canada, Canadian Forest Service, Northern Forestry Centre, in Edmonton, Alberta, Canada. Timothy J. Assal is an assistant professor in the Department of Geography at Kent State University, in Kent, Ohio. Brandon M. Collins is a research scientist at the Center for Fire Research and Outreach, at the University of California, Berkeley, in Berkeley, California, and with the Pacific Southwest Research Station, US Forest Service, in Davis, California. Kimberley T. Davis is a postdoctoral research scientist in the Department of Ecosystem and Conservation Sciences at the University of Montana, in Missoula. Solomon Dobrowski is a professor in the Department of Forest Management at the University of Montana, in Missoula. Donald A. Falk is a professor in the School of Natural Resources and the Environment at the University of Arizona, in Tucson. Paula J. Fornwalt is a research ecologist at the Rocky Mountain Research Station, US Forest Service, in Fort Collins, Colorado. Peter Z. Fulé is a professor in the School of Forestry at Northern Arizona University, in Flagstaff. Brian J. Harvey is an assistant professor in the School of Environmental and Forest Sciences at the University of Washington, in Seattle. Van R. Kane is a research professor in the School of Environmental and Forest Sciences at the University of Washington, in Seattle. Caitlin Littlefield is a postdoctoral research associate at the Rubenstein School of Environment and Natural Resources at the University of Vermont, in Burlington. Ellis Q. Margolis is a research ecologist for the US Geological Survey, New Mexico Landscapes Field Station, in Santa Fe. Malcolm North is a research plant ecologist with the US Forest Service, Pacific Southwest Research Station, in Mammoth Lakes, California. Marc-André Parisien is a research scientist with Natural Resources Canada, Canadian Forest Service, Northern Forestry Centre, in Edmonton, Alberta, Canada. Susan Prichard is a research scientist in the School of Environmental and Forest Sciences at the University of Washington, in Seattle. Kyle C. Rodman is a postdoctoral research associate in the Department of Forest and Wildlife Ecology at the University of Wisconsin, in Madison.