

# **Trees and Logs Important to Wildlife in the Interior Columbia River Basin**

Evelyn L. Bull, Catherine G. Parks, and Torolf R. Torgersen

Companion to General Technical Report  
PNW-GTR-390


Published by:

---

U.S. Department of Agriculture, Forest Service  
Pacific Northwest Research Station  
General Technical Report PNW-GTR-391  
May 1997

---

# INTRODUCTION



More than 80 species of birds, mammals, reptiles, and amphibians use living trees with decay, trees with brooms, hollow trees, snags, and logs in the interior Columbia River basin. Animals use these structures for foraging, nesting, denning, roosting, and resting. Most notable of these wood-using wildlife species are the primary cavity nesters, the woodpeckers and nuthatches, that excavate nest cavities in decayed wood in standing trees. Their vacated cavities are subsequently used by many other birds and mammals.

About 25 percent of the bird species nesting in the northern Rocky Mountain forests are cavity nesters (McClelland and others 1979). Many of the primary and secondary cavity-nesting birds eat forest insects and thus play an important role in regulating their populations. Machmer and Steeger (1995) provide a thorough review of the effect these birds have in reducing numbers of tree-feeding insects.

Tree decay is an important ecological process affecting wildlife habitat. Once begun, the decay process can take hundreds of years as a tree dies, falls to the ground, and decomposes into the forest floor. As it decays, the tree supports many different wildlife groups that use it for foraging substrate, for nesting, and for shelter. The group of wood decomposers is diverse. It includes many arthropods, but in this publication, we focus on decay fungi as the primary organisms in decay. Decay fungi are an essential resource to forest ecosystems in storing and retaining nutrients and in soil development.

This publication provides managers with a description of the ecological processes that create wildlife trees, snags, and logs. It provides qualitative and quantitative information on five distinct structures: living trees with decay (such as internal decay), hollow trees, trees with brooms (misshapen branches), dead trees (snags), and down woody material (logs). We describe the value of these structures to wildlife, the decay or infection processes that produce each type of structure, principles to help in selecting the best structures to retain, and management implications. Although this document cannot prescribe the amount of landscape to manage

for a particular species or a specific, desired population size, it does present information managers can use to make informed decisions regarding the conservation and enhancement of dead wood structures most valuable to wildlife. Forest management practices that fail to properly manage wood components may adversely affect wildlife, soil and stream quality, and forest ecosystem functions.

The information included here applies to coniferous forest lands in the interior Columbia River basin: the area east of the crest of the Cascade Range in Oregon and Washington, all of Idaho, and a portion of western Montana (fig. 1). Tree species characteristic of high-elevation forests, (such as whitebark pine [*Pinus albicaulis*], alpine larch [*Larix lyallii*], mountain hemlock [*Tsuga mertensiana*], and Pacific silver fir [*Abies amabilis*]) are excluded because they typically grow where active management for wildlife trees is seldom needed; these areas are seldom logged. Most of the tree species we deal with here are conifers, although black cottonwood (*Populus trichocarpa*), quaking aspen (*Populus tremuloides*), and paper birch (*Betula papyrifera*) are




Figure 1.—The interior Columbia River basin.

**Snag distribution**—Snags can either be left in clusters or scattered across the landscape, although where to retain them depends largely on where they currently are. Typically, snags are in clumps naturally because the agents of disease, insects, fire, or flooding that kill many trees act in a localized area. Clusters of snags may be easier to protect during management activities than snags evenly distributed across the landscape. Worker safety is an issue when harvesting adjacent to snags. Retaining snags in clusters, as well as the live trees between the snags, keeps workers away from snags and provides canopy cover for wildlife using the snags.

Two studies on woodpecker use of snag clusters reported different results. No difference was found between woodpecker use of snag clusters versus scattered snags in western Oregon (Chambers 1997). In areas where stand-replacement fires had burned in ponderosa pine/Douglas-fir stands in Idaho, Saab and Dudley (1997) found that cavity nesters as a group select clusters of snags for nest sites instead of evenly spaced trees.

3-2  
Most studies recommend managing snags in every 5- to 25-acre patch (Bate 1995, Evans and Martens 1995). Because woodpeckers are territorial, only one pair of each species occupies the same territory; therefore, to attain the maximum density of cavity nesters, snags need to be distributed across the landscape.

In addition, retaining snags on all aspects and on all positions of the slope ensures habitat availability for wildlife species with different preferences. For example, pileated woodpeckers tend to roost in draws during summer but roost higher on slopes in winter (Bull and others 1992). Martens rest or travel in riparian areas a third of the time in summer but only half that often in winter; upper slopes are used half the time in winter but only a third of the time in summer (Bull 1995).

Retaining snags close to living trees provides cover for cavity users. If snags are retained in the middle of a clearcut, species like northern flickers, Lewis' woodpeckers, bluebirds, and American kestrels (*Falco sparverius*) will use them for nesting, but most other woodpeckers will not nest in open areas.

In many areas, fuel-wood cutters are likely to remove snags along roads. This potential problem can be remedied by not leaving snags in areas that are accessible to fuel-wood cutters or by not counting snags left adjacent to accessible roads toward the number retained for a particular stand.

In summary, wherever the objective is to provide viable populations of primary and secondary cavity nesters, prudent managers will provide large-diameter snags when available snags in a variety of structural classes, snags in every 5- to 25-acre stand and in clusters if available, and snags on all slope aspects and positions of the slope, and adjacent to green trees.

**Snag densities**—Limited information is available on numbers of snags to retain for wildlife species in the interior Columbia River basin (Bate 1995, Bull and Holthausen 1993, Dixon 1995, Evans and Martens 1995) (table 1). The guide most widely used in the past, Thomas and others (1979), prescribed the number of nest and roost trees to leave for specified woodpecker populations, but the number was based on a hypothetical, untested model and did not include any snags for foraging. Three studies (Bate 1995, Bull and

Holthausen 1993, Dixon 1995) conducted in eastern Oregon have shown that retaining foraging structure is essential, in addition to nest and roost trees in managed landscapes. The Thomas model provided only two roost trees per pair per year, yet research has shown that individual pairs of pileated woodpeckers and white-headed woodpeckers use considerably more than two per year (Bull and others 1992, Dixon 1995). Radio-telemetry studies have shown that home range sizes of pileated woodpeckers (Bull and Holthausen 1993), white-headed woodpeckers (Dixon 1995), and three-toed and black-backed woodpeckers (Goggans and others 1988) are considerably larger than those used in the Thomas model. Raphael and White (1984) found that the relation between numbers of snags and cavity nesters is not linear, which was assumed in the Thomas model. The substitution factor used in the Thomas model is variable and largely a function of snag density. Neitro and others (1985) thought allowing substitution of snags that reduced the number retained was not appropriate. The Thomas model did not take into account the habitat needs of some of the secondary cavity nesters, like bats and brown creepers, that use such snag features as loose bark. In addition, Bull and Holthausen (1993) found lower densities of pileated woodpeckers in nine study areas than predicted by Thomas and others (1979) based on the number of snags present. The above studies present new data suggesting that some of the assumptions and data used in the Thomas model are not valid, and that the prescribed snag densities need to be revised upward.

We know of only three studies in the interior Columbia River basin that have calculated both the density of snags and woodpeckers in managed and unmanaged landscapes. Bate (1995) used six study areas in ponderosa pine stands and found woodpecker abundance was best predicted by hard snag density, large green tree density, canopy height, and number of canopy layers. Of the nine study areas, only one represented a relatively unmanaged forest with the number of snags that probably approximated historical ones. In this stand, Bate found 3.8 snags greater than 10 inches d.b.h., and 60 percent were larger than 20 inches d.b.h. These snag numbers supported 32.3 woodpeckers per 247 acres and represented the study area that most likely supported viable populations of cavity nesters. Dixon (1995) found similar results in some of the same study areas Bate used.

Another study relating woodpecker density to snag density was conducted in northeastern Oregon (Bull and Holthausen 1993). The two study areas that contained viable populations of pileated woodpeckers had an average of 4.2 and 3.8 snags (10 inches or larger d.b.h.) per acre and 0.5 and 0.4 nesting pairs per 247 acres, respectively. The two study areas were in lightly managed landscapes predominated by mixed-coniferous stands with a high density of logs.

Evans and Martens (1995) recommended densities of snags for retention on the Payette National Forest based on their ecological value, encompassing soil health, seedling regeneration, moisture retention, nutrient recycling, and wildlife use (table 1). The numbers that they recommended were derived from actual snag numbers identified during

timber inventory and vegetation plots but were scaled down to the lower ranges of the snag numbers for the recommendations and include snags 10 inches d.b.h. and larger.

Ideally, data would be available on the exact number of snags required to support specific populations of primary and secondary cavity nesters. Unfortunately, this kind of information is not available. We do know, however, that the snag numbers presented by Thomas and others (1979) are not adequate to support the populations intended because of a lack of foraging strata and invalid assumptions used in the model. If management agencies have an objective to manage for viable populations of woodpeckers, providing numbers of snags that have been shown to support viable populations in the recent studies would be prudent.

Although snags are the key structure typically focused on for cavity nesters, realizing the importance of the forest surrounding the snags also is essential. Some species like the pileated and white-headed woodpeckers and American martens are associated with older forests. For white-headed woodpeckers, the snags must be in association with large-diameter ponderosa pines. For pileated woodpeckers and martens, logs, large-diameter green trees, and a dense canopy are needed. Leaving large-diameter snags or green trees in younger forests, also has value as providing biological legacies.

**Snag retention in burns**—The above management implications apply to healthy green stands. With vast areas of Oregon, Washington, and Idaho burned by wildfires in the 1990s, there is much interest in determining numbers of snags to retain in burned areas after harvesting. Saab and Dudley (1997) reported that woodpeckers selected nest sites with snag densities greater than those measured at random sites in ponderosa pine/Douglas-fir forests with stand-replacing fires. From 10 to 14 snags per acre were retained in burned stands that were harvested; about 38 to 42 snags per acre were in the unharvested burned stands. Seven species of cavity nesters using the burned stands selected nest sites with more than 20 snags per acre, where snags were distributed in clumps. Lewis' and white-headed woodpeckers, and American kestrels selected the largest, most heavily decayed snags available for nesting. Evans and Martens (1995) presented recommendations for numbers of snags to leave in burned stands.

**Green tree replacements**—Snags are a dynamic resource; old snags fall and living trees die to become new snags. Snag-dependent wildlife need a continuous supply of snags over time. To provide a continuum of snag habitat, future snags must be planned for by leaving green trees to eventually become snags in managed stands. To determine the number of green tree replacements, information on fall rate of standing snags, snag density, live stem density, and mortality rate of live stems is required. Various models have been developed to calculate the number of green trees to provide for recruitment when specific stand inventory information is available (Bull and others 1980, Cimon 1983, Schommer and others 1993). If snags are to be created by killing green trees when the stand becomes snag deficient, the number of green trees required is less because the manager does not have to rely on natural mortality to replace

those snags that fall. Creating snags is costly, however, and stands are seldom monitored to determine when trees need to be killed.

### Snag Longevity

Snag longevity, or the amount of time a snag stands, is essential information for managing the snag resource. The length of time snags stand is a primary factor in determining the number of green tree replacements needed to maintain a specific snag density over time. Snag longevity is a function of many factors including species, diameter and height, percentage of heartwood, cause of death, soil type and moisture, forest type, and prevalence of windstorms. Because so many factors influence how long a snag stands, fall rates should be determined by species and size class on each management area. Listed below are studies that report snag fall rates. When possible, the amount of time when 50 percent of snags in a particular study have fallen will be reported.

Several studies have reported fall rates of beetle-killed ponderosa pine. Keen (1955) reported that 50 percent of the ponderosa pines in southern Oregon and northeastern California fell 9 to 10 years after death on pumice soils and 6 to 7 years after death on loam soils. The percentages of snags that fell after 7 years by diameter class, are presented in the following tabulation:

Diameter class (inches)	Percentage
10-18	75
20-28	65
30-38	50
40-48	35
> 50	30

Most of the beetle-killed ponderosa pines in Colorado fall between 7 and 10 years after infestation (Schmid and others 1985). Trees were 7 to 22 inches d.b.h.

In northeastern Oregon, half the ponderosa pines less than 10 inches d.b.h. fell 7 to 8 years after death and half those 10 to 19 inches d.b.h. fell after 8 to 9 years (Bull 1983). Beetle-killed lodgepole pines have similar fall rates, with half of those less than 10 inches d.b.h. falling within 6 to 7 years and half of those 10 inches and larger falling after 7 to 8 years (Bull 1983). In a second study in northeastern Oregon (Bull and Partridge 1986), half of the ponderosa pine snags less than 20 inches d.b.h. fell after 6 years; half of the ponderosa pine snags 20 inches and larger fell after 9 years.

Beetle-killed spruce in Colorado stood much longer than ponderosa pines reported in the studies above. Mielke (1950) found that 84 percent of beetle-killed spruce remain standing after 25 years. Hinds and others (1965) found that 72 percent of beetle-killed spruce remain standing after 20 years.

Fire-killed snags may or may not stand longer than beetle-killed snags, depending on the area. Dahms (1949) reported that a little over half the ponderosa pine snags are down 10 years after a fire in central Oregon. In Montana, nearly half of all lodgepole pine snags fall by the fifth season (Lyon 1977). In California, Kimmey (1955) described the

breakup of a fire-killed stand, although specific fall rates were not presented. In the fourth and fifth year after the fire, many ponderosa and Jeffrey pine (*Pinus jeffreyi*) broke off at the ground or up to 50 feet above the ground. By the fifth year, the general breakup continued until only scattered barkless snags and stubs remain.

In northeastern Oregon, different methods of killing ponderosa pines were investigated (Bull 1996c, Bull and Partridge 1986). Trees that had been topped stand the longest; half of those less than 20 inches d.b.h. are down after 10 to 12 years; half of those 20 inches or larger are down after 12 to 17 years. Half the trees injected with a silvicide are down 7 years after treatment. Half the girdled trees are down 9 years after treatment.

Fall rates of snags that have been created differ. Ponderosa pines infected with dwarf mistletoe were frilled and poisoned with an herbicide in the Southwest; after 10 years, 95 percent of those trees are down (Fairweather 1995).

All of these studies suggest that most ponderosa pine and lodgepole pine that died from fire or beetles in the interior Columbia River basin have fall rates of less than 10 years. Small-diameter snags fall sooner than do larger ones.

## Sampling Techniques

Knowledge of snag density is essential for managing forest stands effectively, as well as for complying with standards and guidelines. Various methods have been used to determine snag densities: area-wide counts, fixed-radius circular plots, variable-radius plots using prisms (Bull and others 1990), and variable-strip transects (Bate and others, in press). Total counts of snags over large areas are very time-consuming but yield the most accurate density information. Fixed-radius plots typically need to be at least 1 acre because snags are usually scarce and variance is high. Many plots are usually required to accurately estimate snag density.

Bate and others (in preparation) have developed a more efficient method of sampling snags than either area-wide counts or fixed-radius plots: data are collected from a pilot sample transect and used in a computer program. The program determines the optimal length and width of a strip transect within a given area; the optimal size is that which minimizes the cost and variance. The program then provides the necessary sample size to obtain a snag density within the desired statistical bounds. Transect lengths are either 164 or 328 feet; transect widths range from 33 to 131 feet. The

**Table 1—Density of snags reported or recommended in four areas; numbers are snags per acre larger than 10 inches d.b.h. (unless otherwise stated), and the percentages represent snags larger than 20 inches d.b.h.**

Forest type	Locality			
	Western Idaho <sup>a</sup>	Central Oregon <sup>b</sup>	Northeast Oregon <sup>c</sup>	North-central Idaho <sup>d</sup>
Ponderosa pine:				
Open canopy <sup>e</sup>	1.6(75%)			
Closed canopy <sup>f</sup>	4.8(44%)	4(60%)		
Mixed conifer:				
Open canopy	2.5(72%)			
Closed canopy	9.0(39%)		3.8(29%)	10-48
Lodgepole pine:				
Open canopy	4.5			
Closed canopy	7.7			
Spruce/fir	9.5(21%)			

<sup>a</sup> Recommendations for snag densities selected from the lower ranges of snag numbers on timber inventory plots and fixed-radius plots on the Payette National Forest; different snag levels recommended for burned and unburned stands (Evans and Martens 1995).

<sup>b</sup> Snag numbers based on variable-width transects conducted in an unmanaged stand of old growth in central Oregon where woodcutting was prohibited (Bate 1995).

<sup>c</sup> Numbers based on density of snags in two 4,000-acre study areas in northeastern Oregon that supported viable numbers of pileated woodpeckers; areas had some timber harvest but no woodcutting (Bull and Holthausen 1993).

<sup>d</sup> Densities based on timber inventory plots in unmanaged stands of large saw timber. The snag density varied depending on what snags were included in the count. The 10 snags per acre included only snags 15 inches d.b.h. or larger and 50 feet or taller. The 48 snags per acre included all snags 15 inches d.b.h. or larger of any height. Western white pine snags were not counted (Craig 1995).

<sup>e</sup> Stands with less than 30 percent canopy closure.

<sup>f</sup> Stands with more than 30 percent canopy closure.

program also provides a "distribution index," which is an estimate of the percentage of the sampled landscape supporting target snags or trees.

### Creating Wildlife Trees

In managed stands, retaining existing snags is the most ecologically sound and economical approach to providing wildlife trees. If stands are devoid of snags, however, an option is to alter or kill living trees. Depending on the method used, killing trees can be costly and may not produce the desired decay conditions. Years may pass before the decay is sufficiently advanced to promote cavity-nester excavations.

Bull and Partridge (1986) investigated six methods of killing ponderosa pine. They determined that topping trees with either a chainsaw or explosives produced snags that stood the longest and received the greatest nest use by woodpeckers (figs. 48 and 49). In this study, girdling, fungal inoculation, and beetles attracted by pheromones did not consistently kill the tree. Trees killed by girdling or silvicides fell over too quickly to provide wildlife nest trees.

In a study in New Mexico, all of the ponderosa pines killed by girdling were used for foraging by woodpeckers. Most of the trees greater than 16 inches d.b.h. were standing after 7 years, and about 30 percent contained woodpecker nest sites (Parks 1996b). Because the climate in the New Mexico sites is drier than in the Oregon study area used by Bull and Partridge (1986), basal decay of killed trees may have been less, so the trees stood longer.

Recent work by Parks and others (1996a, 1996b) documents a new method of inoculating live trees with decay fungi. Six years after inoculation of 60 living western larch, 14 percent contained woodpecker cavities near the point of inoculation. These trees may remain alive for decades with a pocket of decay that woodpeckers can use for nesting. Trees may be logged next to these trees without safety concerns, and live trees are less likely than dead trees to be lost to fuel-wood cutters. These preliminary results for western larch suggest that inoculation produces desirable wildlife trees at a lesser cost than killing trees to create snags.

A variety of wildlife tree structures can be created with a chainsaw in standing, living trees. Trees can be topped leaving a horizontal branch structure on which ospreys (*Pandion haliaetus*) can build nests (fig. 50). A depression can be cut into the top of a tree that has been topped to produce a nest site for great gray owls (fig. 51). Artificial cavities and hollows can be cut into standing and downed trees (figs. 52, 53, and 54). Slits can be cut into trees to create roosting habitat for bats (fig. 55). These techniques and others are described by Brown (1996).

Three primary considerations in creating wildlife trees are the target wildlife species for which the structure is being created; the tree species, size of tree, and habitat locality most likely to be used by the targeted wildlife species; and the method of altering or killing trees that is most likely to produce long-standing trees in a given geographic area. Rainfall, snowfall, incidence of windstorms, and prevalent decay organisms all must be considered.

Thousands of trees have been killed to create snags in the interior Columbia River basin in the last 15 years. Few of them have been monitored, however, to determine whether long-lasting, quality snags were actually produced. Because climate and decay organisms differ with area, results of management activities, such as snag creation, must be monitored to ensure that results warrant the cost of these treatments. We include a data form to monitor wildlife use of created snags in appendix A.

### Review

- Ponderosa pine, western larch, quaking aspen, and paper birch are favored tree species for nest sites of woodpeckers in many areas in the interior Columbia River basin.
- Large-diameter snags provide nest habitat for the greatest variety of cavity nesters and stand longer than smaller snags.
- Snags can be classed into three structural classes that are a simplified version of past decay class categories.
- Snags should be provided in every 5- to 25-acre stand, in clusters if available, on all slope aspects and positions of the slope, and adjacent to green trees.
- Recent studies have shown that the snag model developed by Thomas and others (1979) did not accurately predict woodpecker abundance based on snag abundance. New information suggests that foraging strata (snags and other structures) must be incorporated into any snag model, and that snag numbers need to be revised upward.
- Published data suggest that populations of cavity nesters were viable in stands of ponderosa pine and mixed-conifer forests that contained about four snags per acre, a large component of old-growth stands, and abundant logs.
- Snags can be retained over time by using models for green tree replacements.
- Snag longevity depends on cause of death, tree species, diameter, height, amount of heartwood, geographic area, and site conditions.
- Creating snags and other wildlife structures can mitigate the loss of natural habitat, but retaining existing structures is most cost effective and ecologically sound.

