

# The influence of forest fragmentation and landscape pattern on American martens

Utah

CHRISTINA D. HARGIS, JOHN A. BISSONETTE\* and DAVID L. TURNER†

USDA Forest Service, South-west Forest Science Complex, 2500 S. Pine Knoll, Flagstaff, AZ 86001, USA; \*Utah Cooperative Fish and Wildlife Research Unit (USGS-BRD), Department of Fisheries and Wildlife, Utah State University, Logan, UT 84322-5390, USA; and †USDA Forest Service Intermountain Research Station, 860 N 1200 E, Logan, UT 84321, USA

is a forest interior sp in Utah

## Summary

1. We investigated the effects of forest fragmentation on American martens (*Martes americana* Rhoads) by evaluating differences in marten capture rates (excluding recaptures) in 18 study sites with different levels of fragmentation resulting from timber harvest clearcuts and natural openings. We focused on low levels of fragmentation, where forest connectivity was maintained and non-forest cover ranged from 2% to 42%.

2. Martens appeared to respond negatively to low levels of habitat fragmentation, based on the significant decrease in capture rates within the series of increasingly fragmented landscapes. Martens were nearly absent from landscapes having > 25% non-forest cover, even though forest connectivity was still present.

3. Marten capture rates were negatively correlated with increasing proximity of open areas and increasing extent of high-contrast edges. Forested landscapes appeared unsuitable for martens when the average nearest-neighbour distance between open (non-forested) patches was < 100 m. In these landscapes, the proximity of open areas created strips of forest edge and eliminated nearly all forest interior.

4. Small mammal densities were significantly higher in clearcuts than in forests, but marten captures were not correlated with prey abundance or biomass associated with clearcuts.

5. Conservation efforts for the marten must consider not only the structural aspects of mature forests, but the landscape pattern in which the forest occurs. We recommend that the combination of timber harvests and natural openings comprise < 25% of landscapes  $\geq 9 \text{ km}^2$  in size.  $3.49 \text{ m}^2 = 2220$

6. The spatial pattern of open areas is important as well, because small, dispersed openings result in less forest interior habitat than one large opening at the same percentage of fragmentation. Progressive cutting from a single patch would retain the largest amount of interior forest habitat.

**Key-words:** edge density, habitat fragmentation, *Martes americana*, proximity index, timber harvests.

Journal of Applied Ecology (1999) 36, 157-172

## Introduction

The modification of forested landscapes through land management practices has fostered a growing interest in the effects of habitat fragmentation on wildlife.

Habitat fragmentation, originally defined as the formation of isolated fragments from a formerly continuous habitat (Wilcox 1980; Harris 1984), has taken on a broader meaning in recent times, reflecting an awareness that fragmentation can affect organisms long before the original habitat is reduced to remnant patches. In the fullest sense, fragmentation is disruption in continuity in any temporal or functional

small patches within  
not no value

- 300' canopy  
no marten use

edge = 150' into  
forest

- effect of  
frog will  
increase with  
harshness  
of habitat.

- canopy opening  
causes loss  
of interior  
for

322'

81  
369 km  
341  
= 75.9 yr

818960  
400000  
420000  
at

domain (Lord & Norton 1990). Habitat fragmentation occurs along a continuum, from landscapes dominated by the original cover type to landscapes where the original type is reduced to remnant patches.

The majority of habitat fragmentation studies have focused on extreme examples of fragmentation, after the original cover type has ceased to function as the dominant landscape element. By focusing on remnant habitat patches, these studies have examined the effect that patch area and isolation have on population size, extinction and recolonization rates, and dispersal rates of individual species, as well as on species diversity (Diamond 1975; Whitcomb *et al.* 1981; Harris 1984; Rosenberg & Raphael 1986; Verboom & van Apeldoorn 1990). More recently, investigations have turned to less extreme cases of fragmentation, and have focused on the cohesiveness or contagion of the original habitat type when it is the dominant landscape element (Franklin & Forman 1987; O'Neill *et al.* 1988; Ripple, Bradshaw & Spies 1991; Spies, Ripple & Bradshaw 1994; Wallin, Swanson & Marks 1994). These studies investigate potential restrictions on the movement of organisms, the spread of fire or disease, or the flow of nutrients at the landscape scale (Turner 1987; Turner *et al.* 1989; Bartell & Brenkert 1991; Turner & Dale 1991; Wiens *et al.* 1993; Ims 1995). The area of the original habitat generally is not measured when it is the dominant landscape element, because it forms one continuous patch and may extend beyond the defined landscape boundary. The area and distance between disturbance patches are nevertheless of interest because they serve as barriers to movement and affect the spatial configuration of the original habitat.

Our study focused on low fragmentation of forested landscapes, where the extent of forest has been altered by timber harvests but still comprises the dominant landscape element. We investigated the effects of habitat fragmentation on the American marten *Martes americana* Rhoads, a carnivorous mammal typically associated with mature forest systems. American martens appear sensitive to habitat fragmentation, because populations generally are low in areas fragmented by the clearcut method of timber harvest (Soutiere 1979; Snyder & Bissonette 1987; Thompson & Harestad 1994). However, marten responses to gradual increases in fragmentation are not well known. Most studies of timber harvest effects have been in areas where levels of fragmentation were not differentiated and where fragmentation was quite high, between 41% and 60% clearcut (Soutiere 1979; Steventon & Major 1982; Snyder & Bissonette 1987; Katnik, Harrison & Hodgman 1994; Thompson 1994; Potvin & Breton 1997). In our investigation, we were primarily interested in the response of martens to a gradual increase in fragmentation when forest connectivity was maintained, and therefore focused our study on a series of forested landscapes where non-forest cover ranged from 2% to 42%.

Brainerd (1990) proposed a model of marten response to fragmentation when non-forest cover occupied 55% of a hypothetical landscape. He predicted that habitat quality would increase if cut sizes were fine-grained, permitting marten movements across the clearcuts while taking advantage of increased prey in open areas. Habitat quality was predicted to decrease with medium- and coarse-grained cut sizes because they would restrict marten movements. A similar model was later developed by Thompson & Harestad (1994) that echoed Brainerd's (1990) predictions. Examining the full range of potential fragmentation, they predicted two possible changes in marten carrying capacity with increasing removal of timber. Under the clearcut method of harvest, equivalent to Brainerd's (1990) medium-grained or coarse-grained cut sizes, they predicted a linear negative decline. With selective cutting and patch cuts <3 ha, they predicted an increase in carrying capacity at low cutting levels, followed by a precipitous decline at approximately 20–30% of forest cover removal, due to increased predation of martens in openings, high edge density, and loss of forest interior. Thus, their model differed from Brainerd's (1990) only in predicting a decline at an earlier stage of fragmentation.

Although martens avoid clearcuts and other large openings, especially in the winter (Soutiere 1979; Clark & Campbell 1979; Steventon & Major 1982; Hargis & McCullough 1984), low levels of fragmentation may have little effect on martens as long as forest connectivity is maintained. In fact, a positive response to low fragmentation may be expected, because in some locales martens forage in brushy clearcuts during summer (Steventon & Major 1982; Katnik 1992) and hunt along forest-meadow edges (Simon 1980; Spencer, Barrett & Zielinski 1983). Having a small proportion of the landscape in open areas may be favourable to martens because of the increase in abundance and diversity of small mammals associated with clearcuts, and because marten diets are not restricted to forest-associated prey (Weckwerth & Hawley 1962; Koehler & Hornocker 1977; Baskirk & MacDonald 1984). The initial positive response would be followed eventually by a negative response as forested habitat became limiting.

To test these predictions, we compared marten capture rates across a series of increasingly fragmented landscapes. Our main objective was to determine whether marten abundance changed with incremental increases in habitat fragmentation caused by the combined effects of natural openings and timber clearcuts. If a correlation was found between capture rates and fragmentation, our secondary objective was to look for specific levels of fragmentation that suggested shifts in habitat quality in either a positive or negative direction. Preliminary results from this study were reported in Hargis & Bissonette (1997). This paper reports further analyses on the effects of landscape pattern on martens as measured by several landscape metrics.




Fig. 4. Scatterplot of individual marten captures with (a) percentage of total stems in snags and (b) vole captures 100 trap nights<sup>-1</sup>.

with 0–8 individual captures per site during any given trapping period. Although capture rates differed among sites (described below), we found no significant difference in capture rates among trapping periods within any given site ( $\chi^2 = 2.89$ , d.f. = 3;  $P = 0.41$ ). Sites with high captures were consistently high, and sites with zero captures remained low, resulting in a nearly stable ranking of sites across all years, and confirming that differences in capture rates were due to factors other than seasonal variation or sampling error.

Marten captures were negatively correlated with loss of forest habitat, as measured by the percentage of each site in open areas, and this relationship was significant ( $P = 0.03$ ) when snag abundance was added to the model (Table 5). Capture rates were variable in sites with low fragmentation, but only one marten was captured in 937 trap nights in the four sites having >0.25% open areas (Fig. 5a).

Martens showed significant responses to landscape pattern as well as loss of habitat (Fig. 5b–d and Table 5). Capture rates were lowest in landscapes with large, closely spaced open areas, as measured both by mean proximity index and nearest-neighbour distance. Landscapes with an average distance between open areas <100 m had no marten captures. Also, landscapes with high edge density had fewer marten captures. We did not observe a significant correlation between marten captures and mass fractal dimension, although the trend was for low to zero captures in landscapes where the forest matrix was highly convoluted (Fig. 5e and Table 5).

## COMPARISON OF FRAGMENTATION MODELS

We examined correlations between percentage of each site in openings and the four measures of landscape pattern to assess the degree of additional information provided by these measures (Table 6). Correlations were generally high, with mass fractal dimension having the highest correlation with habitat loss ( $r = -0.97$ ) and nearest-neighbour distance the lowest ( $r = -0.67$ ). In addition, the four measures were interrelated among themselves, with edge density and mass fractal dimension having the strongest correlation ( $r = -0.96$ ).

Partly as a consequence of these interrelationships, nearest-neighbour distance, edge density and percentage of site in openings were similar in their role as explanatory variables of marten capture rates. Used alone, each explained approximated 20% of the variation in marten capture rates and  $P$ -values were similar. When used in combination with snag abundance, percentage of site in openings and edge density became better predictors, but the relationships for nearest-neighbour distance were weakened. Mass fractal dimension provided the poorest fit of all fragmentation measures examined, both when used alone or in combination with snag abundance.

The combined effects of mean proximity index and snag abundance provided the best fit for explaining differences in marten capture rates, based on a comparison of  $P$ -values and  $R^2$  terms from all fragmentation models examined (Table 5). The contribution of mean proximity index was greater than that of percentage of site in openings, which became an insignificant term when both measures were included in a model (mean  $P$  for each variable, respectively, based on 35 bootstrap runs = 0.37 and 0.01).

## Discussion

Martens appeared to respond negatively to low levels of habitat fragmentation, based on a significant decrease in capture rates observed across a series of increasingly fragmented landscapes. Martens were rarely detected in sites with >25% open areas, even though forest connectivity was still present.

Martens were sensitive not only to loss of habitat but also to the size and proximity of open areas. As measured by the mean proximity index, marten capture rates decreased as open patches became more closely situated (Fig. 5b). When the average nearest-neighbour distance between open patches was <100 m, nearly all forest interior had been converted to narrow strips of forest that functioned only as edge habitat (Fig. 1c).

We provide several caveats to aid in interpreting these results. First,  $R^2$  values associated with all significant relationships were low. It may be unrealistic to expect high  $R^2$  values in a study design where landscapes are the units of replication, but nevertheless

**Table 5.** Statistical significance of fragmentation models explaining individual marten capture rates, with mean  $P$ -values (confidence intervals in parentheses) derived from chi-square tests performed on 35 bootstrap data sets per model (error d.f. = 15 in each model). Models with  $P < 0.05$  are indicated with an asterisk. The second column contains both the mean univariate  $P$ -value and the mean multivariate  $P$ -value of the fragmentation measure when snag abundance is added to the model

Model parameters	$P$ -value of fragmentation measure	$P$ -value of snag abundance in multivariate model	$R^2$
% of site in openings	0.06 (0.05, 0.07)		0.23 (0.21, 0.25)
* % of site in openings, snag abundance	0.03 (0.02, 0.04)	0.01 (0.01, 0.01)	0.50 (0.48, 0.52)
* Proximity index	0.01 (0.01, 0.01)		0.48 (0.47, 0.49)
Proximity index, snag abundance	0.04 (0.03, 0.05)	0.07 (0.06, 0.08)	0.59 (0.58, 0.60)
* Nearest-neighbour distance	0.03 (0.03, 0.03)		0.21 (0.20, 0.22)
* Nearest-neighbour distance, snag abundance	0.05 (0.02, 0.08)	0.03 (0.01, 0.05)	0.42 (0.40, 0.44)
Edge density	0.08 (0.07, 0.09)		0.19 (0.17, 0.21)
* Edge density, snag abundance	0.05 (0.01, 0.09)	0.01 (0.00, 0.02)	0.47 (0.45, 0.49)
Mass fractal dimension	0.12 (0.10, 0.14)		0.17 (0.15, 0.19)
Mass fractal dimension, snag abundance	0.11 (0.04, 0.18)	0.02 (0.01, 0.03)	0.43 (0.39, 0.47)

the low  $R^2$  values indicate substantial unexplained variation in marten capture rates.

Secondly, the size chosen for study areas was small relative to marten home range scale. The size of marten home ranges in the Uinta Mountains is unknown, but in the neighbouring state of Wyoming the mean summer home ranges (95% minimum convex polygon) of seven females was 669 ( $\pm 67$  SE) ha and that of eight males was 1820 ( $\pm 153$  SE) ha (O'Doherty, Ruggiero & Henry 1997), sizes that would suggest only 1–2 martens per landscape. Nevertheless, we expected several martens to occur in each landscape because home range boundaries were unlikely to correspond to study area boundaries, and therefore partial home ranges of several martens could be represented. Martens exhibit intrasexual territoriality (reviewed by Powell 1994) but  $>1$  female range typically occurs within those of males. Non-territorial juveniles sometimes share ranges with territorial adults (Archibald & Jessup 1984). Rather than view these landscapes from a home range perspective, we viewed each landscape as a window providing a snapshot of marten numbers at a given point in time.

A third caveat is that marten capture rates represent relative, not absolute, marten numbers. Sites with no marten captures may have contained martens but at densities too low to detect during the 6-night trapping period. To increase our confidence in interpreting zero values, we resampled all sites in which no martens

were caught, and in all cases no martens were captured during these second attempts. Nevertheless, we cannot conclude that no martens were present.

Also, we were unable to determine whether martens in our samples were residents or transients. This is an important consideration, because high numbers of transients could indicate a population sink (Pulliam 1988). Although we do not know how many transients occurred in each site, we do know that sites with high marten captures contained residents. The two sites with the highest number of captures were the only sites where we recaptured the same individuals in subsequent years. These sites also represented two of the four sites with lactating females, denoting reproductive activity. In the site with the highest capture rate, the average weight of males was above average for all sites, and general body condition was fair to excellent. Thus, there is no evidence to suggest that sites with high captures were simply areas of dispersal.

Our final caveat regards the apparent threshold in marten occurrence when open areas were approximately 25% of the total landscape. Martens in our sites may have exhibited a stronger response to low levels of fragmentation than would be expected in geographical areas with less harsh conditions. The clearcuts in our sites provided no habitat for martens because cut areas generally were stripped of both vegetation and logging slash after timber harvest. Moreover, the top soil was considerably disturbed, often




Fig. 5. Scatterplot of marten capture rates with five measures of fragmentation: (a) percentage of site in openings; (b) mean proximity index; (c) mean nearest-neighbour distance; (d) edge density; and (e) mass fractal dimension.

Table 6. Correlation matrix for landscape metrics and snag abundance, with all coefficients >|0.70| highlighted in bold

	Marten	% open	Edge density	Proximity index	Nearest neighbour distance	Mass fractal dimension	Snags
Marten	1.00	-0.50	-0.45	0.59	-0.56	0.41	-0.53
% open		1.00	<b>0.97</b>	<b>0.85</b>	-0.67	-0.97	0.28
Edge			1.00	<b>0.74</b>	-0.72	-0.96	0.26
Proximity index				1.00	-0.50	-0.79	0.28
Nearest-neighbour distance					1.00	0.59	-0.26
Mass fractal dimension						1.00	-0.32
Snags							1.00

retarding the establishment of conifer growth for a decade or more.

Elsewhere, martens apparently can forage in clearcuts containing structure in the form of regenerating deciduous or conifer vegetation and brush (Stevenson

& Major 1982; Katnik 1992; Potvin & Breton 1997) and may tolerate higher levels of fragmentation under these circumstances. Soutiere (1979) found marten using areas that were 60% cut in Maine, although at lower population levels than in uncut areas. We find it

note worthy, however, that Chapin & Harrison (1998), working in the same general vicinity as Soutiere, found no adult female marten territories where >31% of the landscape was in regenerating vegetation. The median percentage of home range area in regenerating patches was 20% for 14 resident adult females and 22% for 13 resident adult males (Chapin & Harrison 1998). Also, Thompson (1994) found resident marten at only one of five sites with logged forests, and clearcuts on these sites were vegetated with shrubs and quaking aspen *Populus tremuloides*. These studies indicate that marten residency, if not marten presence, may be affected by fairly low levels of fragmentation, regardless of the vegetation present in open areas. In areas where openings are poorly vegetated for long periods, we anticipate that martens would respond to fragmentation at approximately the same level that we observed. For example, clearcut management comparable to that of the Uinta Mountains occurs in much of the lodgepole pine and spruce forests of the intermountain states of western North America, and we consider our results relevant to marten conservation in that geographical area.

Our findings may be pertinent to the conservation of the Eurasian pine marten *Martes martes* Linnaeus as well. Of 46 British mammals analysed for vulnerability to fragmentation, the pine marten was considered one of the most vulnerable due to relatively low population density, slow breeding, fairly poor dispersal capabilities and close affinity to semi-natural habitats (Bright 1993). In the former USSR, notable declines in marten abundance were apparent when mature forest was reduced to 25–35% of total area (Grakov 1972). In northern Sweden, pine marten densities were found to be two to three times higher within an old-growth forest reserve than in the surrounding landscape of commercially managed, fragmented forest (Björvall, Nilsson & Norling 1977; cited in Brainerd 1997). In southern Sweden and south-eastern Norway, pine martens consistently preferred later-successional seres of spruce-dominated forest and avoided open habitats throughout the year (Brainerd 1997).

Martens in Scotland appear to tolerate high levels of forest fragmentation, but Balharry (1993) found that the normal rodent diet is supplemented by carrion, mostly from red deer *Cervus elaphus* culling operations and, to a lesser extent, from hill sheep *Ovis aries*. Carrion accounted for 35% of the estimated weight intake of all marten foods in two study sites. Nevertheless, some minimum area of woodland seemed important for establishing breeding territories, because breeding adults had significantly more woodland in their territories than juveniles and non-breeding adults (Balharry 1993). In a comparison of two study sites in Scotland, Balharry (1993) found that male marten home ranges were three times larger in the site having only 3.5% woodland than in the site with 27% woodland.

American martens may be somewhat more sensitive

to fragmentation than pine martens, and may require a certain proportion of forest interior within their home ranges. As summarized by Buskirk & Powell (1994), martens appear to use structural components of mature forests to avoid predators (Drew 1995), to gain access to prey in winter (Hargis & McCullough 1984; Corn & Raphael 1992; Sherburne & Bissonette 1993; Sherburne & Bissonette 1994) and to gain thermal advantages, especially while resting (Buskirk *et al.* 1989; Taylor 1993; Raphael & Jones 1997). Each of these could be affected by loss of forest interior habitat. Forest patches <100 m wide may not provide sufficient escape cover from known predators such as red fox *Vulpes vulpes*, coyote *Canis latrans* and raptors (Hodgman *et al.* 1997; Thompson 1994). Availability of preferred prey may also be a factor. Mills (1995) found lower densities of California red-backed voles *Clethrionomys californicus* near edges than in forest interiors, and Nordström (1995) found lower densities of southern red-backed voles in narrow forest corridors compared to contiguous forest blocks during a low population year. Thermal advantages may also be lost in forests that have been reduced to narrow strips. Chen, Franklin & Spies (1995) found that microclimatic edge effects can extend >240 m into old-growth Douglas-fir forests. Therefore, a forest interspersed with closely spaced open patches may lack the thermal conditions normally associated with mature forests due to landscape pattern.

Edge habitat *per se* is not necessarily detrimental to martens. Marten use of edge may depend on the habitat composition on either side (Buskirk & Powell 1994). In California, martens showed preferential use of edge that bordered mesic meadows (Simon 1980; Spencer, Barrett & Zielinski 1983) and in Maine edge between residual conifer forests and regenerating clearcuts was used in proportion to availability (Chapin 1995; Chapin & Harrison 1998). In our study, the correlation between edge density and mean proximity index made it difficult to determine whether martens responded negatively to edge, or whether the actual response was to the loss of forest interior from closely spaced patches, and edge was simply a covariate.

In addition to loss of interior habitat, martens may avoid landscapes with abundant openings due to the increased energetic costs of circumventing the open areas. It may be energetically prohibitive to defend a territory or forage in a home range that is widely diffused and interspersed with large patches of unsuitable habitat (Thompson & Colgan 1994).

We found no evidence that martens benefit from the increase in abundance and diversity of prey associated with clearcuts. We cannot state this conclusively because we did not collect dietary information, but our trapping data did not indicate a positive response to prey in clearcuts. Although small mammal densities were higher in clearcuts than in forest habitat, marten capture rates were not correlated with biomass of prey found in clearcuts, or with the abundance of any

species associated with clearcuts. Information on red squirrel (*Tamiasciurus hudsonicus* Trouessart) or snowshoe hare abundance, which we did not obtain, would not have changed the relationship between martens and clearcut-associated prey, because both are associated with forests (Raphael 1988; Thompson & Colgan 1994). Seven of the eight incidental captures of snowshoe hares were in forests.

We found a negative correlation between marten capture rates and snag abundance, but martens showed a positive response to snags in other studies (Spencer, Barrett & Zielinski 1983; Lofroth 1993; Drew 1995), partly because of their use of large diameter snags for resting sites (Campbell 1979; Spencer, Barrett & Zielinski 1983). In our study, snags were a common habitat feature in all sites and may not have been limiting to martens, but snag abundance varied among sites due to differences in mortality from mountain pine beetle *Dendroctonus ponderosae* Hopkins. The representation of snags in the overstorey ranged from 9% in an uninfected site to 65% in a heavily infected area. Sites with high beetle mortality were frequently selected for salvage timber harvests, and therefore sites with high snag densities also had the greatest number of clearcuts. Thus, the negative correlation between martens and fragmentation is mirrored by the negative correlation between martens and snag abundance. Because snag abundance improved the fit of all fragmentation models (Table 5), however, it is more than a correlate with fragmentation, and may be influencing marten ecology in ways that our data did not reflect. The more open canopy cover associated with high snag densities may cause edge effects to extend further into forests, causing a greater loss of forest interior at any given level of fragmentation than sites with low snag abundance.

#### RECOMMENDATIONS FOR MANAGEMENT AND RESEARCH

Our study was designed to assess changes in marten capture rates over a range of habitat fragmentation, but we were unable to separate the effects of habitat loss from the effects of landscape pattern. All measures of landscape pattern except mass fractal dimension were significantly correlated with marten captures, but each change in the measure of pattern was associated with a change in available habitat. To isolate the effects of landscape pattern, it would be necessary to sample martens in several landscapes having the same level of habitat loss but differing in landscape pattern. Landscapes with 20–35% of the area in openings appear fruitful for further investigations, based on the notable change in capture rates we encountered at this level. We predict that within this range of fragmentation martens are sensitive to landscape pattern alone, and that landscapes lacking forest interior may not sustain reproducing populations.

The effects of patch size and spatial distribution of forest interior are illustrated using three landscape patterns at the same level of fragmentation, with each landscape having approximately 20% of the area in openings (Fig. 6). The first landscape represents one of our study sites with open areas in a dispersed pattern that has resulted in the loss of much of the forest interior (Fig. 6a). The other images are of two simulated landscapes having 20% openings in which the openings are either clustered (Fig. 6b) or aggregated




Fig. 6. Three landscape patterns with approximately 20% of the area in openings: (a) an actual landscape in Utah; (b) a simulated landscape with clustered patches; and (c) a simulated landscape with openings aggregated into a single patch to maximize forest interior. Black represents forest and grey represents open areas.

into a single patch (Fig. 6c). The simulations were created with a computer program that places patches on a landscape at any specified proportion and spacing (Hargis, Bissonette & David 1997), using digitized clearcut patch shapes from the Uinta Mountains. The comparison clearly illustrates the greater size of forest interior and fewer disruptions in continuity of habitat that occurs when unsuitable habitat is clustered or occupies a single patch.

Assuming American martens do require forest interior for reproductive habitat, management should consider the guidelines developed by Franklin & Forman (1987) for conservation of forest interior species in managed landscapes. They recommend that timber harvests be undertaken progressively outward from a single patch or in clustered cuts so that disturbance patches are consolidated and larger areas of undisturbed forest could be maintained. For martens, progressive cutting rather than clustered cuts would be preferred, because the forested buffers left between tightly clustered clearcuts have little apparent value to martens, and represent additional loss of habitat rather than useable forest. Progressive cutting from a single patch would create maximum aggregation of disturbance and retain the largest amount of interior forest habitat.

#### Acknowledgements

This project was carried out within the Utah Cooperative Fish and Wildlife Research Unit (USGS Biological Resources Division). We thank F. Bartlett for logistical support of this project and K. Paulini and other Forest Service employees who provided us with necessary maps and aerial photos. We are indebted to S. Kuerner, H. Sloan, D. Masters, T. Sajwaj, K. Lischak, A. Krawarik, C. Zank, R. Vinkey, P. Lortz and R. Rood for assisting in collecting data under arduous conditions, and to the martens for being there. B. Milne and T. Keitt provided software for calculations of mass fractal dimension. Thanks to S. Brainerd for sending us P. Majewski's translation of Grakov (1972). We also thank P. Bright, S. W. Buskirk, M. Conover, J. David, S. Durham, J. Long, D. Ramsey and M. Ritchie for their helpful review of manuscript drafts. We acknowledge use of the Khoros image processing environment, a trademark of Khoros Research Inc., and FRAGSTATS, a spatial pattern analysis program in the public domain developed by K. McGarigal and B. Marks.

#### References

- Archibald, W.R. & Jessup, R.H. (1984) Population dynamics of the pine marten (*Martes americana*) in the Yukon Territory. *Northern Ecology and Resource Management* (eds R. Olson, R. Hastings & F. Geddes), pp. 81–97. University of Alberta Press, Edmonton, Canada.
- Balharay, D. (1993) *Factors affecting the distribution and population density of pine martens (Martes martes L.) in*

Scotland. PhD Thesis. University of Aberdeen, Aberdeen, UK.

- Bartell, S.M. & Brenkert, A.L. (1991) A spatial-temporal model of nitrogen dynamics in a deciduous forest watershed. *Quantitative Methods in Landscape Ecology* (eds M.G. Turner & R.H. Gardner), pp. 379–398. Springer-Verlag, New York, NY.
- Björvall, A., Nilsson, E. & Nörling, L. (1977) The importance of old-growth forest for capercaillie and marten. *Fauna Och Flora*, 72, 31–38 [in Swedish].
- Brainerd, S.M. (1990) The pine marten and forest fragmentation: a review and general hypothesis. *Transactions of the 19th International Congress of Game Biologists* (ed. S. Myrberget), pp. 421–434. Trondheim, Norway.
- Brainerd, S.M. (1997) *Habitat selection and range use by the Eurasian pine marten (Martes martes) in relation to commercial forestry practices in southern boreal Scandinavia*. PhD Thesis. Agricultural University of Norway, Norway.
- Bright, P.W. (1993) Habitat fragmentation – problems and predictions for British mammals. *Mammal Review*, 23, 101–111.
- Brown, J.K. (1974) *Handbook for Inventorying Downed Woody Material*. General Technical Report INT-16. USDA Forest Service, Ogden, UT.
- Buskirk, S.W. & MacDonald, S.O. (1984) Seasonal food habits of martens in south-central Alaska. *Canadian Journal of Zoology*, 62, 944–950.
- Buskirk, S.W. & Powell, R.A. (1994) Habitat ecology of fishers and American martens. *Martens, Sables, and Fishers: Biology and Conservation* (eds S.W. Buskirk, A.S. Harestad, M.G. Raphael & R.A. Powell), pp. 283–296. Cornell University Press, Ithaca, NY.
- Buskirk, S.W., Forrest, S.C., Raphael, M.G. & Harlow, H.J. (1989) Winter resting site ecology of martens in the central Rocky Mountains. *Journal of Wildlife Management*, 53, 191–196.
- Campbell, T.M. (1979) *Short-term effects of timber harvests on pine marten ecology*. MS Thesis. Colorado State University, Fort Collins, CO.
- Chapin, T.G. (1995) *Influence of landscape pattern and forest type on use of habitat by marten in Maine*. MS Thesis. University of Maine, Orono, ME.
- Chapin, T.G. & Harrison, D.J. (1998) Influence of landscape pattern on habitat use by American marten in an industrial forest. *Conservation Biology*, 12, 1327–1338.
- Chen, J., Franklin, J.F. & Spies, T.A. (1995) Growing-season microclimatic gradients from clearcut edges into old-growth Douglas-fir forests. *Ecological Applications*, 5, 74–86.
- Clark, T.W. & Campbell, T.M. (1979) Population organization and regulatory mechanisms of pine martens in Grand Teton National Park, Wyoming. *Conference on Scientific Research in National Parks, Vol. 1* (ed. R.M. Linn), pp. 293–295. National Park Service Transactions and Proceedings Series 5, National Park Service, Washington, DC.
- Corn, J.G. & Raphael, M.G. (1992) Habitat characteristics at marten subnivean access sites. *Journal of Wildlife Management*, 56, 442–448.
- Daniel, T.W. & Sterba, H. (1980) Zur ansprache der bestandsdichte. *Allgemeine Forstzeitung*, 91, 155–157.
- Diamond, J.M. (1975) The island dilemma: lessons of modern biogeographic studies for the design of natural reserves. *Biological Conservation*, 7, 129–146.
- Drew, G.S. (1995) *Winter habitat selection by American marten (Martes americana) in Newfoundland: why old growth?* PhD Thesis. Utah State University, Logan, UT.
- Efron, B. & Tibshirani, R.J. (1993) *An Introduction to the Bootstrap*. Chapman & Hall Inc., New York, NY.
- Franklin, J.F. & Forman, R.T.T. (1987) Creating landscape