

As a library, NLM provides access to scientific literature. Inclusion in an NLM database does not imply endorsement of, or agreement with, the contents by NLM or the National Institutes of Health.

Learn more: [PMC Disclaimer](#) | [PMC Copyright Notice](#)

Author Manuscript

Peer reviewed and accepted for publication by a journal



J Health Econ. Author manuscript; available in PMC: 2019 Sep 1.

Published in final edited form as: *J Health Econ.* 2018 Aug 13;61:134–150. doi: [10.1016/j.jhealeco.2018.07.004](https://doi.org/10.1016/j.jhealeco.2018.07.004)

Shale Gas Development and Infant Health: Evidence from Pennsylvania

[Elaine L Hill](#) *

[Author information](#) [Article notes](#) [Copyright and License information](#)

PMCID: PMC6629042 NIHMSID: NIHMS1503878 PMID: [30114565](https://pubmed.ncbi.nlm.nih.gov/30114565/)

The publisher's version of this article is available at [J Health Econ](#)

Abstract

This research exploits the introduction of shale gas wells in Pennsylvania in response to growing controversy around the drilling method of hydraulic fracturing. Using detailed location data on maternal addresses and GIS coordinates of gas wells, this study examines singleton births to mothers residing close to a shale gas well from 2003–2010 in Pennsylvania. The introduction of drilling increased low birth weight and decreased term birth weight on average among mothers living within 2.5 km of a well compared to mothers living within 2.5 km of a future well. Adverse effects were also detected using measures such as small for gestational age and APGAR scores, while no effects on gestation periods were found. These results are robust to other measures of infant health, many changes in specification and falsification tests. In the intensive margin, an additional well is associated

with a 7 percent increase in low birth weight, a 5 gram reduction in term birth weight and a 3 percent increase in premature birth. These findings suggest that shale gas development poses significant risks to human health and have policy implications for regulation of shale gas development.

Keywords: infant health, shale gas development, air pollution, water pollution, low birth weight

The United States (US) holds large unconventional gas reserves in relatively impermeable media such as coal beds, shale, and tight gas sands, which together with Canada account for virtually all commercial shale gas produced in the world ([IEA, 2012](#)).¹ New technologies, such as hydraulic fracturing and directional drilling, have made it economically and practically feasible to extract natural gas from these previously inaccessible geological formations.² In 2010, unconventional gas production was nearly 60% of total gas production in the US ([IEA, 2012](#)). Natural gas from the Marcellus formation, particularly in Pennsylvania, currently accounts for the majority of this production ([Rahm et al., 2013](#)).³ A recent assessment by The Wall Street Journal estimates that over 15 million Americans live within 1 mile of an oil or gas well drilled since 2000 in 11 of the 33 states where drilling is taking place ([Gold and McGinty, 2013](#)). With this expansion, it is becoming increasingly common for shale gas development to take place in close proximity to where people live, work and play.

The expansion of shale gas development (SGD) in the US has brought with it a national debate that seemingly lacks a consensus over its economic, environmental, health and social implications. There is growing evidence that shale gas development creates jobs and generates income for local residents in the short run ([Allcott and Keniston, 2014](#); [Bartik et al., 2016](#); [Feyrer et al., 2017](#); [Hausman and Kellogg, 2015](#); [Mason et al., 2015](#)). In addition to its economic benefits, many claim that a move to natural gas (and away from petroleum- or coal-based energy) will support U.S. energy independence and national security. Shale gas provides an attractive source of energy because it emits fewer pollutants (e.g., carbon dioxide, sulfur dioxide, nitrogen oxides, carbon monoxide and particulate matter) when burned than coal and other fossil-fuel energy sources per unit of heat produced ([Chen et al., 2017](#)). Globally, the shale boom has improved ambient air quality and displaced coal-based electricity, especially for areas with coal-fired power plants ([Johnsen et al., 2016](#)). However, these benefits may come with local costs associated with drilling activity in communities where it takes place. These costs may include reduced environmental quality through local air pollution ([Colborn et al., 2012](#); [Litovitz et al., 2013](#); [Witter et al., 2013](#)), water contamination ([Warner et al., 2012](#); [Olmstead et al., 2013](#); [Hill and Ma, 2017](#)), increased truck traffic ([Graham et al., 2015](#)) and health. Concerns over perceived ground water contamination have caused a discount of housing prices to compensate for the risk and an approximately \$19 million increase in bottled water purchases in 2010 in response to SGD in Pennsylvania ([Muehlenbachs et al., 2015](#); [Wrenn et al., 2016](#)). This is further supported by a recent

cost-benefit analysis that found substantial environmental costs associated with health damages from air pollution emitted by SGD totaling \$27.2 billion ([Loomis and Haefele, 2017](#)).

In utero exposure to air pollution has been linked to adverse birth outcomes, lower educational attainment, labor market outcomes and future health problems (See [Currie and Schmieder \(2009\)](#); [Currie \(2009\)](#); [Currie et al. \(2014b\)](#) for summaries of this research). In particular, a large literature has linked air pollution (e.g. particulate matter (PM), carbon monoxide (CO), sulfur dioxide (SO₂), nitrogen oxide (NO_x)) from coal-fired power plants with low birth weight, premature birth and infant mortality both within the US and in the developing world.⁴ With natural gas touted as a transition fuel between coal-based electricity and renewable options, infant health is one way to compare costs across alternative options. While coal is undeniably worse than natural gas with respect to resource extraction and energy generation, concerns regarding emissions associated with shale gas should be studied ([Chen et al., 2017](#)).

The impact of shale gas development on health has become the focus of a growing body of literature. To my knowledge, [Hill \(2012\)](#) is the first study to assess the impact of shale gas development on infant health. Concurrent health studies include case studies ([Bamberger and Oswald, 2012](#)), health impact assessments ([McKenzie et al., 2012](#)), toxicological assessments of specific chemicals ([Colborn et al., 2011](#)), self-reported health symptoms ([Ferraro et al., 2013](#)) and studies exploiting administrative records such as birth certificates, hospital records or electronic medical records (EMR) to study asthma, pneumonia, fatigue, migraine, sinus effects, and birth outcomes ([Hill, 2013](#); [McKenzie et al., 2014](#); [Stacy et al., 2015a](#); [Rasmussen et al., 2016](#); [Casey et al., 2016](#); [Tustin et al., 2017](#); [Currie et al., 2017](#); [Whitworth et al., 2017](#); [Peng et al., 2018](#)).⁵ All but one of the infant health studies find a positive association between drilling and poor birth outcomes measured by premature/preterm birth (PTB) or low birth weight (LBW). Due to a lack of consistency in outcomes, proximity, and exposure metrics used, it is challenging to compare findings across these studies.

To assess the impact of shale gas development on infant health, I build a unique database that contains the longitude and latitude of all shale gas wells, the street address (geocoded) of all new mothers, and data on whether the mother's address falls within public water service areas. To define a treatment variable, I exploit both the timing of drilling activity (using the "spud date," or the date the drilling rig begins to drill a well) and the exact locations of well heads relative to residences. I then use as a comparison group mothers who live in proximity to future wells, as designated by well permits. The exact locations of both wells and mothers' residences allow me to exploit variation in the effect of shale gas drilling within small, relatively homogeneous socio-economic groups, and the timing of the start of drilling allows me to confirm the absence of substantive pre-existing differences. Through this method, I am able to provide robust estimates of the impact of maternal exposure to shale gas development

during pregnancy on birth outcomes.

The main results suggest both statistically and economically significant effects on infant health. I find that shale gas development increased the incidence of low birth weight and small for gestational age in the vicinity of a shale gas well by 25 percent and 18 percent, respectively. Furthermore, term birth weight and birth weight were decreased by 49.6 grams (1.5 percent) and 46.6 grams (1.4 percent), on average, respectively and the prevalence of APGAR scores less than 8 increased by 26 percent. Results for premature birth were mixed and sensitive to specification. The difference-in-differences research design, which relies on the common trends assumption, is tested by examining the observable characteristics of the mothers in these two groups before and after development, testing for pretrends in the outcome variables using the sample before drilling, permit dates only, and future wells only, and using a random date to define treatment. The research design is robust to these tests as well as a range of specifications. I examine mobility using the group of mothers with more than one birth and find that there is little evidence of moms moving in response to drilling. I perform a back of the envelope calculation on the costs of these activities using my estimates and the estimated population within 1 mile of drilling from the Wall Street Journal (e.g. 15 million Americans) and estimate that drilling costs more than \$230 million per year in the 11 out of 33 gas producing states. This estimate is likely to be a lower bound given that this assessment doesn't include all states with development and that I use a lower bound estimate of the costs associated with low birth weight.

This paper contributes to the literature using a quasi-experimental design and is a combination of the strengths of both the epidemiologic and economic literature described above. First, I improve upon the epidemiologic literature by employing a difference-in-differences design. In particular, I exploit the exogeneity of drilling conditional on leasing and permitting, which results in statistically homogenous treated and comparison groups. This provides a more stable comparison group than in [Currie et al. \(2017\)](#) that compares to those living within 3–15km. Second, I improve upon the economics literature by using the strengths of the epidemiologic literature by looking at multiple measures of adverse infant health outcomes which may be indicative of different aspects of drilling exposure. Preterm birth is indicative of preterm premature rupture of membranes, which can result from genetics, stress or low socio-economic status (SES) ([Goldenberg et al., 2008](#)). Low birth weight and small for gestational age (SGA) are more related to intrauterine growth restriction (IUGR), which is more consistently related to air pollution ([Stieb et al., 2012b](#); [Sun et al., 2015](#); [WHO, 2005](#)). Congenital abnormalities indicate exposure to a teratogen during pregnancy. Given the inconsistency in measured outcomes in existing studies, I simultaneously estimate impacts for all outcomes within the same sample and identification strategy. This is particularly useful for policy given the mixed findings in the existing studies and that none of these studies directly test exposure mechanisms. Third, I improve upon the economics literature by thoroughly controlling for predictors of infant health and estimating the extensive and

intensive margins of drilling. I include controls for insurance status, WIC, previous risky pregnancy, parity, and smoking status. I also measure heterogeneity across SES subgroups and test whether moms are moving in response to drilling. Importantly, I contribute to the literature by measuring the effect of an additional well on birth outcomes, which is perhaps more relevant to policy-making than simple binary measurements of exposure.

The rest of the paper proceeds as follows: [section I](#) presents background and context and [section II](#) describes the data. [Section III](#) presents graphical evidence and [section IV](#) describes the estimation strategy. Sections [V](#) and [VI](#) presents results and robustness checks. Section VII provides interpretation and discussion of the results. [Section VIII](#) concludes.

I. Background

I.1. A Brief Shale Gas Overview for Pennsylvania

In Pennsylvania, shale gas development involves primarily high-volume hydraulically fractured horizontal wells drilled into the Marcellus Shale and more recently, the Utica Shale. Hydraulic fracturing is a process to stimulate a well that uses water to fracture the rock or shale beneath the ground. On average, in Pennsylvania, it involves injecting approximately 4–8 million gallons of water mixed with sand and fracturing chemicals into the well and using pressure to fracture the shale about 6,500–7,500 ft below the surface ([Chen and Carter, 2016](#)). Shale plays are heterogeneous and so the distance drilled and quantity of water required differ across varied geological formations. The entire process of completing a natural gas well takes, on average, 3–9 months to finish: access road and well pad construction occurs for a month (0–4 weeks) prior to the spud date, drilling the well takes about 30 days (vertical drilling for 0–2 weeks and horizontal drilling for 4–8 weeks), preparation for hydraulic fracturing takes 1–2 months, hydraulic fracturing takes about 7 days, flowback occurs for 2–8 weeks and clean up and testing takes about a month before the well goes into production ([Casey et al., 2015](#); [Graham et al., 2015](#)). During the first few months, diesel trucks bring in materials required for the drilling process, averaging 1500–2000 truck trips per well completion in Pennsylvania. During the first 30 days after well stimulation, it is estimated that approximately 30–70% of the water used during the drilling process returns to the surface (called flowback) and is collected in ground level water impoundments and then taken to be treated at a waste water facility ([Kondash et al., 2017](#)).

Most wells are drilled on private property that has been leased to oil and gas companies.¹ After the land is leased by the mineral owner, a company applies for a permit to drill on that property. The state government approves permits and once a company has a permit, the drilling often commences quickly

thereafter. There are many layers of decision-making independent of the mineral owner that determine exactly which leases become permits and which permits become a well. This research uses only those locations that are permitted by the state to reduce selection bias in the estimates that follow.

The identification strategy used in this paper depends on the assumption that drilling is exogenous relative to locations that are permitted but not yet drilled. However, areas that are permitted but not drilled may be different from areas that experience active drilling. For example, areas without active drilling may not have as many property owners willing to lease mineral rights or the industry may prioritize leasing in areas with the most productive shale. Appendix Figure A1 overlays the parcels with leases from Drillinginfo with the strata of shale depth from EIA. For counties where we have lease data, the extent of leasing is densest along the deepest contours and more sparse along the shallower contours, except in the northeastern part of the state such as Bradford County. To examine this further, I linked the lease and depth data to the wells and permits used in these analyses to test whether there are substantial differences.⁷ There are no differences in leasing defined by the proportion of acres leased within Census block groups between permitted and drilled wells. The average Census block group in the data is 40 percent leased for both permitted and drilled locations. In the top 10 drilled counties, this jumps to 60 percent, but is again the same across permitted and drilled locations. Permits that are drilled seem to be explained by shale depth as opposed to some difference in community preference as proxied for by leasing activity.

I.II. Shale Gas Development As A Potential Pollution Source

Preliminary evidence indicates that shale gas development may produce waste that could contaminate the air, aquifers, waterways, and ecosystems that surround drilling sites or areas where water treatment facilities treat the waste water from the drilling process. Below I review the current state of the scientific evidence.

I.II.1. Water Pollution

There are a number of mechanisms by which shale gas development might contaminate ground and surface water sources and thereby impact either public or private drinking water. According to a recent assessment by EPA, these mechanisms include: spills of hydraulic fracturing (HF) fluids prior to mixing with large quantities of water or produced water after hydraulic fracturing has taken place, injection of hydraulic fracturing fluids into wells with inadequate mechanical integrity (e.g. faulty well casings), injection of HF fluids directly into groundwater sources, discharge of inadequately treated hydraulic fracturing wastewater to surface water, and disposal or storage of hydraulic fracturing wastewater in

unlined pits ([EPA, 2016](#); [Osborn et al., 2011](#); [Jackson et al., 2013](#); [Olmstead et al., 2013](#); [Warner et al., 2013](#)).⁸ The EPA report identified 1,084 chemicals reported to be used in hydraulic fracturing fluids and 599 chemicals detected in produced water ([EPA, 2016](#)). Of the 599 chemicals detected in produced water, only 77 were also reported to be used in hydraulic fracturing fluid– which is not a great match. The report found that chemicals used in HF fluid varied greatly across regions, which limits external validity ([EPA, 2016](#)).⁹ [Elliott et al. \(2017\)](#) provides a review of these chemicals for reproductive and developmental toxicity.¹⁰

The lack of reliable information about what chemicals are used leaves the scientific community testing many different chemicals across regions, with little overlap among detected chemicals. Studies of groundwater contamination have primarily used private drinking water wells and assessed proximity to shale gas wells to assess contamination (e.g. within 5 km of gas wells versus larger distances) ([Hildenbrand et al., 2016](#); [Osborn et al., 2011](#); [Jackson et al., 2013](#)). Studies have found increases in organics (many naturally occurring such as chlorides, bromides and iodides, arsenic, selenium, manganese, strontium, barium, heavy metals, beryllium), volatile and semivolatile organic compounds (e.g. BTEX, 2-Butanone), diesel range organic compounds, solvents (e.g. methanol, dichloromethane), and methane ([Drollette et al., 2015](#); [Hildenbrand et al., 2015, 2016](#); [Yan et al., 2016](#); [Alawattegama et al., 2015](#); [Burton et al., 2016](#)). Some studies have not found any evidence of contamination, leaving whether SGD impacts water quality a hotly debated question ([Li et al., 2016](#)). One study assessing groundwater-sourced public water systems' water quality found that SGD wells were associated with an increase in SGD-related chemicals for wells drilled within 1 km of the groundwater source ([Hill and Ma, 2017](#)).

Surface water impacts are more likely to be associated with the handling of shale gas waste. Waste water treatment and discharge is associated with elevated levels of barium, strontium, bromides, chlorides, benzene, and total dissolved solids exceeding the maximum contaminant level for drinking water ([Olmstead et al., 2013](#); [Vengosh et al., 2014](#); [Hladik et al., 2014](#); [Lester et al., 2015](#); [Ferrar et al., 2013](#)). Treated produced water (containing naturally occurring bromide and iodide) are potential sources of toxic disinfection byproducts (DBPs): iodinated trihalomethanes (THMs) and brominated haloacetonitriles (HANs) in surface water ([Parker et al., 2014](#)).¹¹ Endocrine disrupting chemicals measured in surface water near waste effluent in Colorado and West Virginia are of concern for reproductive health ([Kassotis et al., 2015](#)).

I.II.2. Air Pollution

Despite less attention in the media, air pollution is gaining more recent attention by researchers. All stages of shale gas development have the potential to produce hazardous air pollution emissions

([Kargbo et al., 2010](#); [Schmidt, 2011](#)). Air pollution has become a more immediate concern following studies in Colorado that discovered higher levels of volatile organic compounds (VOCs), methane and other hydrocarbons near drilling sites ([Colborn et al., 2012](#); [Pétron et al., 2012](#)). Other emissions associated with combustion include particulate matter, poly-cyclic aromatic hydrocarbons, sulfur oxides and nitrogen oxides ([Colborn et al., 2012](#)). More recent studies have also assessed the air pollution contribution of the many truck trips necessary to build and fracture a well ([McCawley, 2017](#); [Goodman et al., 2016](#)).

Studies of air pollution in Pennsylvania are suggestive of increased emissions associated with shale gas development, but have produced inconsistent results. For example, the Pennsylvania Department of Environmental Protection (PA DEP) has conducted three short-term (1 week) air pollution studies in three regions of the state but found little evidence of air pollution concentrations that would likely trigger air-related health issues associated with Marcellus Shale drilling activities ([PADEP, 2010b, 2011b, a](#)). But the air emissions inventory for the unconventional natural gas industry, starting in 2011, indicates modest emissions of CO, NO_x, PM₁₀, SO_x and VOCs ([PADEP, 2013a](#)).¹² These results were verified by a recent RAND study that used the PA DEP data and other sources to estimate the emissions from shale gas in Pennsylvania ([Litovitz et al., 2013](#)). The most significant pollutants, according to the authors, were NO_x and VOCs, which were equivalent to or larger than some of the largest single emitters in the state and the low-end estimates of nitrogen oxide emissions were 20–40 times higher than the level that would be defined as a “major” emissions source. During the same time period, due to the conversion of electricity from coal to natural gas in the state, the overall pollution for all the criteria pollutants measured decreased substantially and more than outweighed the new pollution related to shale gas development. These data, however, indicate a more nuanced picture of air emissions from drilling activities and show that shale gas development is now a significant source of air pollution in rural counties with few other point-sources of pollution. For example, the 2,600 tons and 2,440 tons of shale-related NO_x emitted in Bradford County and Susquehanna County, respectively in 2011 make up one-third of the statewide shale-related NO_x of 16,500 tons ([PADEP, 2013b](#)). These levels surpass the single largest industrial source of NO_x pollution in the 11-county northeast region, a coal-fired power plant in Northampton County that emitted 2,000 tons in 2011 ([Legere, 2013](#)).

As mentioned above, Pennsylvania DEP began requiring companies drilling Marcellus shale gas wells to report annual estimates of air emission to an inventory starting in 2011. In [Table 1](#), I estimate the intensive margin of the number of wells in a zip code on the annual tons of each pollutant aggregated to that zip code from 2011 to 2015. I also estimate tertiles of wells to capture intensity. Each additional well contributes an average of 0.5 tons of CO, 2 tons of NO_x, 0.07 tons of PM_{2.5}, 0.03 tons of SO_x, and 0.17 tons of VOCs per year. The average zip code in 2011 experienced 14 tons of CO, 41 tons of NO_x, 1.4 tons of PM_{2.5}, 0.5 tons of SO_x, and 8 tons of VOCs. In the subset of wells that were spudded prior to

2011, the average well produced 2 tons of CO, 4.7 tons of NO_x, 0.14 tons of PM_{2.5}, 0.04 tons of SO_x, and 0.63 tons of VOCs in 2011. The top tertile (14–213 wells) of zip codes experience an average of 28 tons of carbon monoxide (CO), 90 tons of NO_x, 2.6 tons of PM_{2.5}, 1.8 tons of SO_x, and 9 tons of volatile organic compounds (VOC) per year. Babies exposed to shale gas development within 10 km face an average of 24 wells (max of 240) in 2010 and is fairly similar to the tertiles used in [Table 1](#) Although there isn't a direct way to measure the contribution of these emissions to ambient air quality, they do represent a modest and potentially significant amount of emissions for these rural areas.

Table 1:

Pollution Per Well and Tertiles Aggregated to Zip Code 2011–2015

	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)
	CO	CO	NO _x	NO _x	PM _{2.5}	PM _{2.5}	SO _x	SO _x
# wells	0.526*** (0.0567)		2.048*** (0.171)		0.0639*** (0.00543)		0.0325*** (0.00500)	
3–5 wells		3.271 (2.928)		13.28 (9.060)		0.395 (0.289)		0.514* (0.256)
6–13 wells		13.30*** (3.305)		45.02*** (10.23)		1.472*** (0.326)		1.271* (0.289)
14–213 wells		27.47*** (3.934)		89.83*** (12.17)		2.630*** (0.388)		1.777* (0.344)
log prod	0.552*** (0.193)	0.443** (0.202)	1.806*** (0.580)	1.627*** (0.626)	0.0633*** (0.0185)	0.0598*** (0.0200)	0.0336** (0.0170)	0.0241 (0.017)
Observations	1,172	1,172	1,172	1,172	1,172	1,172	1,172	1,172
R-squared	0.697	0.688	0.730	0.707	0.724	0.699	0.500	0.494
Dep. Var	15.35	15.35	45.85	45.85	1.482	1.482	0.507	0.507
Mean								

[Open in a new tab](#)

Notes: Data are from the PA DEP Air Emissions Inventory for Unconventional Natural Gas Operations 2011–2015. Units are tons/year. Emissions are aggregated to zip code-year. Regressions include year and zip code fixed effects. First column for each pollutant is number of reported wells in that zip code-year. Second column provides tertile estimates. Significance:

* $p < 0.10$,
** $p < 0.05$,
*** $p < 0.01$.

Of interest is whether wells continue to produce emissions after drilling and entering into production. To test this, I estimate the amount of reported emissions per year per pollutant using years since spud date as the regressors for all wells reported in the emissions inventory from 2011–2015 ([Appendix Table A1](#)). For the most part, emissions are largest for the year of the spud date and the first year after drilling occurred, but emissions continue for most pollutants out to years 4 or 5. Due to this evidence, I estimate models using wells drilled from 2006–2010 and determine exposure by wells drilled prior to birth as opposed to restricting just to drilling activity during gestation.

I.III. Pollution and Health Literature

[Stillerman et al. \(2008\)](#) review the epidemiological literature and find associations between low birth weight and maternal exposures to PM, SO₂, CO, NO_x, VOCs and ozone. Most of the studies cited looked at these pollutants in isolation, but with shale gas development mothers are likely exposed to many at the same time and there is little research that examines any compounding effects.¹³ All of the air pollutants emitted by shale gas development described above have been associated with adverse birth outcomes (see Online Appendix for more detail). Unfortunately, many of the epidemiological studies do not take into account socio-economic status and so the observed relationships could reflect unobserved factors that may be correlated with pollution and infant health outcomes (i.e. urban areas). The epidemiological literature relating water pollution to reproductive health is more limited (see [Quansah et al. \(2015\)](#) and [Nieuwenhuijsen et al. \(2013\)](#) for recent reviews).

There is a growing literature within health economics that addresses the most common air pollutants associated with SGD described above utilizing quasi-experimental designs and rich controls for potential confounders to identify the infant health effects of ambient air pollution. See [Currie et al. \(2014b\)](#) for a review of the economics literature on short and long term impacts of early life exposure to pollution. For example, [Currie and Walker \(2011\)](#) estimate that reductions in air pollution from E-Z Pass result in reductions of low birth weight (LBW) between 8.5–11.3 percent and [Zahran et al. \(2012\)](#) utilize the natural experiment of benzene content in gasoline from 1996 to 1999 in the US and found exposure to benzene reduces birth weight by 16.5 g and increases the odds of a very low birth weight event by a multiplicative factor. [Lavaine and Neidell \(2013\)](#) use the natural experiment of a strike that affected oil refineries in France to explore the temporary reductions in SO₂ and find that the reductions increased birth weight by 75 grams, on average (2.3 percent increase) and reduced low birth weight by

2 percentage points for residences within 8 km of the air pollution monitor.

With natural gas touted as a transition fuel between coal-based electricity and renewable options, infant health is one way to compare costs across alternative options. To date, even within the epidemiological literature, studies of the effects of living near coal mining (underground or mountain top) on birth outcomes are extremely limited. All three studies focus on WV: one found an increased risk of low birth weight (16 percent increase in most intensive areas) and one study found an increased risk of congenital anomalies with mountain top removal mining associated with worse outcomes, but was later refuted by the third study when the authors controlled for hospital of birth ([Ahern et al., 2011b, a](#); [Lamm et al., 2015](#)). See [Hendryx \(2015\)](#) and [Boyles et al. \(2017\)](#) for systematic reviews of the public health literature. However, recent papers in the economics literature have exploited plant openings and closings or being downwind from a plant to identify the causal impact of coal-fired power plants on infant health and have found adverse birth outcomes: a 5 percent reduction in continuous birth weight as the grid transitioned from nuclear to coal in Tennessee ([Severnini, 2017](#)), a 6 percent increase in low birth weight for infants 20 miles downwind of a power plant ([Yang et al., 2017](#)), 15 percent decreased risk for low birth weight once the plant closed ([Yang and Chou, 2017](#)), and 3,500 infant deaths per year as of 1962 associated with the expansion of the power grid between 1938 and 1962 ([Clay et al., 2016](#)). A recent paper focused on storage of coal at power plant locations found that a 10 percent increase in PM2.5 from coal storage increased infant mortality rates by 6.6 percent ([Jha and Muller, 2017](#)).

I.III.1. SGD and Health Literature

Most of the studies to date that address potential health impacts of shale gas development measure pollutants at drilling sites or in drilling fluids and then identify the health implications based upon expected exposure to these chemicals (e.g. toxicological assessment). For example, [Colborn et al. \(2011\)](#) find that more than 75% of the chemicals could affect the skin, eyes, and other sensory organs, and the respiratory and gastrointestinal systems. Chronic exposure is particularly concerning because approximately 40–50% could affect the brain/nervous system, immune and cardiovascular systems, and the kidneys; 37% could affect the endocrine system; and 25% could cause cancer and mutations. These may have long-term health effects that are not immediately expressed after a well is completed. Recent studies have found increased hospitalizations for cardiac conditions ([Jemielita et al., 2015](#)), increased risk of three types of asthma measures ([Rasmussen et al., 2016](#)), increased risk of hospitalization for pneumonia ([Peng et al., 2018](#)), and increased prevalence of fatigue, migraine and sinus effects for residents living near development ([Tustin et al., 2017](#)).

A growing body of literature has attempted to address the potential reproductive health effects of shale

gas development. All of these studies are retrospective analyses of birth certificate records or electronic medical record data and focus on proximity to maternal residences as the definition of “exposure.” In Colorado, [McKenzie et al. \(2014\)](#) find an increased risk of congenital heart defects with the highest quartile of exposure compared with the absence of any gas wells within a 10-mile radius of the maternal residence. They also found a reduction in premature birth and low birth weight for the highest quartile of exposure. [Hill \(2013\)](#) finds an increase in the latter two measures of around 30 percent for oil, natural gas and coalbed methane wells. Using a similar research design in Texas, [Whitworth et al. \(2017\)](#) finds an increase in premature birth of 14 percent and an increase in fetal death upwards of 50 percent. Using a case-control analysis, Whitworth et al. (2199) find a 20 percent increase and 15 percent increase in preterm birth for any wells and producing wells within 0.5 miles of the maternal residence, respectively.

Focusing on the three studies in Pennsylvania, [Stacy et al. \(2015a\)](#) study three counties in Southwestern Pennsylvania from 2007–2010 and [Casey et al. \(2016\)](#) study two hospitals in the Geisinger Health System from 2009–2013.¹⁴ [Currie et al. \(2017\)](#) study birth records from Pennsylvania from 2004–2013. [Stacy et al. \(2015a\)](#) use inverse distance weighted number of wells within 10 miles of the maternal residence and create quartiles to define exposure (compare 4th to 1st quartiles; omitting mothers with no wells within 10 miles). [Casey et al. \(2016\)](#) create an “activity index” and use quartiles of the index (compare 4th (average 124 wells, median 8) to 1st quartile (average 6 wells, median 0), but include those with no wells within 20 km).¹⁵ [Currie et al. \(2017\)](#) utilize a difference-in-difference study design comparing close (e.g. 0–1, 1–2, 2–3km) versus further away (e.g. all PA or 3–15km) in Pennsylvania using county fixed effects. [Stacy et al. \(2015a\)](#) find a reduction in birth weight and an increase in small for gestational age (SGA) of 34 percent. [Casey et al. \(2016\)](#) find an increase in premature birth that ranges from 40 to 90 percent and an increase in the prevalence of risky pregnancies. [Currie et al. \(2017\)](#) find a 25 percent increase in low birth weight for the 0–1km group. The 2–3km buffer suggests a 16 percent increase in low birth weight. The 1–2km buffer is not as consistent or statistically precise as the 0–1 or 2–3km buffers. Other measures studied include continuous birth weight and a health index. [Currie et al. \(2017\)](#) further estimate their models using maternal fixed effects but these models are not statistically significant, nor are they consistent with all of their primary findings.


In the discussion section ([Section VII](#)), I compare and contrast my results with those cited above and also provide discussion of interpretation.

II. Data

My analysis is based upon a data set acquired from the Pennsylvania Department of Environmental

Protection (PA DEP) that contains GIS information for all of the wells drilled in the state of Pennsylvania since 2000 and define whether it is a Marcellus shale well. For the analysis that follows, the spud date (date when the drilling rig begins drilling the well) is used as the temporal identification of treatment. In total, the analysis uses 2,459 natural gas wells spudded between 2006 and 2010. In addition to the existing gas well data, this study also makes use of the permit data on the PA DEP website. This allows for the identification of permits that do not become a well during the sample time frame; approximately 40 percent of permits do not become a well (author calculation from PA DEP data). This information is used to define a potential control group for those infants born to residences close to existing gas wells. The assumption is that these residences are a potential counterfactual group: those who have the potential to live close to a gas well in the future, but have not yet had a well drilled as of the timing of the data collection. [Figure 1](#) shows drilled and permitted wells through 2010 along the strata of shale depth. For the most part, wells that are drilled are clustered along the deepest shale strata and permitting is more random.

Figure 1:



[Open in a new tab](#)

Map of Shale Gas Development and Permitting through 2010

My second source of data comes from restricted-access vital statistics natality and mortality data from Pennsylvania for the years 2003 to 2010. The restricted-access version of these birth certificate records contain residential addresses geocoded to latitude and longitude and unique identifiers for the mother, father and infant. This precision is essential to my identification strategy because the consequences of drilling are highly localized. To construct the analysis data set, I combine the spatially identified wells and maternal residences and calculate proximity to the nearest wells.

The vital statistics contain important maternal characteristics such as race, education, age, marital status, WIC status, insurance type, previous risky pregnancy and whether the mother smoked during her pregnancy. In the empirical analyses that follow, I control explicitly for these, as well as month of birth, year of birth, the interaction, and gender of the child.¹ I exclude multiple births in all analyses because plural births are more likely to have poor reproductive health independent of exposures to environmental pollution.

I focus on low birth weight (LBW), premature birth and term birth weight (TBW) as the primary outcomes of interest. Low birth weight, defined as birth weight less than 2500 grams, and premature birth, defined as gestation length less than 37 weeks, are commonly used as key indicators of infant health and have been shown to predict adult health and well-being.¹ I also present the continuous measure of term birth weight, defined as birth weight for infants who reach full term at 37 weeks gestation, to study whether there is an average effect on the birth weight distribution as opposed to these more extreme health outcomes. Other birth outcomes that I examine include the continuous measure of birth weight, gestation (measured in weeks), small for gestational age (SGA; defined as 10th percentile of weight distribution for the gestational week of birth), an indicator for whether the APGAR score is less than 8 to predict an increased need for respiratory support, congenital anomalies, an infant health index and infant mortality (death in the first year).¹⁸

[Table 2](#) provides summary statistics for the universe of births in Pennsylvania from 2003-2010. The first column reports characteristics of all births and the second column reports average characteristics of births for mothers' residences within 2.5 km of where a shale gas well has been drilled or will be drilled. The localized data I use in this analysis is actually quite similar to the characteristics of the rest of the state. Mothers who live close to shale gas development are less likely to be African American and Hispanic, slightly better off in terms of health outcomes, younger, better educated and more likely to be married at the time of birth compared with the state average. The mothers in the analysis sample are also more likely to smoke than the average for the state. Columns (3) and (4) provide summary statistics for the primary difference-in-difference (DD) analysis sample; the sample is restricted to those mothers' residences within 2.5 km of a gas well or permit and I compare residences before and after drilling. Most of the statistically significant differences between these two samples are arguably not very economically important. Mothers with infants born after drilling are less likely to be over the age of 35, more likely to receive WIC, and more likely to receive Medicaid, on average, likely to do with the shale gas boom coinciding with the Great Recession. However, [Table 3](#) suggests no changes in these economic variables after shale gas development.¹⁹

Table 2:

Summary Statistics by Sample

	All Births	Residences within 2.5 km of well			T-Stat
		Total	Before	After	for difference
Characteristics of birth					
Birth weight (grams)	3321	3340	3343.23	3310.30	2.70**
Term birth weight (grams)	3407	3415	3418.39	3383.15	3.30***
Gestation in weeks	38.77	38.76	38.76	38.71	1.33
Premature	0.08	0.08	0.076	0.077	−0.09
Low birth weight (LBW)	0.07	0.06	0.055	0.063	−1.52
Small for gestational age (SGA)	0.11	0.10	0.098	0.106	−1.25
APGAR 5 minute	8.81	8.89	8.886	8.885	0.07
Female	0.49	0.49	0.485	0.495	−0.95
Mother’s Characteristics					
Drop Out	0.164	0.113	0.112	0.118	−0.88
High School	0.270	0.296	0.297	0.288	0.93
Some college	0.260	0.299	0.299	0.293	0.64
College plus	0.298	0.290	0.289	0.299	−1.07
Teen Mom	0.057	0.048	0.047	0.049	−0.34
Mom Aged 19–24	0.265	0.268	0.267	0.274	−0.65
Mom Aged 25–34	0.527	0.547	0.545	0.559	−1.31
Mom Aged 35 and older	0.150	0.137	0.140	0.117	3.03**
Mom Black	0.156	0.025	0.025	0.024	0.15
Mom Hispanic	0.092	0.011	0.011	0.010	0.57

	All Births	Residences within 2.5 km of well			T-Stat
		Total	Before	After	for difference
Married at time of birth	0.575	0.632	0.633	0.626	0.71
Mom Smoked While Pregnant	0.227	0.299	0.299	0.300	-0.13
Received WIC	0.385	0.398	0.395	0.427	-2.94**
Medicaid	0.272	0.326	0.320	0.376	-5.45***
Private Insurance	0.576	0.567	0.569	0.549	1.84
Wells within 2.5 km					
# of wells before birth	0.000	0.333	0.000	2.89	-19.30***
# of wells during gestation	0.000	0.188	0.000	1.714	-93.13***
Observations	1098884	21610	19246	2364	

[Open in a new tab](#)

Notes: The samples described here include only singleton births.

Significance:

* $p < 0.10$,

** $p < 0.05$,

*** $p < 0.01$.

Table 3:

Post- Drilling Differences in Average Characteristics of Mothers Close to Wells

	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)
	Teen Mom	Dropout	Black	Smoked	WIC	Medicaid	Born PA	Mov
	Differences in characteristics for analysis sample using DD estimator							
Within 2.5 km_* post- drilling	0.000550	-0.0132	0.00343	0.00277	-0.00501	-0.0204	-0.0222	0.01
	(0.00666)	(0.0118)	(0.00308)	(0.0196)	(0.0246)	(0.0282)	(0.0163)	(0.01
Observations	21646	21646	21646	21646	21469	21646	21646	215
R ²	0.012	0.039	0.016	0.026	0.061	0.078	0.020	0.04
Pre-drilling Mean	0.0496	0.117	0.0243	0.307	0.404	0.323	0.815	0.07

[Open in a new tab](#)

Notes: Each coefficient is from a different regression. Pre-drilling (post-drilling) refers to births that occur before (after) the spud date of the closest well. Robust standard errors are clustered at the mother's residence county. All regressions include indicators for month and year of birth, birth*year and residence county fixed effects.

Significance:

* p<0.10,


** p<0.05,

*** p<0.01.

III. Graphical Evidence

If living close to a drilled well has a negative impact on infant health, we should see average prevalence of low birth weight for mother's residences in close proximity to wells increase subsequent to when drilling begins. Moreover, we should observe larger impacts for homes closest to drilling activity (e.g. dose response). [Figure 2](#) shows the low birth weight (LBW) and premature birth gradients of distance to closest well before and after drilling. LBW prevalence is on average higher for those residences close to drilled wells, compared with those who are close to permitted wells. The primary effect appears to be within 2.5 km but persists out to almost 5 km (consistent with regression results). In contrast, we do not see a clear trend in premature birth over distance (regression results are mixed depending on extensive or intensive measures).

Figure 2:




[Open in a new tab](#)

Distance Gradients of Infant Health by Nearest Well Results from a local polynomial regressions of low birth weight on distance from closest well's future/current location or on days before/after spud date. Observations within 5 km of a well.

In [Figure 3](#), I explore pre-trends in these two outcomes across treatment (e.g. drilled wells) and control (e.g. permitted wells) groups, which addresses the validity of my difference-in-difference design. Prior to drilling in 2008, trends appear parallel and indicate a diverging trend once drilling begins.

Figure 3:




[Open in a new tab](#)

Time Trends of Infant Health Within 2.5 km of Drilled and Permitted Wells Results are from a regression with an interaction term for drilled well * year including county, birth month and year fixed effects. Observations are the main difference-in-differences sample or those mothers within 2.5 km of a drilled well or permitted well.

A primary threat to my identification strategy is that the population of mothers may change in response to drilling. One way to test this is to graph the gradient in observable maternal characteristics. In [Figure 4](#), I graph this gradient out to 20 km.²⁰ The gradient is very similar within 5 km of the nearest gas well before and after drilling. If anything, moms after drilling may be more college educated, which is consistent with my regression results. However, the characteristics change meaningfully beyond 5 km, and moms who live more than 5 km from a gas well before or after drilling are more likely to be college educated, less likely to have their birth paid for by Medicaid, less likely to participate in WIC and less likely to smoke. This suggests selection into living very close to drilling/future drilling and that those who live closer may have lower SES than those who live 15–20 km away. This could drive adverse outcomes related to living very close to drilling, which is why I use permitted locations that are similarly close to mothers' residences since these groups are more homogeneous and statistically similar.

Figure 4:



[Open in a new tab](#)

Distance Gradients of Maternal Characteristics by Nearest Well Distance bins are 0.5 km, smoothed using “lpoly” (degree 0, bandwidth 15).

IV. Empirical Strategy

I exploit the variation over time and across space in the introduction of shale gas wells in Pennsylvania during 2003–2010. Combining gas well data and vital statistics allows the comparison of infant health outcomes of those living near a gas well and those living there before drilling began. Rather than compare aggregated areas, I know specific locations where shale gas drilling has taken place and the

dates of when drilling began. The specific location data allow me to compare reproductive health within very small areas in which mothers are likely to be more homogeneous in observable and unobservable characteristics than in aggregate comparisons.

Relying on cross-sectional variation alone, however, would be problematic if mother characteristics vary within the small radius of interest that are unobservable to the researcher. If, for example, the location of gas drilling occurs where the neighborhoods are already economically distressed, then the variation in health outcomes may reflect socio-economic status, as opposed to living in close proximity to shale gas development. I therefore examine localized reproductive health outcomes before and after shale gas development exploiting permitted but not-yet-drilled wells as a comparison. I use 2.5 km (approximately 1.5 miles) as the primary distance of interest for the main specifications that follow due to my graphical analyses as well as due to the precision of the effect at this distance for robustness checks.²¹

My primary model is a difference-in-difference model – in which mothers living within 2.5 km from a shale gas well or permit before drilling are used as a control for those exposed after drilling began – to estimate the impact of exposure to shale gas development on birth outcomes. Thus, the counterfactual change in infant health for mother's residences close to a shale gas well is estimated using births prior to drilling at the same distance from the well bore location or permitted location (e.g. those permits that become a well by 2011 are treated differently than those permits that are not drilled by 2011). These models take the following form:

$$Outcome_{it} = \beta_1[Well \leq X]_{it} + \beta_2[Post]_{it} + \beta_3[Well \leq X]_{it} * [Post]_{it} + \beta_4 X_{it} + \gamma_t + \chi_c + \epsilon_{it} \quad (1)$$

where $Outcome_{it}$ is either low birth weight, prematurity and other measures of reproductive health for each infant i born in month-year t . $[Well \leq X]_{it}$ is either an indicator for any gas well or the number of gas wells within X km of the mother's residence. $[Post]_{it}$ is an indicator for whether the birth occurs after the spud date of the nearest well of the maternal residence. The estimated impact of shale gas development on infant health is given by the coefficient β_3 and is the difference-in-differences estimator comparing before and after drilling holding the distance X km fixed for wells, future wells and permits.²² The vector X_{ict} contains mother and child characteristics including indicators for whether the mother is African American, Hispanic, four mother education categories (less than high school (left out category), high school, some college, and college or more), mother age categories (teen mom (left out category), 19–24, 25–34 and 35+), indicators for smoking during pregnancy, an indicator for receipt of Women, Infants, and Children (WIC), three health care payment method categories (Medicaid, private insurance, and self-pay), mother's marital status, parity, previous risky pregnancy

and an indicator for sex of the child. Indicators for missing data for each of these variables were also included. γ_t are indicators for the year, month and year*month to allow for systematic trends. χ_c are indicators for each mother's county of residence. Standard errors are clustered at the county.²³

V. Results

V.I. Differences in Characteristics of Mothers Close to a Well

To test the validity of my research design, I estimate [equation \(1\)](#) and use the difference-in-difference estimator to see if there are any changes in mother characteristics after drilling began well would face similar ex ante conditions as those born close to a permit that did become a well during the period I have gas well data for (2003–2011). Infants born to mothers who reside close to potential wells are likely to be the most similar comparison group when it comes to family, geological formation and community characteristics. The decision for which permits become a well is arguably exogenous to the families in these locations. This should account for both observable characteristics, as well as unobservable characteristics, such as economic factors that promote gas drilling in a community and the unobserved geology of the shale underneath these communities. I test these assumptions and do not find any observable differences in the characteristics of mothers who live close to a future well versus a permitted and not yet drilled well.

(e.g. replace birth outcomes with indicators for maternal characteristics). In [Table 3](#): Panel B, I do not find any indication that maternal characteristics are changing in response to shale gas development. In [Appendix Table A2](#), I show that there are no statistically significant differences in maternal characteristics for any potential proximities (e.g. 2km-3.5km).

V.II. The Impact of Shale Gas Development on Birth Outcomes

[Table 4](#) shows the results from estimating ([equation 1](#)) on low birth weight, term birth weight and premature birth. Distance to a well, including future and permitted, is held fixed at 2.5 km for these models. Each coefficient represents an estimate of β_3 – the difference-in-difference estimator – from a separate regression. Columns (1), (3) and (5) show a model that controls only for month and year of birth, month*year and county fixed effects. Adding controls for observable characteristics of the mother should only reduce the sampling variance while leaving the coefficient estimates qualitatively unchanged. Columns (2), (4) and (6) add maternal characteristics and show that controlling for maternal characteristics has little effect on the estimated coefficients for low birth weight and term birth weight. I find a statistically significant increase in low

birth weight of 1.36 percentage points and a reduction in term birth weight of 49.58 grams, on average. I do not find any statistically significant effect for premature birth. Thus, mothers who give birth after drilling are more likely to have reduced weight babies, but they come to term. This difference indicates an overall increase in low birth weight of 24 percent (base of 5.7 percent) and a decrease in term birth weight of 1.5 percent (base of 3416 grams), on average.^{[25](#)}

Table 4:

Impact of Well Location on Birth Outcomes

	(1)	(2)	(3)	(4)	(5)	(6)
	Low Birth	Weight	Term Birth	Weight	Premature	
Within 2.5 km [*] post- drilling	0.0144 ^{**} (0.00537)	0.0136 ^{**} (0.00511)	TM 47.82 ^{***} (15.12)	TM 49.58 ^{***} (14.04)	0.00118 (0.00597)	0.000354 (0.00664)
Observations	21610	21610	19978	19978	21,189	21,189
R-squared	0.008	0.021	0.013	0.075	0.008	0.012
Pre-drilling Mean	0.057	0.057	3416	3416	0.079	0.079
Maternal Characteristics	No	Yes	No	Yes	No	Yes

[Open in a new tab](#)

Notes: Each coefficient is from a different regression. The sample is limited to singleton births and to the sample with a well/permit within 2.5 km. All regressions include indicators for month and year of birth, month*year, residence county indicators, an indicator for drilling before birth (defined by closest well), an indicator for residence within 2.5 km of a well or future well and the interaction of interest of Within 2.5km*post-drilling. Maternal characteristics include mother black, mother Hispanic, mother education (hs, some college, college), mother age (19–24, 25–34, 35+), female child, WIC, smoking during pregnancy, marital status and payment type (private insurance, medicaid, selfpay, other). Indicators for missing data

Significance:

* $p < 0.10$,

** $p < 0.05$,

*** $p < 0.01$.

The results are qualitatively similar when I estimate [equation \(1\)](#) for other distances up to 5 km from a gas well or permit (See [Appendix Table A3](#)). As the buffer of exposure expands, the point estimates become smaller, indicating a dose response relationship, with effects dissipating beyond 3.5 km. The advantage of using permits as the counterfactual is that I can look at only residences that are going to be very close to gas wells at some point in the observable future, which should account for the economic benefits for households receiving lease royalties from the industry.²⁶

[Table 5](#) presents estimates of ([equation 1](#)) for changes in birth weight, 5 minute APGAR scores less than 8, gestation (weeks), small for gestational age (SGA), congenital anomaly, and an index for infant health due to having multiple outcomes of interest.²⁷ As before, each column presents estimates from a separate regression, comparing outcomes before and after drilling at 2.5 km from a well head or permit. I present results with maternal controls due to there being little appreciable difference for the models without these controls (results available upon request). Looking across all reproductive health measures, these estimates are consistent with shale gas development being detrimental to infant health. The introduction of shale gas development reduced birth weight by 46.6 grams (1.4 percent reduction), which is consistent with the findings for term birth weight. Five minute APGAR scores were also affected by drilling; drilling increased scores less than 8 by 2.51 percentage points or an overall increase of 26 percent. Small for gestational age (SGA), a strong indicator of intrauterine growth restriction (IUGR), increased by 1.81 percentage points or an increase of 18 percent from the mean. Perhaps surprisingly, given that low birth weight is often correlated with premature birth, gestation shows no difference with the introduction of SGD (similar to the findings for premature birth). I do not find any impact on congenital anomaly, despite [McKenzie et al. \(2014\)](#) finding an increase in Colorado. A drilled shale gas well has a small and statistically significant effect on the summary index, increasing the probability of an adverse reproductive health outcome by 0.026 standard deviations. This result is consistent with the finding that living within 1 mile of an operating toxic plant increased the probability of a poor health outcome by 0.016–0.017 standard deviations ([Currie et al., 2014a](#)).

Table 5:

Difference-in-Difference Estimates of the Effect of Drilling on Alternative Health Measures

	(1)	(2)	(3)	(4)	(5)	(6)	(7)
	Birth Weight	APGAR < 8	Gestation	SGA	Congenital Anomaly	Summary Index	
Within 2.5 km	-47.02***	0.0251**	-0.0143	0.0181**	-0.00193	0.0264**	
* post-drilling							
	(12.16)	(0.0101)	(0.0664)	(0.00764)	(0.00189)	(0.0101)	
Observations	21,583	21646	21,631	21524	21,646	21646	
R-squared	0.061	0.029	0.020	0.040	0.008	0.045	
Pre-drilling Mean	3340	0.104	38.74	0.0993	0.00562	-0.0372	

[Open in a new tab](#)

Notes: Each coefficient is from a different regression. See [Table 4](#) for details about included covariates.

Significance:

* p<0.10,

** p<0.05,

*** p<0.01.

V.III. Well Density

Given the finding that the introduction of shale gas development adversely affects birth outcomes in a binary or extensive margin framework, it follows to consider how the density of well development might impact the main outcomes of interest. For the primary sample used in [Table 4](#), the average number of wells drilled at 2.5 km prior to birth is 0.6 wells (s.d. 2.12) with a range of 0 to 35. When

limited to those who have at least one well drilled within 2.5 km prior to birth (the “treatment group”) the average increases to 2.98 wells (s.d. 2.62). In [Table 6](#), I present findings that regress infant health on well density. I find that for each additional shale gas well drilled prior to birth within 2.5 km, low birth weight increases by 0.3 percentage points and term birth weight is reduced by 5 grams. Unlike the previous specification, I also find that each additional well increases premature birth by a similar 0.3 percentage points.^{[28](#)}

Table 6:

Impact of Well Density on Birth Outcomes

	(1)	(2)	(3)	(4)	(5)	(6)
	Low Birth Weight		Term Birth Weight		Premature	
Wells within 2.5 km [*] post	0.00308 ^{***} (0.000868)	0.00306 ^{***} (0.000931)	TM 4.864 ^{***} (1.783)	TM 5.386 ^{***} (1.632)	0.00266 ^{**} (0.00121)	0.00257 ^{**} (0.00123)
Observations	21610	21610	19978	19978	21,189	21,189
R ²	0.009	0.021	0.013	0.076	0.008	0.011
Pre-drilling Mean	0.057	0.057	3416	3416	0.079	0.079
Maternal Characteristics	No	Yes	No	Yes	No	Yes

[Open in a new tab](#)

Notes: Each coefficient is from a different regression. The sample is limited to singleton births and to having a well or permit within 2.5 km. All regressions include an indicator for drilling before birth (defined by closest well), number of wells within 2.5km (including future wells) and the interaction of interest: number of wells within 2.5km *post-drilling. See [Table 4](#) for details about other included covariates.

Significance:

* p<0.10,

** p<0.05,

*** p<0.01.

As before, these findings are consistent across proximity buffers from 2 to 5 km, as shown in [Appendix Table A4](#), and also show some degree of dose response for low birth weight and premature birth. At 2 km, estimates for LBW and preterm birth are about 0.4 percentage points and drop to about 0.02

percentage points at 5 km. The relationship for term birth weight shows less of a dose response, but peaks at 2.5 km with 5 grams and drops to < 1 gram at 5 km.

VI. Robustness Checks and Heterogeneity of Impacts

VI.I. Heterogeneity by Maternal Characteristics

The economics literature measuring health effects of pollution considers avoidance behavior to be an important factor to explore ([Currie \(2009\)](#); [Neidell \(2004\)](#); [Currie et al. \(2014b\)](#)). If families engage in avoidance behavior (e.g. move, use water purification or purchase bottled water ([Wrenn et al., 2016](#)), avoid going outside during drilling), then the health effects measured could be a lower bound. To assess this, the literature tests heterogeneity across characteristics to determine whether there are differential impacts by SES ([Currie et al., 2013b](#); [Sanders and Stoecker, 2015](#)). This would not reflect a biological difference, but would provide evidence for or against maternal behavioral responses to shale gas. [Table 7](#) contains estimates of heterogeneity for three primary measures of infant health: low birth weight, term birth weight, and premature birth (each reported as a separate panel). Each column and coefficient represents an estimate of β_3 in [equation \(1\)](#) from a separate regression to explore whether the effects of exposure to shale gas drilling are the same for different subgroups of the population. For the most part, the results for low birth weight and term birth weight indicate that there is not much heterogeneity of impacts across demographic groups—shale gas development has detrimental impacts on all subgroups. However, high school dropouts and moms on Medicaid do experience larger impacts with increases in low birth weight of about 4 percentage points and college educated mothers have slightly smaller impacts of about 1 percentage point.²⁹ No subgroups have statistically significant impacts for prematurity and similar to before, the signs of the coefficients are not consistently positive or negative.

Table 7:

Shale Gas Development on Maternal Subgroups

	(1)	(2)	(3)	(4)	(5)	(6)
	High School dropout	Smoker	Nonsmoker	Medicaid	WIC	College
Panel A: Low Birth Weight						
Within 2.5 km [*] post	0.0432	0.0186	0.0122 ^{**}	0.0413 ^{***}	0.0138 ^{**}	0.0105
	(0.0268)	(0.0132)	(0.00470)	(0.0120)	(0.00645)	(0.00995)
Observations	2,434	6,465	15,145	7,047	8,541	6,260
R-squared	0.072	0.034	0.018	0.029	0.024	0.029
Pre-drilling Mean	0.0847	0.0830	0.0456	0.0747	0.064	0.0414
Panel B: Term Birth Weight						
Within 2.5 km [*] post	-42.09	-56.15	-51.36 ^{**}	-62.97 [*]	-38.30	-49.61 [*]
	(41.26)	(37.10)	(19.04)	(36.70)	(29.02)	(28.45)
Observations	2,191	5,773	13,763	6,375	7,748	5,699
R-squared	0.131	0.064	0.042	0.077	0.076	0.055
Pre-drilling Mean	3305	3272	3479	3325	3349	3494
Panel C: Premature						
Within 2.5 km [*] post	0.0181	-0.00393	-0.000441	-0.00579	-0.00160	0.000744
	(0.0233)	(0.00950)	(0.00753)	(0.0136)	(0.0142)	(0.0134)
Observations	2,409	6,338	14,851	6,973	8,418	6,122
R-squared	0.070	0.026	0.015	0.027	0.021	0.030
Pre-drilling Mean	0.0896	0.0867	0.0749	0.0859	0.0782	0.0713

Notes: Each coefficient is from a different regression. See [Table 4](#) for details about included covariates.

Significance:

* $p < 0.10$,

** $p < 0.05$,

*** $p < 0.01$.

In [Hill \(2012\)](#), I also report estimates of maternal mobility for the sample of mothers who have multiple singleton births and those who have ever resided within 2.5 km of a well or future well during 2003–2010. I found that moms may be moving in response to shale gas development (an increase of 2.2. percentage points), but it was not statistically significant. Despite some potential increased mobility of these mothers, I found that the results are qualitatively similar for those who stay as those who move and indicate that the main results are not driven by maternal mobility.

VI.II. Sensitivity Analyses

Additional robustness checks were performed to make sure the main specifications are robust to different counterfactual groups, additional controls and subsets of counties associated with production and drilling. These results are reported in [Appendix Table A6](#). First, I limit the sample to mothers who were born in Pennsylvania to test whether migration from out of state is driving the main findings. The results are very similar for the 83 percent of moms who were born in PA.^{[30](#)}

Next, I report the estimates using the 10 most drilled counties and the 10 most producing counties (these are not the same) and find similar results indicating that it is not just drilling or production driving these findings.^{[31](#)}

Another difference-in-difference model commonly used in the environmental health literature is to compare observed health close to a pollution source versus slightly further away. For example, ([Currie and Walker, 2011](#)) compared mothers within 2 km of a toll plaza to mothers who are 2–10 km from a toll plaza, before and after the adoption of E-Z Pass in Pennsylvania and New Jersey. In [Hill \(2012\)](#), I compared residences close to a well (a range of proximities as before of 2–3.5km) and residences a little further away (5, 10 and 15km), before and after drilling.^{[32](#)} The results are consistent with the main findings for low birth weight and term birth weight, but as described in the graphical evidence

section, there may be selection into proximity and so this is not a preferred specification.

VI.III. Falsification Tests

My analysis shows little evidence of any preexisting differences in communities located close to drilled wells relative to communities close to permits or future wells. It is theoretically possible that the increase in low birth weight after drilling is driven by differential trends in fertility or migration post-drilling among mothers who do not have multiple births during the sample. I investigate this possibility by estimating [equation \(1\)](#) using permit dates to define exposure, instead of spud dates. I also create a placebo test using a random date for the closest well. In these specifications, I find no evidence of a spurious effect ([Table 8](#)). I also run models on future wells and repeat the well density models using number of future wells. These models are also consistent with no impact and are consistent with the conclusion that shale gas development has an adverse impact on birth outcomes.

Table 8:

Falsification Tests on Impact of Well Location

	(1)	(2)	(3)	(4)	(5)	(6)
	Permit Date			Random date		
	Low Birth Weight	Term Birth Weight	Premature	Low Birth Weight	Term Birth Weight	Premature
Within 2.5 km _ post	-0.000106	-5.03	-0.00149	0.00103	-1.152	-0.00654
	(0.00682)	(12.382)	(0.00897)	(0.00303)	(11.5)	(.00789)
Sample Size	19246	17795	18854	21610	19978	21204
R ²	0.009	0.013	0.009	0.021	0.075	0.012

[Open in a new tab](#)

Notes: See [Table 4](#) for included covariates. Each panel is a separate regression. All regressions include controls for maternal characteristics and time trends and county fixed effects. Columns (1)- (3) use permit date to define “treatment” and the coefficient reported is the interaction between an indicator for whether the permit was within 2.5 km from the mother’s residence and whether the birth occurred after (post) the permit date. Columns (4)-(6) use a random date to define post birth.

Significance:

* p<0.10,

**p<0.05,

***p<0.01.

VII. Discussion

My results suggest that shale gas development can have adverse effects on the health of people living nearby, namely that of prenatal infants. For the extensive margin, babies born of mothers who lived within 2.5 km of at least one gas well during pregnancy experienced adverse birth outcomes. I find supportive evidence that these effects persist out to 3.5 km of a mother's address and are consistent across multiple specifications. For the intensive margin, or estimating the impact of well density, I find that each additional well drilled within 2.5 km of the mother's residence increases low birth weight and premature birth by 0.4 percentage points and reduces term birth weight by 5 grams.

These results are reasonable for three reasons. First, most areas with shale gas development in Pennsylvania are rural areas with relatively low prevalence of low birth weight (5.7 percent) compared to the state average of 7 percent (for singleton births only).³³ The studies cited in this paper that assess low birth weight impacts of air emissions from other sources (e.g. EZ-Pass, mountain-top coal mining) report baseline average prevalence of low birth weight of 9 or more percent ([Currie and Walker, 2011](#); [Ahern et al., 2011b](#)) and therefore mechanically lower relative effect sizes. However, the average birth weight in this population is almost identical to the state average and is 1.5 percent relative to the mean, which is not large, and is very similar or smaller than the average impact on birth weight of exposure to air emissions in other studies ([Severnini, 2017](#); [Lavaine and Neidell, 2013](#); [Yang and Chou, 2017](#)). Second, most of the existing literature has studied the effects of air pollution on infant health on a pollutant-by-pollutant basis. In this case, I am identifying the health effects of exposure to the disamenity itself, which according to the air emissions inventory emits a wide variety of pollutants. Some, such as NO_x, are much higher than the largest pre-drilling emitter in the region.³⁴ Each of these contaminants have been separately associated with the birth outcomes measured in this paper, while SGD increases exposure to all of these during active drilling and production. Thus, it is not surprising that my estimates are larger than some of those found in the literature, especially those that are studying one pollutant. Finally, these results are smaller than or similar in magnitude to the existing literature studying the infant health impacts of shale gas development ([Stacy et al., 2015b](#); [Casey et al., 2016](#); [Currie et al., 2017](#); [Whitworth et al., 2017](#), 2199).

My study builds upon the existing literature measuring the infant health impacts of shale gas development. Due to inconsistency in measures used across existing studies, it is challenging to compare and interpret measured impacts. My results are consistent with [Currie et al. \(2017\)](#) for low birth weight and [Stacy et al. \(2015a\)](#) for small for gestational age. While I do not find an impact on premature birth in the extensive margin, my intensive margin results indicate that premature birth may be impacted, especially at the highest tertile of exposure. This most closely relates to the inverse distance weighted quartile measures used in the epidemiologic literature and is consistent with [Casey et al. \(2016\)](#) and [Whitworth et al. \(2017\)](#). Although exact mechanisms are difficult to ascertain with the data currently available, the increase in small for gestational age and low birth weight in the extensive

margin without a symmetric increase in premature birth indicates that infants born to mothers exposed to any drilling are coming to full term, but are small, as would be the case where drilling persistently increases local air or water pollution. Whereas, preterm labor may be induced by air pollution or stress at higher intensities of drilling and therefore explain the symmetric intensive margin impacts on preterm birth and low birth weight ([Dole et al., 2003](#); [Stieb et al., 2012b](#); [Sun et al., 2015](#)).

These results suggest that requiring air and water pollution monitoring of drilling sites could assist researchers and public health officials in efforts to ascertain exposure pathways for residents living nearby and inform policies to mitigate any risks that are likely to be very localized. In 2011, PA DEP began requiring the shale gas industry to report emissions of these pollutants into an emissions inventory so that policy makers can better address these exposures in the future.

The effects of gas drilling are larger for lower SES children. There is prior evidence that in some cases this is explained by the fact that lower SES women take fewer measures to avoid pollution. I do not, however, detect heterogeneous responses as measured by mothers moving. As previously mentioned, early shocks to a child's health can persist for many years, hence if poorer families are unable to mitigate the risks of drilling activity their children's health is likely to suffer, which is reflected in literature that finds pollution to be one potential mechanism by which SES affects health ([Neidell, 2004](#)). Given the wealth of studies that identify a causal link between birth weights and long-run outcomes, these impacts are likely to persist throughout these children's lives.

VII.I. Cost Estimates

While the economic benefits and costs of shale gas development are quantifiable, the public health benefits and costs might be more difficult to assess. This paper provides evidence that maternal exposure within at least 1.5 miles of SGD is detrimental to fetal development. Due to shale gas development occurring only recently in Pennsylvania, the number of infants observed close to existing wells is quite small relative to other more populated areas with SGD. This translates to a cost of \$4.1 million.³⁵ As a back-of-the envelope estimate, there are more than 2.8 million American women of reproductive age with a well within a mile of their homes ([Gold and McGinty, 2013](#); [Howden and Meyer, 2010](#)).³⁶ Using the current fertility rate of 64 per 1,000 women in this age group nationally ([Martin et al., 2012](#)), there are over 170,000 pregnant women living within 1 mile of a well in these states. Using the estimates in this paper as a benchmark, oil and gas development in these communities could amount to over 2,000 additional low birth weight infants each year which amounts to a cost of more than \$230 million per year in these 11 states.

VIII. Conclusions

My study seeks to understand and quantify the impacts of shale gas development on infant health. As a first step, I assembled a unique data set with the latitude and longitude of new mothers' residences and the locations of shale gas wells and permits in Pennsylvania. I examine the impacts of living in close proximity to shale gas development on low birth weight, term birth weight and other measures of infant health.

These results suggest that shale gas wells are associated with reduced average birth weight among infants born to mothers living within a 2.5 km radius from a shale gas well; this implies a monetized cost of \$4.1 million. The impacts associated with shale gas studied in this paper are large but not implausible given the estimates found in the literature for air pollution impacts on low birth weight and term birth weight. The strength of this approach is in exploiting a natural experiment that controls for unobservable characteristics and the results are robust across a variety of specifications, providing evidence on the credibility of the research design.

It is clear from these results that policies intended to mitigate the risks of shale gas development can have significant health benefits. I find detectable effects of shale gas development on low birth weight and term birth weight more than 3.5 km from the well head (more than 2 miles or over 11,000 ft). This finding is of significant independent interest and an important contribution of this paper.

Current required set back distances (distance between well head and nearby residences, hospitals and schools) range from 300 ft to 800 ft across the 33 states where shale gas development is taking place. With detectable infant health effects up to 2 miles away, these set back distances may be deemed insufficient to protect human health. The impacts of shale gas development estimated in this paper are independent of drinking water source and suggest that the mechanism by which shale gas development adversely affects reproductive health is through the pathway of air pollution. This finding also adds impetus for regulators to increase regulations that reduce air pollution emissions from drilling operations and for industry actors to increase voluntary action to reduce air pollution emissions.


Since I have focused on only the infant health effects of shale gas development, the total health effects of drilling exposure are likely to be much greater. Further research on the longer term health impacts of shale gas development on all members of our society –as well as the probable mechanisms and how best to mitigate them– is warranted.

Acknowledgments

I am grateful to the Cornell Population Center for their generous financial support. These data were supplied by the Bureau of Health Statistics & Research, Pennsylvania Department of Health, Harrisburg, Pennsylvania. The Pennsylvania Department of Health specifically disclaims responsibility for any analyses, interpretations or conclusions. Thank you to Amy Farrell and James Rubertone of Pennsylvania Department of Health for facilitating access to the data. Gratitude to anonymous reviewers, David Sahn, Jordan Matsudaira, John Cawley, Alison Buttenheim, Joanna Upton, Julia Berazneva, Kira Villa, David Goldsmith, Nick Sanders, Reed Walker, Matt Neidell, Seth Berrin Shonkoff, Doug Almond, Chris Timmins, Lucija Muehlenbachs, Alan Krupnick, Nicolas Ziebarth, Lala Ma, Andrew Boslett, Alina Denham, Mary Willis and seminar participants at TREE Seminar, Cornell University, Columbia University, University of Pittsburgh, University of Maryland College Park, Resources For the Future, PAA 2012, and ISEE 2014 for invaluable comments. A previous version of this paper was circulated as “Unconventional Natural Gas Development and Infant Health: Evidence from Pennsylvania.”

Appendices

Figure A1:



[Open in a new tab](#)

Map of Leasing through 2010

Table A1:

Emissions from Shale Gas Wells First 5 Years after Spud Date

	(1)	(2)	(3)	(4)	(5)	(6)
	co	nox	pm10	pm25	sox	voc
Year of Spud	2.188*** (0.0517)	7.938*** (0.136)	0.282*** (0.00614)	0.259*** (0.00537)	0.107*** (0.00538)	0.585*** (0.0463)
One Year Since Spud	2.241*** (0.0532)	6.709*** (0.140)	0.225*** (0.00632)	0.202*** (0.00552)	0.0656*** (0.00558)	1.008*** (0.0473)
Two Years Since Spud	0.595*** (0.0577)	1.351*** (0.152)	0.0612*** (0.00685)	0.0550*** (0.00596)	0.00860 (0.00607)	0.719*** (0.0501)
Three Years Since Spud	0.378*** (0.0603)	0.661*** (0.158)	0.0289*** (0.00715)	0.0256*** (0.00622)	0.00985 (0.00628)	0.427*** (0.0523)
Four Years Since Spud	0.321*** (0.0737)	0.438** (0.193)	0.0213** (0.00874)	0.0172** (0.00760)	0.00334 (0.00765)	0.502*** (0.0648)
Five Years Since Spud	0.178* (0.100)	0.250 (0.264)	0.0107 (0.0119)	0.00882 (0.0104)	0.00101 (0.0104)	0.731*** (0.0892)
Observations	13,650	13,650	13,610	13,555	13,472	14,073
R-squared	0.215	0.299	0.204	0.218	0.038	0.067
Dep. Var Mean	1.242	3.805	0.136	0.123	0.0436	0.675

[Open in a new tab](#)

Table A2:

Differences in characteristics for analysis sample using DD estimator by Distance

	(1)	(2)	(3)	(4)	(5)	(6)	(7)	
	Teen Mom	Dropout	Black	Smoked	WIC	Medicaid	Born PA	Mile
Within 2km * post-drilling	0.00464 (0.00704)	-0.00150 (0.00927)	0.00181 (0.00457)	-0.00366 (0.0254)	-0.0195 (0.0276)	-0.0288 (0.0273)	-0.0198 (0.0133)	-0.000000 (0.000000)
Observations	14,131	14,131	14,131	14,131	14,026	14,131	14,131	14,131
R-squared	0.015	0.046	0.022	0.031	0.072	0.098	0.025	0.000
Within 2.5 km * post- drilling	0.000550 (0.00666)	-0.0132 (0.0118)	0.00343 (0.00308)	0.00277 (0.0196)	-0.00501 (0.0246)	-0.0204 (0.0282)	-0.0222 (0.0163)	0.000000 (0.000000)
Observations	21646	21646	21646	21646	21469	21646	21646	21646
R-squared	0.012	0.039	0.016	0.026	0.061	0.078	0.020	0.000
Within 3km * post-drilling	-0.00351 (0.0108)	-0.0206 (0.0193)	0.00443 (0.00550)	-0.0210 (0.0234)	-0.0221 (0.0304)	-0.0426 (0.0371)	-0.0209 (0.0139)	0.000000 (0.000000)
Observations	28,910	28,910	28,910	28,910	28,655	28,910	28,910	28,910
R-squared	0.010	0.032	0.016	0.025	0.061	0.073	0.017	0.000
Within 3.5km * post- drilling	-0.0140 (0.0108)	-0.0258 (0.0217)	-0.000432 (0.00694)	-0.0234 (0.0266)	-0.0451 (0.0349)	-0.0451 (0.0419)	-0.0160 (0.0173)	0.000000 (0.000000)

	(1)	(2)	(3)	(4)	(5)	(6)	(7)	
	Teen Mom	Dropout	Black	Smoked	WIC	Medicaid	Born PA	M
Observations	36,447	36,447	36,447	36,447	36,100	36,447	36,447	3
R-squared	0.009	0.029	0.015	0.024	0.057	0.069	0.015	(

[Open in a new tab](#)

Notes: See [Table 3](#) for specification details.

Significance:

* $p < 0.10$,

** $p < 0.05$,

*** $p < 0.01$.

The Effect of Shale Gas Development on Infant Health by Distance

[illegible]

	(1)	(2)	(3)	(4)	(5)	(6)	(7)
	<i>d</i> < 2 km	<i>d</i> < 2:5 km	<i>d</i> < 3 km	<i>d</i> < 3:5 km	<i>d</i> < 4 km	<i>d</i> < 4:5 km	<i>d</i> < 5 km
Well in 'd' km * post-drilling	-0.00962** (0.00403)	0.000354 (0.00664)	0.00460 (0.00455)	-0.00184 (0.00483)	-0.000704 (0.00564)	0.000242 (0.00503)	0.00273 (0.00446)
Observations	13,843	21,189	28,309	35,661	43,741	51,139	57,981
R-squared	0.017	0.012	0.010	0.010	0.009	0.009	0.008
Pre-drilling Mean	0.0802	0.0785	0.0791	0.0791	0.0782	0.0783	0.0786

[Open in a new tab](#)

Notes: See [Table 4](#) for specification details.

Significance:

* $p < 0.10$,

** $p < 0.05$,

*** $p < 0.01$.

Table A4:

Impact of Number of Wells by Proximity

	(1)	(2)	(3)	(4)	(5)	(6)	(7)
	$d < 2 \text{ km}$	$d < 2:5 \text{ km}$	$d < 3 \text{ km}$	$d < 3:5 \text{ km}$	$d < 4 \text{ km}$	$d < 4:5 \text{ km}$	$d < 5 \text{ km}$
Panel A: Low Birth Weight							
Wells in 'd' km * post- drilling	0.00410 [*] (0.00231)	0.00306 ^{***} (0.000931)	0.00232 ^{***} (0.000758)	0.00122 ^{**} (0.000509)	0.000266 (0.000433)	0.000194 (0.000302)	0.000194 (0.000302)
Observations	14,049	21,524	28,756	36,241	44,442	51,994	58,994
R-squared	0.023	0.021	0.020	0.019	0.018	0.018	0.018
Pre-drilling Mean	0.0583	0.0570	0.0578	0.0578	0.0575	0.0573	0.0573
Panel B: Term Birth Weight							
Wells in 'd' km * post- drilling	-3.857 (2.609)	-5.386 ^{***} (1.632)	-4.716 ^{***} (1.331)	-3.152 ^{***} (0.818)	-2.429 ^{***} (0.644)	-1.438 ^{**} (0.570)	-0.918 [*] (0.457)
Observations	12,694	19,463	25,969	32,692	40,067	46,822	53,994
R-squared	0.080	0.076	0.078	0.077	0.079	0.076	0.076
Pre-drilling Mean	3415	3416	3415	3412	3412	3415	3415
Panel C: Premature							
Wells in 'd' km * post- drilling	0.00366 [*]	0.00257 ^{**}	0.00212 ^{**}	0.000889	0.000281	0.000235	0.000235

	(1)	(2)	(3)	(4)	(5)	(6)	(7)
	$d < 2 \text{ km}$	$d < 2.5 \text{ km}$	$d < 3 \text{ km}$	$d < 3.5 \text{ km}$	$d < 4 \text{ km}$	$d < 4.5 \text{ km}$	$d < 5 \text{ km}$
drilling							
	(0.00210)	(0.00123)	(0.000889)	(0.000718)	(0.000602)	(0.000398)	(0.000210)
Observations	13,784	21,109	28,206	35,519	43,506	50,825	57,120
R-squared	0.017	0.011	0.010	0.010	0.009	0.008	0.007
Pre-drilling	0.0803	0.0785	0.0790	0.0789	0.0781	0.0781	0.0781
Mean							

[Open in a new tab](#)

Notes: See [Table 6](#) for specification details.

Significance:

* $p < 0.10$,

** $p < 0.05$,

*** $p < 0.01$.

Table A5:

Robustness Check: Future Number of Wells by Proximity

	(1)	(2)	(3)	(4)
	$d < 2 \text{ km}$	$d < 2.5 \text{ km}$	$d < 3 \text{ km}$	$d < 3.5 \text{ km}$
Panel A: Low Birth Weight				
Wells in 'd' km [*] future	-0.000223 (0.000449)	-0.000133 (0.000341)	8.19e-05 (0.000172)	6.12e-06 (0.000139)
Observations	14,049	21,524	28,756	36,241
R-squared	0.023	0.021	0.020	0.019
Panel B: Term Birth Weight				
Wells in 'd' km [*] future	0.977 (1.342)	0.318 (0.588)	0.410 (0.359)	0.730 ^{**} (0.272)
Observations	12,694	19,463	25,969	32,692
R-squared	0.080	0.076	0.078	0.077
Panel C: Premature				
Wells in 'd' km [*] future	0.000394 (0.000412)	0.000172 (0.000476)	0.000352 (0.000273)	0.000290 (0.000227)
Observations	13,784	21,109	28,206	35,519
R-squared	0.017	0.011	0.010	0.010

[Open in a new tab](#)

Notes: See [Table 6](#) for specification details. Instead of existing wells, this table looks at future wells.

Significance:

* $p < 0.10$,

** $p < 0.05$,

*** $p < 0.01$.

Table A6:

Robustness Checks

	(1)	(2)	(3)
	Low Birth Weight	Term Birth Weight	Premature Birth
Panel A: Mom Born in Pennsylvania			
Within 2.5 km _{post}	0.0128*** (0.00466)	-50.87*** (15.99)	-0.00523 (0.00645)
Observations	17,491	15,814	17,155
R-squared	0.022	0.081	0.012
Pre-drilling Mean	0.0576	3415	0.0791
Panel B: Top 10 Major Production Counties			
Within 2.5 km _{post}	0.0160* (0.00726)	-44.52*** (12.03)	-0.00303 (0.0104)
Observations	15,052	13,627	14,789
R-squared	0.025	0.081	0.017
Pre-drilling Mean	0.0573	3415	0.0790
Panel C: Top 10 Major Drilling Counties			
Within 2.5 km _{post}	0.0175** (0.00576)	-46.66*** (12.36)	0.000296 (0.00978)
Observations	13,208	11,951	12,957
R-squared	0.024	0.076	0.016
Pre-drilling Mean	0.0559	3423	0.0783

Notes: Each coefficient is from a different regression. The sample is limited to singleton births, the sample with a well/permit within 2.5 km and to the panel headings listed. All regressions include indicators for month and year of birth, month*year, residence county indicators, an indicator for drilling before birth (defined by closest well), an indicator for residence within 2.5 km of a well or future well and the interaction of interest of Within 2.5km*post-drilling. Maternal characteristics include mother black, mother Hispanic, mother education (hs, some college, college), mother age (19–24, 25–34, 35+), female child, WIC, smoking during pregnancy, marital status and payment type (private insurance, medicaid, self-pay, other). Indicators for missing data for these variables are also included. Standard errors are in parentheses and clustered at the mother's residence county.

Significance:

* $p < 0.10$,

** $p < 0.05$,

*** $p < 0.01$.

Footnotes

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

¹The International Energy Agency (IEA) defines unconventional gas as sources of gas trapped in impermeable rock deep underground.

²Hydraulic fracturing (popularly known as “fracking” or “fracing”) stimulates the well using a combination of large quantities of water (“high-volume”), fracturing chemicals (“slick water”) and sand that are injected underground at high pressure. This process fractures the rock and causes the resource to be released.

³Pennsylvania experienced very rapid development of shale gas, with 4,272 shale gas wells drilled from 2007–2010 ([PADEP, 2010a](#)).

⁴See [Chay and Greenstone \(2003a\)](#); [Currie and Neidell \(2005\)](#); [Jayachandran \(2009\)](#); [Tanaka \(2015\)](#); [Knittel et al. \(2015\)](#); [Sanders and Stoecker \(2015\)](#); [Clay et al. \(2016\)](#); [Arceo Eva et al. \(2016\)](#); [Yang et al. \(2017\)](#); [Yang and Chou \(2017\)](#); [Severnini \(2017\)](#); [Jha and Muller \(2017\)](#). For example, [Yang et al. \(2017\)](#) found that after a power plant in PA closed down, low birth weight declined by 15 percent and premature birth by 28 percent due to reductions in PM2.5 and SO₂.

⁵See [Colborn et al. \(2011\)](#) regarding health effects of fracturing chemicals; see [McKenzie et al. \(2012\)](#) for a review of studies investigating the effects of inhalation exposure; see [Vengosh et al. \(2014\)](#) for a review of the likely effects of water contamination from SGD; see [Werner et al. \(2015\)](#), [Stacy \(2017\)](#), and [Balise et al. \(2016\)](#) for recent reviews of SGD and health related studies.

⁶ To date, there are no estimates in Pennsylvania of how many properties are “split estate”- the condition where surface owners do not own the mineral rights.

⁷Available upon request.

⁸Scientists face challenges in assessing the potential for contamination due to limited baseline data on water quality, lack of publicly available data regarding the chemicals used in fracturing uid, the sheer number of chemicals use and naturally occurring contaminants returning to the surface in the process of drilling and hydraulic fracturing.

⁹See [Chen et al. \(2017\)](#) for more information about specific chemicals of concern. The EPA Report has a large appendix characterizing each chemical with citations.

¹⁰Toxicity information was lacking for 781 (76%) chemicals. Of the remaining 240 substances, toxicological studies suggested reproductive toxicity for 103 (43%), developmental toxicity for 95 (40%), and both for 41 (17%). Of these 157 chemicals, 67 had or were proposed for a federal water quality standard or guideline.

¹¹This is also true for groundwater public drinking water systems that treat their water prior to distribution.

¹²According to this emissions inventory, shale gas wells emit carbon monoxide, NO_x, PM₁₀, PM_{2.5}, SO_x, volatile organic compounds (VOCs), Benzene, ethylbenzene, formaldehyde, hexane, toluene, xylene, trimethylbenzene, CO₂, and Methane (Author's calculations of wells drilled 2011–2016).

¹³See [Currie et al. \(2009\)](#); [Shah and Balkhair \(2011\)](#); [Stieb et al. \(2012a\)](#); [Glinianaia et al. \(2004\)](#); [Sram et al. \(2005\)](#) for other reviews of past literature related to air pollution and birth outcomes.

¹⁴Both of these study populations are contained within the population studied in this paper.

¹⁵According to the authors, the index does not distinguish between pregnant women living near several producing wells versus well pads under development.

¹⁶ I also test whether drilling activity has affected these characteristics directly by changing fertility and/or the composition of families living near shale gas development and I find few economically significant changes.

¹⁷ [Johnson and Schoeni \(2011\)](#) use national data from the US and find that low birth weight increases the probability of dropping out of high school by one-third, lowers labor force participation by 5 percentage points, and reduces earnings by almost 15 percent. More recently, [Figlio et al. \(2014\)](#) use linked birth and schooling records in Florida and find that birth weight has a significant impact on schooling outcomes for twin births.

¹⁸Small for gestational age (SGA) is used to determine the immediate health care needs of the infant and is used increasingly to predict long-term adverse health outcomes and potential exposure to environmental pollution ([Callaghan and Dietz, 2010](#)). This paper uses the World Health Organization weight percentiles calculator ([WHO, 2011](#)). Another potential measure of reproductive health is the 5 minute American Pediatric Gross Assessment Record (APGAR) score. The physician rates the infant a 0, 1, or 2 on each of 5 dimensions (heart rate, breathing effort, muscle tone, reex initiability, and color), and then sum the scores, giving an APGAR score of 0–10, where 10 is best. This discrete measure is highly correlated (when the score is low) with the need for respiration support at birth ([Almond et al., 2005](#)). Most of these outcomes has been previously examined in both the epidemiological and economics literature (e.g., [Currie and Walker \(2011\)](#)). Following [Currie et al. \(2014a\)](#), I also construct a single standardized measure to address examining multiple outcomes and multiple hypothesis tests. I first convert each birth measure so that an increase is “adverse” and then standardize the measure to a mean of zero and standard deviation of 1. I then construct the summary measure by taking the mean over the standardized outcomes, weighting them equally.

¹⁹ An examination of fertility over time suggests a consistent number of births within 2.5 km of the well head. [Muehlenbachs et al. \(2015\)](#) do not find any changes in neighborhood composition using Census data at the tract level from 2000–2012 in Pennsylvania.

²⁰This is the largest distance used as a treated group in related studies. [McKenzie et al. \(2014\)](#) use 10 miles, [Stacy et al. \(2015b\)](#) use 10 miles, [Casey et al. \(2016\)](#) uses 20km, [Whitworth et al. \(2017\)](#) use 10 miles and [Currie et al. \(2017\)](#) use 15 km.

²¹In [Appendix Tables A3](#) and [A4](#), I report different proximities to gas wells for the definition of treatment and show that for distances up to 5 km, the results are fairly robust.

²²By including permitted wells not drilled, this estimation strategy becomes more than just a pre-post analysis. This identification strategy assumes that infants born within a similar distance to a permit that is a potential future

²³Due to the localized nature of this estimation strategy, there is little variation within zip codes to allow for zip code fixed effects. Models with zip code fixed effects are qualitatively similar but less precisely estimated. Results available upon request.

²⁴ Only one maternal characteristic shows a significant change with drilling: mothers observed after drilling are more educated than those observed prior to drilling (results not shown). Increased college completions among mothers would potentially improve observed infant health in these communities. However, this does suggest some selection and so I include these and other controls in all the subsequent results. The time frame of interest is during the onset of the Great Recession. It may indicate that the opportunity cost of going to college, or becoming a mother, has reduced and so more educated mothers are having children. Other research has linked recessions to improved infant health outcomes, so it is unlikely to be the driver of impacts reported in the next section ([Chay and Greenstone, 2003b](#); [Dehejia and Lleras-Muney, 2004](#)).

²⁵Overall prevalence is calculated as follows: $0.0136/0.057=23.9$ percent low birth weight and $49.6/3416 = 1.5$ percent reduction in term birth weight.

²⁶Permitted wells must have already gone through the leasing process and households that lease their mineral rights will have received signing bonuses previously. These benefits can only reach an approximate 3km buffer where horizontal drilling can reach minerals and would result in royalties. At very close proximities (e.g. < 1km), I see some indication that birth outcomes are improved by drilling. There is a large and growing literature that suggests positive income shocks can have a positive effect on birth outcomes ([Almond et al., 2011](#); [Hoynes et al., 2015](#)) and so this ending would be consistent with that hypothesis. Royalties may mitigate the risks of close exposure.

²⁷Following [Currie et al. \(2014a\)](#), I address the issue of precision using a summary index measure of infant health. I first convert each birth measure so that an increase is “adverse” and then standardize the measure to a mean of zero and standard deviation of 1. I then construct the summary measure by taking the mean over the standardized outcomes, weighting them equally.

²⁸I also estimate models using tertiles of wells and find that the top tertile (> 3 wells) has a similar sized effect as the extensive margin results for low birth weight and term birth weight, however, the top tertile increases premature birth by 2 percentage points, in contrast to the null finding in the extensive margin results.

²⁹The pre-drilling mean for these three groups are substantially different from the overall average. The percent changes relative to the mean for both HS dropouts and Medicaid recieve a 50 percent increase, while the effect for college educated moms recieve a 25 percent increase, which is the same as the main effect. I tested the differences between these and the main results and only the results for Medicaid are statistically different [pvalue=0.01]

³⁰This does not perfectly address this question since migration can also occur within PA.

³¹Other robustness checks were reported in [Hill \(2012\)](#). First, I showed the results for restricting the sample to infants born within 2 years (before and after) of the spud date for the closest well. This specification is designed to address any possible concerns about unequal prior and post observation periods for each location or concerns about unobserved and differential sorting in the mothers living close to drilled versus permitted wells. The point estimates are somewhat smaller, but qualitatively similar to the estimates in Tables [4](#) and [5](#). Next I showed the results using the sample of births from 2008 to 2010, when most of the shale gas development took place during the sample frame. This point estimate is slightly larger for low birth weight (LBW) indicating a 1.89 percentage point increase. Finally, I reported the results from adding the continuous distance to the closest well, as well as the number of wells drilled within 5 km of the maternal residence. Again, the point estimates are very similar to those reported in Tables [4](#) and [5](#) and suggest most of the effect is driven by proximity to the closest well.

³²In [Hill \(2012\)](#), I used up to 15 km as the comparison group and reported it as a lower-bound estimate; shale gas development increases the overall prevalence of low birth weight by 12.5 percent and reduces term birth weight by 0.6 percent, on average. Depending on the scale of shale gas development, it is possible that other aspects of drilling activity will influence infant health within 15 km of a well and could explain these smaller estimates. For example, communities with shale gas development are exposed to increased truck traffic, pipelines, water storage, compressor stations and

general increased localized economic activity. These community level effects are less likely to influence the estimates in the main results of the paper that use permitted/future wells as the comparison group.

³³Using the pre-drilling mean of low birth weight for the analysis sample, the effect size is 24 percent relative to the mean, whereas the effect size is 19 percent relative to the state average.

³⁴As mentioned in the background section of the paper, the largest industrial source of NO_x in the 11-county region is a power plant that produces 2,000 tons per year. Shale wells in 2011 produced 16,000 tons of NO_x in aggregate.

³⁵Combining hospital costs attributable to low birth weight (\$15,100 in additional hospital costs) ([Russell et al., 2007](#)), estimates for special education services (\$5,200) ([Chaikind and Corman, 1991](#); [Augenblick et al., 2007](#)) and decreased earnings (\$76,800) ([Currie et al., 2013a](#)), an arguably conservative estimate is \$96,500 in added cost for each low birth weight child. This figure excludes medical bills after the first year, parental lost earnings and other costs and is, hence, a lower bound estimate of costs.

³⁶Using The Wall Street Journal estimate that over 15 million Americans live within 1 mile of an oil or gas well drilled since 2000, and using a rough estimate that half of those people are women and forty percent of them are ages 18–44.

References

1. Ahern Melissa, Mullett Martha, MacKay Katherine, and Hamilton Candice. 2011b. “Residence in Coal-Mining Areas and Low-Birth-Weight Outcomes.” *Maternal and Child Health Journal*, 15(7): 974–979, URL: <https://link.springer.com/article/10.1007/s10995-009-0555-1> , DOI: 10.1007/s10995-009-0555-1. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
2. Ahern MM, Hendryx M, Conley J, Fedorko E, Ducatman A, and Zullig KJ. 2011a. “The association between mountaintop mining and birth defects among live births in central Appalachia, 1996–2003.” *Environmental research*, 111(6): 838–846. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
3. Alawattagama Shyama K, Tetiana Kondratyuk, Renee Krynock, Bricker Matthew, Rutter Jennifer K, Bain Daniel J, and Stolz John F. 2015. “Well water contamination in a rural community in southwestern Pennsylvania near unconventional shale gas extraction.” *Journal of*

Environmental Science and Health, Part A, 50(5): 516–528. [DOI] [PubMed] [Google Scholar]

4. Allcott Hunt, and Keniston Daniel. 2014. “Dutch disease or agglomeration? The local economic effects of natural resource booms in modern America.” Technical report, National Bureau of Economic Research. [Google Scholar]

5. Almond D, Chay KY, and Lee DS. 2005. “The Costs of Low Birth Weight.” The Quarterly Journal of Economics, 120(3): 1031–1083. [Google Scholar]

6. Almond Douglas, Hoynes Hilary W, and Schanzenbach Diane Whitmore. 2011. “Inside the war on poverty: The impact of food stamps on birth outcomes.” The Review of Economics and Statistics, 93(2): 387–403. [Google Scholar]

7. Eva Arceo, Rema Hanna, and Paulina Oliva. 2016. “Does the Effect of Pollution on Infant Mortality Differ Between Developing and Developed Countries? Evidence from Mexico City.” The Economic Journal, 126(591): 257–280, URL: [https:// onlinelibrary-wiley-com.ezp.lib.rochester.edu/doi/full/10.1111/econj.12273](https://onlinelibrary-wiley-com.ezp.lib.rochester.edu/doi/full/10.1111/econj.12273) , DOI: 10.1111/econj.12273. [DOI] [Google Scholar]

8. Augenblick Palaich, Water G Van de, and Myer JL. 2007. “Costing Out the Resources Needed to Meet Pennsylvania’s Public Education Goals.” Pennsylvannia State Board of Education, URL: <http://ridley.schoolwires.net/cms/lib2/PA01001042/Centricity/Domain/5/Costing-OutStudyRevisedFinalReport12-10-07.pdf> . [Google Scholar]

9. Balise Victoria D, Chun-Xia Meng, Cornelius-Green Jennifer N, Kassotis Christopher D, Kennedy Rana, and Nagel Susan C. 2016. “Systematic review of the association between oil and natural gas extraction processes and human reproduction.” Fertility and Sterility, 106(4): 795–819. [DOI] [PMC free article] [PubMed] [Google Scholar]

10. Bamberger M, and Oswald RE. 2012. “Impacts of Gas Drilling on Human and Animal Health.” NEW SOLUTIONS: A Journal of Environmental and Occupational Health Policy 51–77. [DOI] [PubMed] [Google Scholar]

11. Bartik Alexander Wickman, Currie Janet, Greenstone Michael, and Christopher R Knittel. 2016. “The Local Economic and Welfare Consequences of Hydraulic Fracturing.” [Google Scholar]

12. Boyles Abee L., Blain Robyn B., Rochester Johanna R., Avanası Raghavendhran, Goldhaber Susan B., Sofie McComb Stephanie D. Holmgren, Masten Scott A., and Thayer Kristina A.. 2017.

“Systematic review of community health impacts of mountaintop removal mining.”
Environment International, 107 163–172, URL: <http://www.sciencedirect.com/science/article/pii/S016041201730140X> , DOI: 10.1016/j.envint.2017.07.002. [DOI] [PMC free article] [PubMed] [Google Scholar]

13. Burton Taylour G, Hanadi S Rifai, Hildenbrand Zacariah L, Carlton Doug D, Fontenot Brian E, and Schug Kevin A. 2016. “Elucidating hydraulic fracturing impacts on groundwater quality using a regional geospatial statistical modeling approach.” Science of The Total Environment, 545 114–126. [DOI] [PubMed] [Google Scholar]

14. Callaghan WM, and Dietz PM. 2010. “Differences in birth weight for gestational age distributions according to the measures used to assign gestational age.” American journal of epidemiology, 171(7): 826–836. [DOI] [PubMed] [Google Scholar]

15. Casey Joan A, Savitz David A, Rasmussen Sara G, Ogburn Elizabeth L, Pollak Jonathan, Mercer Dione G, and Schwartz Brian S. 2015. “Unconventional Natural Gas Development and Birth Outcomes in Pennsylvania, USA..” Epidemiology (Cambridge, Mass.). [DOI] [PMC free article] [PubMed] [Google Scholar]

16. Casey Joan A, Savitz David A, Rasmussen Sara G, Ogburn Elizabeth L, Pollak Jonathan, Mercer Dione G, and Schwartz Brian S.. 2016. “Unconventional natural gas development and birth outcomes in Pennsylvania, USA.” Epidemiology (Cambridge, Mass.), 27(2): 163–172, URL: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4738074/>, DOI: 10.1097/EDE.0000000000000387. [DOI] [PMC free article] [PubMed] [Google Scholar]

17. Chaikind Stephen, and Corman Hope. 1991. “The impact of low birthweight on special education costs.” Journal of Health Economics, 10(3): 291 – 311. [DOI] [PubMed] [Google Scholar]

18. Chay KY, and Greenstone M. 2003a. “Air quality, infant mortality, and the Clean Air Act of 1970.” Technical report, National Bureau of Economic Research. [Google Scholar]

19. Chay KY, and Greenstone M. 2003b. “The Impact of Air Pollution on Infant Mortality: Evidence from Geographic Variation in Pollution Shocks Induced by a Recession*.” The Quarterly journal of economics, 118(3): 1121–1167. [Google Scholar]

20. Chen Huan, and Carter Kimberly E. 2016. “Water usage for natural gas production through hydraulic fracturing in the United States from 2008 to 2014.” Journal of environmental management, 170 152–159. [DOI] [PubMed] [Google Scholar]

21. Chen Lu, Miller Shelie A., and Ellis Brian R.. 2017. "Comparative Human Toxicity Impact of Electricity Produced from Shale Gas and Coal." *Environmental Science & Technology*, 51(21): 13018–13027, URL: 10.1021/acs.est.7b03546, DOI: <http://dx.doi.org/10.1021/acs.est.7b03546> . [DOI] [PubMed] [Google Scholar]
22. Clay Karen, Lewis Joshua, and Severnini Edson. 2016. "Canary in a Coal Mine: Infant Mortality, Property Values, and Tradeoffs Associated with Mid-20th Century Air Pollution." Technical Report w22155, National Bureau of Economic Research. [Google Scholar]
23. Colborn T, Kwiatkowski C, Schultz K, and Bachran M. 2011. "Natural gas operations from a public health perspective." *Human and Ecological Risk Assessment: An International Journal*, 17(5): 1039–1056. [Google Scholar]
24. Colborn Theo, Schultz Kim, Herrick Lucille, and Kwiatkowski Carol. 2012. "An Exploratory Study of Air Quality near Natural Gas Operations." *Human and Ecological Risk Assessment: An International Journal*. [Google Scholar]
25. Currie J 2009. "Healthy, wealthy, and wise: Socioeconomic status, poor health in childhood, and human capital development." *Journal of Economic Literature*, 47(1): 87–122. [Google Scholar]
26. Currie J, and Neidell M. 2005. "Air Pollution and Infant Health: What Can We Learn from California's Recent Experience?" *Quarterly journal of economics*, 120(3): 1003–1030. [Google Scholar]
27. Currie J, Neidell M, and Schmieder JF. 2009. "Air pollution and infant health: Lessons from New Jersey." *Journal of health economics*, 28(3): 688–703. [DOI] [PMC free article] [PubMed] [Google Scholar]
28. Currie J, and Schmieder JF. 2009. "Fetal Exposures to Toxic Releases and Infant Health." *The American Economic Review*, 99(2): 177–183. [DOI] [PMC free article] [PubMed] [Google Scholar]
29. Currie J, and Walker R. 2011. "Traffic Congestion and Infant Health: Evidence from EZPass." *American Economic Journal: Applied Economics*, 3(1): 65–90. [Google Scholar]
30. Currie Janet, Davis Lucas, Greenstone Michael, and Walker Reed. 2013a. "Do Housing Prices Reflect Environmental Health Risks? Evidence from More than 1600 Toxic Plant Openings and Closings." Technical report, National Bureau of Economic Research. [Google Scholar]

31. Currie Janet, Davis Lucas, Greenstone Michael, and Walker Reed. 2014a. "Do housing prices reflect environmental health risks? Evidence from more than 1600 toxic plant openings and closings." [DOI] [PMC free article] [PubMed] [Google Scholar]
32. Currie Janet, Greenstone Michael, and Meckel Katherine. 2017. "Hydraulic fracturing and infant health: New evidence from Pennsylvania." *Science Advances*, 3(12):, p. e1603021, URL: <http://advances.sciencemag.org/content/3/12/e1603021> , DOI: 10.1126/sciadv.1603021. [DOI] [PMC free article] [PubMed] [Google Scholar]
33. Currie Janet, Joshua Graff Zivin Jamie Mullins, and Neidell Matthew. 2014. b. "What Do We Know About Short- and Long-Term Effects of Early-Life Exposure to Pollution?" *Annual Review of Resource Economics*, 6(1): 217–247, URL: 10.1146/annurev-resource-100913-012610, DOI: 10.1146/annurev-resource-100913-012610. [DOI] [Google Scholar]
34. Currie Janet, Graff Zivin Joshua S., Meckel Katherine, Neidell Matthew J., and Schlenker Wolfram. 2013b. "Something in the Water: Contaminated Drinking Water and Infant Health." [DOI] [PMC free article] [PubMed] [Google Scholar]
35. Dehejia Rajeev, and Lleras-Muney Adriana. 2004. "Booms, Busts, and Babies' Health." *The Quarterly Journal of Economics*, 119(3): 1091–1130. [Google Scholar]
36. Dole Nancy, Savitz David A. Hertz-Picciotto abd Irva, Siega-Riz Anna Maria, McMahon Michael J., and Buekens Pierre. 2003. "Maternal stress and preterm birth." *American journal of Epidemiology*, 157(1): 14–24. [DOI] [PubMed] [Google Scholar]
37. Drollette Brian D, Hoelzer Kathrin, Warner Nathaniel R, Darrah Thomas H, Karatum Osman, O'Connor Megan P, Nelson Robert K, Fernandez Loretta A, Reddy Christopher M, Vengosh Avner et al. 2015. "Elevated levels of diesel range organic compounds in groundwater near Marcellus gas operations are derived from surface activities." *Proceedings of the National Academy of Sciences*, p. 201511474. [DOI] [PMC free article] [PubMed] [Google Scholar]
38. Elliott Elise G, Ettinger Adrienne S, Leaderer Brian P, Bracken Michael B, and Deziel Nicole C. 2017. "A systematic evaluation of chemicals in hydraulic-fracturing fluids and wastewater for reproductive and developmental toxicity." *Journal of Exposure Science and Environmental Epidemiology*, 27(1): 90–99. [DOI] [PubMed] [Google Scholar]
39. EPA. 2016. "Hydraulic Fracturing for Oil and Gas: Impacts from the Hydraulic Fracturing Water Cycle on Drinking Water Resources in the United States (Final Report)." URL: <https://www.epa.gov/hfstudy> , Accessed January, 2016.

40. Ferrar Kyle J, Kriesky Jill, Christen Charles L, Marshall Lynne P, Malone Samantha L, Sharma Ravi K, Michanowicz Drew R, and Goldstein Bernard D. 2013. "Assessment and longitudinal analysis of health impacts and stressors perceived to result from unconventional shale gas development in the Marcellus Shale region." *International journal of occupational and environmental health*, 19(2): 104–112. [DOI] [PubMed] [Google Scholar]
41. Feyrer James, Mansur Erin T, and Sacerdote Bruce. 2017. "Geographic Dispersion of Economic Shocks: Evidence from the Fracking Revolution." *American Economic Review*. [Google Scholar]
42. Figlio David, Guryan Jonathan, Karbownik Krzysztof, and Roth Jeffrey. 2014. "The effects of poor neonatal health on children's cognitive development." *The American Economic Review*, 104(12): 3921–3955. [DOI] [PubMed] [Google Scholar]
43. Glinianaia SV, Rankin J, Bell R, Pless-Mulloli T, and Howel D. 2004. "Particulate air pollution and fetal health: a systematic review of the epidemiologic evidence." *Epidemiology*, 15(1):, p. 36. [DOI] [PubMed] [Google Scholar]
44. Gold Russell, and Tom McGinty. 2013. "Energy Boom Puts Wells in America's Backyards." URL: <http://online.wsj.com/news/articles/SB10001424052702303672404579149432365326304> , Accessed October, 2013. [Google Scholar]
45. Goldenberg Robert L, Culhane Jennifer F, Iams Jay D, and Romero Roberto. 2008. "Epidemiology and causes of preterm birth." *The Lancet*, 371(9606): 75–84, URL: <http://www.sciencedirect.com/science/article/pii/S0140673608600744> , DOI: 10.1016/S0140-6736(08)60074-4. [DOI] [PMC free article] [PubMed] [Google Scholar]
46. Goodman Paul S, Galatioto Fabio, Thorpe Neil, Namdeo Anil K, Davies Richard J, and Bird Roger N. 2016. "Investigating the traffic-related environmental impacts of hydraulic-fracturing (fracking) operations." *Environment International* , 89-90 248–260, URL: <http://www.sciencedirect.com/science/article/pii/S0160412016300277> , DOI: 10.1016/j.envint.2016.02.002. [DOI] [Google Scholar]
47. Graham Jove, Irving Jennifer, Tang Xiaoqin, Sellers Stephen, Crisp Joshua, Horwitz Daniel, Muehlenbachs Lucija, Krupnick Alan, and Carey David. 2015. "Increased traffic accident rates associated with shale gas drilling in Pennsylvania." *Accident Analysis & Prevention*, 74 203–209. [DOI] [PubMed] [Google Scholar]
48. Hausman Catherine, and Kellogg Ryan. 2015. "Welfare and Distributional Implications of

Shale Gas.” Brookings Papers on Economic Activity. [[Google Scholar](#)]

49. Hendryx Michael. 2015. “The public health impacts of surface coal mining.” *The Extractive Industries and Society*, 2(4): 820–826. [[Google Scholar](#)]

50. Hildenbrand Zacariah L, Carlton Doug D, Fontenot Brian E, Meik Jesse M, Walton Jayme L, Thacker Jonathan B, Korlie Stephanie, Shelor C Phillip, Kadjo Akinde F, Clark Adelaide et al. 2016. “Temporal variation in groundwater quality in the Permian Basin of Texas, a region of increasing unconventional oil and gas development.” *Science of The Total Environment*, 562 906–913. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

51. Hildenbrand Zacariah L, Carlton Doug D Jr, Fontenot Brian E, Meik Jesse M, Walton Jayme L, Taylor Josh T, Thacker Jonathan B, Korlie Stephanie, Shelor C Phillip, Henderson Drew et al. 2015. “A comprehensive analysis of groundwater quality in the Barnett Shale region.” *Environmental science & technology*, 49(13): 8254–8262. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

52. Hill Elaine. 2012. “Unconventional Natural Gas Development and Infant Health: Evidence from Pennsylvania.” Cornell Dyson School Working Paper. [[Google Scholar](#)]

53. Hill Elaine. 2013. “The Impact of Oil and Gas Development on Infant Health in Colorado.” Working Paper. [[Google Scholar](#)]

54. Hill Elaine, and Ma Lala. 2017. “Shale Gas Development and Water Quality.” *American Economic Review, Papers and Proceedings*. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

55. Hladik Michelle L, Focazio Michael J, and Engle Mark. 2014. “Discharges of produced waters from oil and gas extraction via wastewater treatment plants are sources of disinfection by-products to receiving streams.” *Science of the Total Environment*, 466 1085–1093. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

56. Howden Lindsay M, and Meyer Julie A. 2010. “Age and sex composition: 2010” US CENSUS BUREAU. [[Google Scholar](#)]

57. Hoynes Hilary, Miller Doug, and Simon David. 2015. “Income, the earned income tax credit, and infant health.” *American Economic Journal: Economic Policy*, 7(1): 172–211. [[Google Scholar](#)]

58. IEA. 2012. “Golden Rules for a Golden Age of Natural Gas.” Technical report, International Energy Agency. [[Google Scholar](#)]

59. Jackson Robert B, Vengosh Avner, Darrah Thomas H, Warner Nathaniel R, Down Adrian, Poreda Robert J, Osborn Stephen G, Zhao Kaiguang, and Karr Jonathan D. 2013. "Increased stray gas abundance in a subset of drinking water wells near Marcellus shale gas extraction." *Proceedings of the National Academy of Sciences*. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
60. Jayachandran S 2009. "Air Quality and Early-Life Mortality." *Journal of Human Resources*, 44(4): 916–954. [[Google Scholar](#)]
61. Jemielita Thomas, George L Gerton Matthew Neidell, Chillrud Steven, Yan Beizhan, Stute Martin, Howarth Marilyn, Saberi Poun'e, Fausti Nicholas, Penning Trevor M et al. 2015. "Unconventional gas and oil drilling is associated with increased hospital utilization rates." *PloS one*, 10(7):, p. e0131093. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
62. Jha Akshaya, and Muller Nicholas Z.. 2017. "Handle with Care: The Local Air Pollution Costs of Coal Storage.." Working Paper 23417, National Bureau of Economic Research, DOI: 10.3386/w23417. [[DOI](#)] [[Google Scholar](#)]
63. Johnsen Reid, LaRiviere Jacob S, and Wolff Hendrik. 2016. "Estimating Indirect Benefits: Fracking, Coal and Air Pollution." SSRN Scholarly Paper ID 2834220, Social Science Research Network, Rochester, NY. [[Google Scholar](#)]
64. Johnson Rucker C., and Schoeni Robert F.. 2011. "The influence of early-life events on human capital, health status, and labor market outcomes over the life course." *The BE journal of economic analysis and policy*, 11(3):. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
65. Kargbo David M, Wilhelm Ron G, and Campbell David J. 2010. "Natural gas plays in the Marcellus shale: Challenges and potential opportunities." *Environmental Science & Technology*, 44(15): 5679–5684. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
66. Kassotis Christopher D, Tillitt Donald E, Chung-Ho Lin, McElroy Jane A, and Nagel Susan C. 2015. "Endocrine-Disrupting Chemicals and Oil and Natural Gas Operations: Potential Environmental Contamination and Recommendations to Assess Complex Environmental Mixtures." *Environmental health perspectives*. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
67. Knittel Christopher R., Miller Douglas L., and Sanders Nicholas J.. 2015. "Caution, Drivers! Children Present: Traffic, Pollution, and Infant Health." *The Review of Economics and Statistics*, 98(2): 350–366, URL: 10.1162/REST_a_00548, DOI: 10.1162/REST_a_00548. [[DOI](#)] [[Google Scholar](#)]

68. Kondash, Andrew J, Elizabeth Albright, and Avner Vengosh. 2017. "Quantity of flowback and produced waters from unconventional oil and gas exploration." *Science of the Total Environment*, 574 314–321. [DOI] [PubMed] [Google Scholar]
69. Lamm Steven H., Li Ji, Robbins Shayhan A., Dissen Elisabeth, Chen Rusan, and Feinleib Manning. 2015. "Are residents of mountain-top mining counties more likely to have infants with birth defects? The west virginia experience." *Birth Defects Research Part A: Clinical and Molecular Teratology*, 103(2): 76–84, URL: <http://onlinelibrary.wiley.com/doi/10.1002/bdra.23322/abstract>, DOI: 10.1002/bdra.23322. [DOI] [PubMed] [Google Scholar]
70. Lavaine Emmanuelle, and Neidell Matthew J.. 2013. "Energy Production and Health Externalities: Evidence from Oil Refinery Strikes in France." [Google Scholar]
71. Legere Laura. 2013. "Northern Tier counties top state list of Marcellus air pollution." URL: <http://stateimpact.npr.org/pennsylvania/jp/northern-tier-counties-top-state-list-of-marcellus-air-pollution/>, Accessed March 2013. [Google Scholar]
72. Lester Yaal, Ferrer Imma, Thurman E Michael, Sitterley Kurban A, Korak Julie A, Aiken George, and Linden Karl G. 2015. "Characterization of hydraulic fracturing flowback water in Colorado: Implications for water treatment." *Science of the Total Environment*, 512 637–644. [DOI] [PubMed] [Google Scholar]
73. Li Huishu, Son Ji-Hee, and Carlson Kenneth H. 2016. "Concurrence of aqueous and gas phase contamination of groundwater in the Wattenberg oil and gas field of northern Colorado." *Water research*, 88 458–466. [DOI] [PubMed] [Google Scholar]
74. Litovitz Aviva, Curtright Aimee, Abramzon Shmuel, Burger Nicholas, and Samaras Constantine. 2013. "Estimation of regional air-quality damages from Marcellus Shale natural gas extraction in Pennsylvania." *Environmental Research Letters*, 8(1);, p. 014017. [Google Scholar]
75. Loomis John, and Haefele Michelle. 2017. "Quantifying Market and Non-market Benefits and Costs of Hydraulic Fracturing in the United States: A Summary of the Literature." *Ecological Economics*, 138 160–167, URL: <http://www.sciencedirect.com/science/article/pii/S092180091631285X>, DOI: 10.1016/j.ecolecon.2017.03.036. [DOI] [Google Scholar]
76. Martin J, B Hamilton S Ventura M Osterman, Wilson E, and Mathew TJ. 2012. "Births: Final Data for 2010." *National Vital Statistics Report*. [PubMed] [Google Scholar]
77. Mason Charles F, Lucija Muehlenbachs, and Olmstead Sheila M. 2015. "The economics of

shale gas development.” Resources for the Future Discussion Paper 14–42. [[Google Scholar](#)]

78. McCawley Michael A. 2017. “Does increased traffic flow around unconventional resource development activities represent the major respiratory hazard to neighboring communities?: knowns and unknowns.” Current Opinion in Pulmonary Medicine, 23(2): 161–166, URL: <http://content.wkhealth.com/linkback/openurl?sid=WKPTLP:landingpage&an=00063198-900000000-99324> , DOI: 10.1097/MCP.0000000000000361. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

79. McKenzie Lisa M, Ruixin Guo, Witter Roxana Z, Savitz David A, Newman Lee S, and Adgate John L. 2014. “Birth Outcomes and Maternal Residential Proximity to Natural Gas Development in Rural Colorado.” Environmental health perspectives. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

80. McKenzie LM, Witter RZ, Newman LS, and Adgate JL. 2012. “Human health risk assessment of air emissions from development of unconventional natural gas resources.” Science of The Total Environment. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

81. Muehlenbachs Lucija, Spiller Elisheba, and Timmins Christopher. 2015. “The Housing Market Impacts of Shale Gas Development.” American Economic Review, 105(12): 3633–59. [[Google Scholar](#)]

82. Neidell Matthew J. 2004. “Air pollution, health, and socio-economic status: the effect of outdoor air quality on childhood asthma.” Journal of health economics, 23(6): 1209–1236. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

83. Nieuwenhuijsen Mark J, Payam Dadvand, James Grellier, David Martinez, and Martine Vrijheid. 2013. “Environmental risk factors of pregnancy outcomes: a summary of recent meta-analyses of epidemiological studies.” Environmental Health, 12, p. 6, URL: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3582445/>, DOI: 10.1186/1476-069X-12-6. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

84. Olmstead Sheila, Muehlenbachs Lucija, Shih Jhih-Shyang, Chu Ziyan, and Krupnick Alan. 2013. “Shale Gas Development Impacts On Surface Water Quality in Pennsylvania.” Proceedings of the National Academy of Sciences. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

85. Osborn Stephen, Vengosh Avner, Warner Nathaniel, and Jackson Robert. 2011. “Methane Contamination of Drinking Water Accompanying Gas Well Drilling and Hydraulic Fracturing.” Proceedings of the National Academy of Sciences, 108(20): 8172–8176. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

86. PADEP. 2010a. "Marcellus Permits Issued and Wells Drilled." URL: <http://www.dep.state.pa.us/dep/deputate/minres/oilgas/2010PermitDrilledmaps.htm> .
87. PADEP. 2010b. "Southwestern Pennsylvania Marcellus Shale Short-Term Ambient Air Sampling Report." URL: http://www.dep.state.pa.us/dep/deputate/airwaste/aq/aqm/docs/Marcellus_SW_11-01-10.pdf , Accessed January, 2012.
88. PADEP. 2011a. "Northcentral Pennsylvania Marcellus Shale Short-Term Ambient Air Sampling Report." URL: http://www.dep.state.pa.us/dep/deputate/airwaste/aq/aqm/docs/Marcellus_NC_05-06-11.pdf , Accessed January, 2012.
89. PADEP. 2011b. "Northeastern Pennsylvania Marcellus Shale Short-Term Ambient Air Sampling Report." URL: http://www.dep.state.pa.us/dep/deputate/airwaste/aq/aqm/docs/Marcellus_NE_01-12-11.pdf , Accessed January, 2012.
90. PADEP. 2013a. "Air Emissions Inventory Data for the Unconventional Natural Gas Industry." URL: http://files.dep.state.pa.us/Air/AirQuality/AQPortalFiles/Unconventional_Natural_Gas_Emissions_Well-Station-All.xlsx , Accessed June, 2013.
91. PADEP. 2013b. "Summary of Unconventional Natural Gas Emissions by County." URL: http://files.dep.state.pa.us/Air/AirQuality/AQPortalFiles/Unconventional_Natural_Gas_Emissions-County.xlsx , Accessed June, 2013.
92. Parker Kimberly M, Zeng Teng, Harkness Jennifer, Vengosh Avner, and Mitch William A. 2014. "Enhanced Formation of Disinfection Byproducts in Shale Gas Wastewater Impacted Drinking Water Supplies." *Environmental science & technology*, 48(19): 11161–11169. [DOI] [PubMed] [Google Scholar]
93. Peng Lizhong, Meyerhoefer Chad, and Chou ShinYi. 2018. "The health implications of unconventional natural gas development in Pennsylvania." *Health Economics*, 0(0):, URL: <https://onlinelibrary-wiley-com.ezp.lib.rochester.edu/doi/10.1002/hec.3649> , DOI: 10.1002/hec.3649. [DOI] [PubMed] [Google Scholar]
94. Pétron Gabrielle, Frost Gregory, Miller Benjamin R, Hirsch Adam I, Montzka Stephen A, Karion Anna, Trainer Michael, Sweeney Colm, Andrews Arlyn E, Miller Lloyd et al. 2012. "Hydrocarbon emissions characterization in the Colorado Front Range: A pilot study." *Journal of Geophysical Research: Atmospheres* (1984–2012), 117(D4):. [Google Scholar]
95. Quansah Reginald, Frederick Ato Armah, Essumang David Kofi, Luginaah Isaac, Clarke Edith, Marfoh Kissinger, Samuel Jerry Cobbina Edward Nketiah-Amponsah, Proscovia Bazanya

- Namujju Samuel Obiri, and Dzodzomenyo Mawuli. 2015. "Association of Arsenic with Adverse Pregnancy Outcomes/Infant Mortality: A Systematic Review and Meta-Analysis." *Environmental Health Perspectives*, 123(5): 412–421, URL: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4421764/>, DOI: 10.1289/ehp.1307894. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
96. Rahm Brian G, Bates Josephine T, Bertoia Lara R, Galford Amy E, Yoxthimer David A, and Riha Susan J. 2013. "Wastewater management and Marcellus Shale gas development: Trends, drivers, and planning implications." *Journal of environmental management*, 120 105–113. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
97. Rasmussen Sara G, Ogburn Elizabeth L, McCormack Meredith, Casey Joan A, Bandeen-Roche Karen, Mercer Dione G, and Schwartz Brian S. 2016. "Association between unconventional natural gas development in the Marcellus Shale and asthma exacerbations." *JAMA Internal Medicine*, 176(9): 1334–1343. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
98. Russell Rebecca B., Green Nancy S., Steiner Claudia A., Meikle Susan, Howse Jennifer L., Poschman Karalee, and Dias Todd et al. 2007. "Cost of hospitalization for preterm and low birth weight infants in the United States." *Pediatrics*, 120(1). [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
99. Sanders Nicholas J., and Stoecker Charles. 2015. "Where have all the young men gone? Using sex ratios to measure fetal death rates." *Journal of Health Economics*, 41 30–45, URL: <http://www.sciencedirect.com/science/article/pii/S0167629614001520>, DOI: 10.1016/j.jhealeco.2014.12.005. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
100. Schmidt Charles W. 2011. "Blind rush? Shale gas boom proceeds amid human health questions." *Environmental Health Perspectives*, 119(8): . [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
101. Severnini Edson. 2017. "Impacts of nuclear plant shutdown on coal-fired power generation and infant health in the Tennessee Valley in the 1980s." *Nature Energy*, 2(4):, p. 17051, URL: <https://www.nature.com/articles/nenergy201751>, DOI: 10.1038/nenergy.2017.51. [[DOI](#)] [[Google Scholar](#)]
102. Shah Prakesh S., and Balkhair Taiba. 2011. "Air pollution and birth outcomes: a systematic review." *Environment international*, 37(2):. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
103. Sram Radim J., Binkova Blanka, Dejmek Jan, and Bobak Martin. 2005. "Ambient air pollution and pregnancy outcomes: a review of the literature." *Environmental Health*

Perspectives, 113(4):. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

104. Stacy Shaina L. 2017. "A Review of the Human Health Impacts of Unconventional Natural Gas Development." *Current Epidemiology Reports*, 4(1): 38–45. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

105. Stacy Shaina L, Brink LuAnn L, Larkin Jacob C, Sadovsky Yoel, Goldstein Bernard D, Pitt Bruce R, and Talbott Evelyn O. 2015a. "Perinatal outcomes and unconventional natural gas operations in Southwest Pennsylvania." *PloS one*, 10(6):, p. e0126425. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

106. Stacy Shaina L., Brink LuAnn L., Larkin Jacob C., Sadovsky Yoel, Goldstein Bernard D., Pitt Bruce R., and Talbott Evelyn O.. 2015b.. "Perinatal Outcomes and Unconventional Natural Gas Operations in Southwest Pennsylvania." *PLOS ONE*, 10(6):, p. e0126425, URL: <http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0126425>, DOI: 10.1371/journal.pone.0126425. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

107. Stieb David M., Chen Li, Eshoul Maysoon, and Judek Stan. 2012a. "Ambient air pollution, birth weight and preterm birth: A systematic review and meta-analysis." *Environmental research*. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

108. Stieb David M., Chen Li, Eshoul Maysoon, and Judek Stan. 2012b. "Ambient air pollution, birth weight and preterm birth: A systematic review and meta-analysis." *Environmental Research*, 117 100–111, URL: <https://www.sciencedirect.com/science/article/pii/S0013935112001764>, DOI: 10.1016/j.envres.2012.05.007. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

109. Stillerman Karen Perry, Mattison Donald R., Giudice Linda C., and Woodruff Tracey J.. 2008. "Environmental exposures and adverse pregnancy outcomes: a review of the science." *Reproductive Sciences*, 15(7): 631–650. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]

110. Sun Xiaoli, Luo Xiping, Zhao Chunmei, Rachel Wai Chung Ng, Chi Eung Danforn Lim Bo Zhang, and Liu Tao. 2015. "The association between fine particulate matter exposure during pregnancy and preterm birth: a meta-analysis." *BMC Pregnancy and Childbirth*, 15, p. 300, URL: 10.1186/s12884-015-0738-2, DOI: <http://dx.doi.org/10.1186/s12884-015-0738-2>. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

111. Tanaka Shinsuke. 2015. "Environmental regulations on air pollution in China and their impact on infant mortality." *Journal of Health Economics*, 42 90–103, URL: <http://www.sciencedirect.com/science/article/pii/S0167629615000284>, DOI: <http://dx.doi.org/>

[10.1016/j.jhealeco.2015.02.004](https://doi.org/10.1016/j.jhealeco.2015.02.004) . [DOI] [PubMed] [Google Scholar]

112. Tustin Aaron W, Hirsch Annemarie G, Rasmussen Sara G, Casey Joan A, Bandeen-Roche Karen, and Schwartz Brian S. 2017. "Associations between unconventional natural gas development and nasal and sinus, migraine headache, and fatigue symptoms in Pennsylvania." *Environmental health perspectives*, 125(2): , p. 189. [DOI] [PMC free article] [PubMed] [Google Scholar]

113. Vengosh Avner, Robert B Jackson Nathaniel Warner, Darrah Thomas H, and Kondash Andrew. 2014. "A critical review of the risks to water resources from unconventional shale gas development and hydraulic fracturing in the United States." *Environmental science & technology*. [DOI] [PubMed] [Google Scholar]

114. Warner Nathaniel, Jackson Robert, Darrah Thomas, Osborn Stephen, Down Adrian, Zhao Kaiguang, White Alissa, and Vengosh Avner. 2012. "Geochemical Evidence for Possible Natural Migration of Marcellus Formation Brine to Shallow Aquifers in Pennsylvania." *Proceedings of the National Academy of Sciences*, 109(30): 11961–11966. [DOI] [PMC free article] [PubMed] [Google Scholar]

115. Warner Nathaniel R, Cidney A Christie, Jackson Robert B, and Vengosh Avner. 2013. "Impacts of shale gas wastewater disposal on water quality in Western Pennsylvania." *Environmental science & technology*, 47(20): 11849–11857. [DOI] [PubMed] [Google Scholar]

116. Werner Angela K, Sue Vink, Kerrienne Watt, and Paul Jagals. 2015. "Environmental health impacts of unconventional natural gas development: a review of the current strength of evidence." *Science of the Total Environment*, 505 1127–1141. [DOI] [PubMed] [Google Scholar]

117. Whitworth Kristina W., Marshall Amanda K., and Symanski Elaine. 2017. "Maternal residential proximity to unconventional gas development and perinatal outcomes among a diverse urban population in Texas." *PLOS ONE*, 12(7): , p. e0180966, URL: <http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0180966> , DOI: <http://dx.doi.org/10.1371/journal.pone.0180966>. [DOI] [PMC free article] [PubMed] [Google Scholar]

118. Whitworth Kristina Walker, Amanda Kaye Marshall, and Elaine Symanski., "Drilling and Production Activity Related to Unconventional Gas Development and Severity of Preterm Birth." *Environmental Health Perspectives*, 37006, p. 1. [DOI] [PMC free article] [PubMed] [Google Scholar]

119. WHO. 2005. "Effects of air pollution on children's health and development: a review of the evidence." [[Google Scholar](#)]
120. WHO. 2011. "WHO Weight Percentages Calculator." URL: http://www.who.int/entity/reproductivehealth/topics/best_practices/weight_percentiles_calculator.xls . [[Google Scholar](#)]
121. Witter Roxana, Lisa McKenzie Kaylan Stinson, Scott Kenneth, Newman Lee, and Adgate John. 2013. "The Use of Health Impact Assessment for a Community Undergoing Natural Gas Development." American Journal of Public Health. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
122. Wrenn Douglas H, Klaiber H Allen, and Jaenicke Edward C. 2016. "Unconventional Shale Gas Development, Risk Perceptions, and Averting Behavior: Evidence from Bottled Water Purchases." Journal of the Association of Environmental and Resource Economists, 3(4): 779–817. [[Google Scholar](#)]
123. Yan Beizhan, Stute Martin, Reynold A Panettieri James Ross, Mailloux Brian, Matthew J Neidell Lissa Soares, Howarth Marilyn, Liu Xinhua, Poun'e Saberi et al. 2016. "Association of groundwater constituents with topography and distance to unconventional gas wells in NE Pennsylvania." Science of The Total Environment. [[DOI](#)] [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
124. Yang Muzhe, Bhatta Rhea A., Chou Shin-Yi, and Cheng-I Hsieh. 2017. "The Impact of Prenatal Exposure to Power Plant Emissions on Birth Weight: Evidence from a Pennsylvania Power Plant Located Upwind of New Jersey." Journal of Policy Analysis and Management: [the Journal of the Association for Public Policy Analysis and Management], 36(3): 557–83. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]
125. Yang Muzhe, and Chou Shin-Yi. 2017. "The Impact of Environmental Regulation on Fetal Health: Evidence from the Shutdown of a Coal-Fired Power Plant Located Upwind of New Jersey." Journal of Environmental Economics and Management. [[Google Scholar](#)]
126. Zahran Sammy, Weiler Stephan, Mielke Howard W., and Alves Pena Anita. 2012. "Maternal benzene exposure and low birth weight risk in the United States: A natural experiment in gasoline reformulation." Environmental research, 112 139–146. [[DOI](#)] [[PubMed](#)] [[Google Scholar](#)]