

Data Submitted (UTC 11): 7/9/2025 8:16:03 PM

First name: Michael

Last name: Garrity

Organization:

Title:

Comments: Please find our comments attached.

?

July 8, 2025

Amy Hass

District Ranger

Beartooth Ranger District

6811 Highway 212

RedcLodge, Montana 59068

Dear District Ranger Hass,

Thank you for this opportunity to comment. Please accept these comments from me on behalf of the Alliance for the Wild Rockies, Native Ecosystems Council, Council on Wildlife and Fish, and the Center for Biological Diversity on the proposed Burnt Mountain Project.

We believe because of the size of the project and the cumulative effects of past current and future logging by the Forest Service and private logging and mining in the area the Forest Service must complete a full environmental impact statement (EIS) for this Project.

1

The scope of the Project will likely have a significant individual and cumulative impact on the environment. Alliance has reviewed the statutory and regulatory requirements governing National Forest Management projects, as well as the relevant case law, and compiled a check- list of issues that must be included in the EIS for the Project in order for the Forest Service's analysis to comply with the law. Following the list of necessary elements, Alliance has also included a general narrative discussion on possible impacts of the Project.

I. NECESSARY ELEMENTS FOR PROJECT EIS OR and EA
IF YOU REFUSE TO WRITE AN EIS: ?

?

A. Disclose all Custer-Gallatin National Forest Plan requirements for logging/burning projects and explain how the Project complies with them;

B. Will this project comply with forest plan big game hiding cover standards?

2

C. Disclose the acreages of past, current, and reasonably foreseeable logging, grazing, and road building activities within the Project area;

D. Solicit and disclose comments from the Montana Department of Fish, Wildlife, and Parks regarding the impact of the Project on wildlife habitat;

E. Solicit and disclose comments from the Montana Department

of Environmental Quality regarding the impact of the Project on water quality;

F. Disclose the biological assessment for the candidate, threatened, or endangered species with potential and/or actual habitat in the Project area;

G. Disclose the biological evaluation for the sensitive and management indicator species with potential and/or actual habitat in the Project area;

3

H. Disclose the snag densities in the Project area, and the method used to determine those densities;

I. Disclose the current, during-project, and post-project road densities in the Project area;

J. Disclose the Custer-Gallatin National Forest's record of compliance with state best management practices regarding stream sedimentation from ground-disturbing management activities;

K. Disclose the Custer-Gallatin National Forest's record of compliance with its monitoring requirements as set forth in its Forest Plan;

L. Disclose the Custer-Gallatin National Forest's record of compliance with the additional monitoring requirements set forth in previous DN/FONSIs and RODs on the Custer-Gallatin National Forest;

4

M. Disclose the results of the field surveys for threatened, endangered, sensitive, and rare plants in each of the proposed units;

N. Please formally consult with the US FWS on the impacts of this project on candidate, threatened, or endangered species and plants;

O. Please consult with the US FWS on the impacts of this project on lynx critical habitat and potential lynx critical habitat; (The US District Court just ordered the US FWS to redo their designation of lynx critical based on where lynx were in 2000 when lynx were listed.

P. Will this Project exacerbate existing noxious weed infestations and start new infestations?

Q. Do unlogged old growth forest store more carbon than the wood products that would be removed from the same forest in a logging operation?

5

R. What is the cumulative effect of National Forest logging on U.S. carbon stores? How many acres of National Forest lands are logged every year? How much carbon is lost by that logging?

S. Is this Project consistent with "research recommendations (Krankina and Harmon 2006) for protecting carbon gains against the potential impacts of future climate change? That study recommends "[i]ncreasing or maintaining the forest area by avoiding de-forestation," and states that "protecting forest from logging or clearing offer immediate benefits via prevented emis-

sions." That study also states that "[w]hen the initial condition of land is a productive old-growth forest, the conversion to forest plantations with a short harvest rotation can have the opposite effect lasting for many decades" The study does state that thinning may have a beneficial effect to stabilize the forest and avoid stand-

6

replacing wildfire, but the study never defines thinning. In this Project, where much of the logging is clear-cutting and includes removing large trees without any diameter limit, and where the removal of small diameter surface and ladder fuels is an unfunded mandate to the tune of over \$3 million dollars, it is dubious whether the prescriptions are the same type of "thinning" envisioned in Krankina and Harmon (2006).

T. Please list each visual quality standard that applies to each unit and disclose whether each unit meets its respective visual quality standard. A failure to comply with visual quality Forest Plan standards violates NFMA.

U. For the visual quality standard analysis please define "ground vegetation," i.e. what age are the trees, "reestablishes," "short-term," "longer term," and "revegetate."

V. Please disclose whether you have conducted surveys in the Project area for this Project for wolverines, grizzly bears, pine

7

martins, northern goshawk, whitebark pine, monarch butterflies, and lynx as required by the Forest Plan.

W. Please disclose how often the Project area has been surveyed for wolverines, pine martins, northern goshawks, grizzlies, whitebark pine, monarch butterflies, and lynx.

X. Is it impossible for wolverines, pine martins, northern goshawks, grizzlies, whitebark pine, monarch butterflies, to inhabit the Project area?

Y. Would the habitat be better for wolverines, pine martins, northern goshawks, grizzlies, whitebark pine, monarch butterflies, if roads were removed in the Project area?

Z. What is the U.S. FWS position on the impacts of this Project on wolverines, pine martins, northern goshawks, grizzlies, whitebark pine, monarch butterflies, and lynx? Have you conducted ESA consultation?

8

AA. Please provide us with the full BA for the wolverines, pine martins, northern goshawks, grizzlies, whitebark pine, monarch butterflies, and lynx.

BB. What is wrong with uniform forest conditions? ?

?

CC. Has the beetle kill contributed to a diverse landscape?

DD. Why are you trying to exclude stand replacement fires when these fires help aspen and whitebark pine?

EE. Please disclose what is the best available science for restoration of whitebark pine.

FF. Disclose the level of current noxious weed infestations in the

Project area and the cause of those infestations;

GG. Disclose the impact of the Project on noxious weed infestations and native plant communities;

9

HH. Disclose the amount of detrimental soil disturbance that currently exists in each proposed unit from previous logging and grazing activities;

II. Disclose the expected amount of detrimental soil disturbance in each unit after ground disturbance and prior to any proposed mitigation/remediation;

JJ. Disclose the expected amount of detrimental soil disturbance in each unit after proposed mitigation/remediation;

KK. Disclose the analytical data that supports proposed soil mitigation/ remediation measures;

LL. Disclose the timeline for implementation;

MM. Disclose the funding source for non- commercial activities proposed;

NN. Disclose the current level of old growth forest in each third order drainage in the Project area;

10

OO. Disclose the method used to quantify old growth forest acreages and its rate of error based upon field review of its predictions;

PP. Disclose the historic levels of mature and old growth forest in the Project area;

QQ. Disclose the level of mature and old growth forest necessary to sustain viable populations of dependent wildlife species in the area;

RR. Disclose the amount of mature and old growth forest that will remain after implementation;

SS. Disclose the amount of current habitat for old growth and mature forest dependent species in the Project area;

TT. Disclose the amount of habitat for old growth and mature forest dependent species that will remain after Project implementation;

11

UU. Disclose the method used to model old growth and mature forest dependent wildlife habitat acreages and its rate of error based upon field review of its predictions;

VV. Disclose the amount of big game (moose and elk) hiding cover, winter range, and security currently available in the area;

WW. Disclose the amount of big game (moose and elk) hiding cover, winter range, and security cover during Project implementation;

XX. Disclose the amount of big game (moose and elk) hiding cover, winter range, and security after implementation;

YY. Disclose the method used to determine big game hiding cover, winter range, and security, and its rate of error as determined by field review;

ZZ. Disclose and address the concerns expressed by the ID Team in the draft Five-Year Review of the Forest Plan regar-

ding the failure to monitor population trends of MIS, the inade-
12

quacy of the Forest Plan old growth standard, and the failure to compile data to establish a reliable inventory of sensitive species on the Forest;

AAA. Disclose the actions being taken to reduce fuels on private lands adjacent to the Project area and how those activities/or lack thereof will impact the efficacy of the activities proposed for this Project;

BBB. Disclose the efficacy of the proposed activities at reducing wildfire risk and severity in the Project area in the future, including a two-year, five-year, ten-year, and 20-year projection;

CCC. Disclose when and how the Custer Gallatin National Forest made the decision to suppress natural wildfire in the Project area and replace natural fire with logging and prescribed burning;

13

DDD. Disclose the cumulative impacts on the Forest-wide level of the Custer Gallatin National Forest's policy decision to replace natural fire with logging and prescribed burning;

EEE. Disclose how Project complies with the Roadless Rule;

FFF. Disclose the impact of climate change on the efficacy of the proposed treatments;

GGG. Disclose the impact of the proposed project on the carbon storage potential of the area;

HHH. Disclose the baseline condition, and expected sedimentation during and after activities, for all streams in the area;

III. Disclose maps of the area that show the following elements:

1. Past, current, and reasonably foreseeable logging units in the Project area;

2. Past, current, and reasonably foreseeable grazing allotments in the Project area;

14

3. Density of human residences within 1.5 miles from the Project unit boundaries;

4. Hiding cover in the Project area according to the Forest Plan definition;

5. Old growth forest in the Project area;

6. Big game security areas;

7. Moose winter range;

The best available science, Christensen et al (1993), recommends elk habitat effectiveness of 70% in summer range and at least 50% in all other areas where elk are one of the primary resource considerations. According to Figure 1 in Christensen et al (1993), this equates to a maximum road density of approximately 0.7 mi/sq mi. in summer range and approximately 1.7 mi/sq mi. in all other areas.

Do any of the 6th Code watersheds in the Project area meet either of these road density thresholds? It appears the Project area

15

as a whole also far exceeds these thresholds. Please disclose this

type of Project level or watershed analysis on road density. Christensen et al (1993) state that if an area is not meeting the 50% effectiveness threshold of 1.7 mi/sq mi, the agency should admit that the area is not being managed for elk: "Areas where habitat effectiveness is retained at lower than 50 percent must be recognized as making only minor contributions to elk management goals. If habitat effectiveness is not important, don't fake it. Just admit up front that elk are not a consideration." The Project EIS does not make this admission.

The Forest Service should provide an analysis of how much of the Project area, Project area watersheds, affected landscape areas, or affected Hunting Districts provide "elk security area[s]" as defined by the best available science, Christensen et al (1993) and Hillis et al (1991), to be comprised of contiguous 250 acre blocks of forested habitat 0.5 miles or more from open

16
roads with these blocks encompassing 30% or more of the area.

Please provide a rational justification for the deviation from the Hillis security definition and numeric threshold that represent the best available science on elk security areas.

We believe that best available science shows that Commercial Logging does not reduce the threat of Forest Fires. What best available science supports the action alternatives?

Please find Schoennagel et al (2004) attached. Schoennagel states: "we are concerned that the model of historical fire effects and 20th-century fire suppression in dry ponderosa pine forests is being applied uncritically across all Rocky Mountain forests, including where it is inappropriate.

Schoennagel et al (2004) states: "High-elevation subalpine forests in the Rocky Mountains typify ecosystems that experience infrequent, high-severity crown fires []. . . The most extensive

17
subalpine forest types are composed of Engelmann spruce (*Picea engelmannii*), subalpine fir (*Abies lasiocarpa*), and lodgepole pine (*Pinus contorta*), all thin-barked trees easily killed by fire. Extensive stand-replacing fires occurred historically at long intervals (i.e., one to many centuries) in subalpine forests, typically in association with infrequent high-pressure blocking systems that promote extremely dry regional climate patterns."

Schoennagel et al (2004) states: "it is unlikely that the short period of fire exclusion has significantly altered the long fire intervals in subalpine forests. Furthermore, large, intense fires burning under dry conditions are very difficult, if not impossible, to suppress, and such fires account for the majority of area burned in subalpine forests.

Schoennagel et al (2004) states: "Moreover, there is no consistent relationship between time elapsed since the last fire and fuel abundance in subalpine forests, further undermining the idea

18

that years of fire suppression have caused unnatural fuel buildup in this forest zone."

Schoennagel et al (2004) states: "No evidence suggests that spruce-fir or lodgepole pine forests have experienced substantial shifts in stand structure over recent decades as a result of fire suppression. Overall, variation in climate rather than in fuels appears to exert the largest influence on the size, timing, and severity of fires in subalpine forests []. We conclude that large, infrequent stand replacing fires are 'business as usual' in this forest type, not an artifact of fire suppression."

Schoennagel et al (2004) states: "Contrary to popular opinion, previous fire suppression, which was consistently effective from about 1950 through 1972, had only a minimal effect on the large fire event in 1988 []. Reconstruction of historical fires indicates that similar large, high-severity fires also occurred in the early 1700s []. Given the historical range of variability of fire regimes in high-elevation subalpine forests, fire behavior in Yellow-

stone during 1988, although severe, was neither unusual nor surprising."

Schoennagel et al (2004)(emphasis added) states: "Mechanical fuel reduction in sub-alpine forests would not represent a restoration treatment but rather a departure from the natural range of variability in stand structure."

Schoennagel et al (2004) states: "Given the behavior of fire in Yellowstone in 1988, fuel reduction projects probably will not substantially reduce the frequency, size, or severity of wildfires under extreme weather conditions."

Schoennagel et al (2004) states: "The Yellowstone fires in 1988 revealed that variation in fuel conditions, as measured by stand age and density, had only minimal influence on fire behavior. Therefore, we expect fuel-reduction treatments in high-elevation forests to be generally unsuccessful in reducing fire frequency, severity, and size, given the overriding importance of

extreme climate in controlling fire regimes in this zone. Thinning also will not restore subalpine forests, because they were dense historically and have not changed significantly in response to fire suppression. Thus, fuel-reduction efforts in most Rocky Mountain sub-alpine forests probably would not effectively mitigate the fire hazard, and these efforts may create new ecological problems by moving the forest structure outside the historic range of variability."

Likewise, Brown et al (2004) states: "At higher elevations, forests of subalpine fir, Engelmann spruce, mountain hemlock, and lodgepole or whitebark pine predominate. These forests also have long fire return intervals and contain a high proportion of fire sensitive trees. At periods averaging a few hundred years, extreme drought conditions would prime these forests for large, severe fires that would tend to set the forest back to an early successional stage, with a large carry-over of dead trees as

a legacy of snags and logs in the regenerating forest . . . natural

21

ecological dynamics are largely preserved because fire suppression has been effective for less than one natural fire cycle. Thinning for restoration does not appear to be appropriate in these forests. Efforts to manipulate stand structures to reduce fire hazard will not only be of limited effectiveness but may also move systems away from pre-1850 conditions to the detriment of wildlife and watersheds." "Fuel levels may suggest a high fire 'hazard' under conventional assessments, but wildfire risk is typically low in these settings."

Likewise, Graham et al (2004) states: "Most important, the fire behavior characteristics are strikingly different for cold (for example, lodgepole pine, Engelmann spruce, subalpine fir), moist (for example, western hemlock, western redcedar, western white pine), and dry forests. Cold and moist forests tend to have long fire-return intervals, but fires that do occur tend to be high-intensity, stand-replacing fires.

22

Dry forests historically had short intervals between fires, but most important, the fires had low to moderate severity." According to Graham et al (2004), thinning may also increase the likelihood of wildfire ignition in the type of forests in this Project area: "The probability of ignition is strongly related to fine fuel moisture content, air temperature, the amount of shading of surface fuels, and the occurrence of an ignition source (human or lightning caused) . . . There is generally a warmer, dryer microclimate in more open stands (fig. 9) compared to denser stands. Dense stands (canopy cover) tend to provide more shading of fuels, keeping relative humidity higher and air and fuel temperature lower than in more open stands. Thus, dense stands tend to maintain higher surface fuel moisture contents compared to more open stands. More open stands also tend to allow higher wind speeds that tend to dry fuels compared to dense stands. These factors may increase

23

probability of ignition in some open canopy stands compared to dense canopy stands."

Please see the attached study by DellaSala et. al. 2022 reviewed 1500 wildfires between 1984 and 2014 found that actively managed forests had the highest level of fire severity. While those forests in protected areas burned, on average, had the lowest level of fire severity. In other words, the best way to reduce severe fires is to protect the land as wilderness, not "manage" it. A study of over 1,500 fires covering 23 million acres by Bradley et al. 2016, (attached) concluded the more "thinning" of trees in a forest the more quickly and intensely a wildfire burns. One author writes, "Dense, mature forests tend to burn less intensely ... because they have higher canopy cover and more shade, which creates a cooler, more moist microclimate. The higher density of trees of all sizes can act as a windbreak, buffering gust-driven

flames. Thinning and other activities that remove trees, especially mature trees, reverse those effects, creating hotter, drier, and

24

windier conditions." Even a USFS review of the largest fire in New Mexico history, a mismanaged prescribed burn, agreed. Please see the attached paper by Schoennagel et al. 2017, concluded that "fuels reduction cannot alter regional wildfire trends."

Intentional burning produces massive amounts of deadly air pollution. Please see the attached paper by Maji et al. 2024 concluded America's "prescribed burns" annually cause about 10,000 premature deaths and \$100 billion in health costs, about the same as from wildfires.

How many premature deaths will this project cause and how much will it increase health care costs?

The Project will violate the NEPA if there are no valid snag surveys done for the project area both within and outside proposed harvest units.

The project will violate the NEPA if there are no valid surveys for old growth habitat within each project area, as identified by Green et al. 1992; old growth types need to be defined and quan-

25

tified by timber types, such as lodgepole pine, Douglas-fir, mixed conifer, spruce, subalpine fir, and limber pine.

Please demonstrate the project will maximize the retention of old-growth and large trees as required by Section 605 of the Healthy Forest Restoration Act.

The project will likely violate the NEPA if the mitigation measures for MIS, sensitive species, and Montana Species of Concern (birds, mammals including bats) are not clearly defined, and demonstrated to be effective as per the current best science. We request a careful analysis of the impacts to fisheries and water quality, including considerations of sedimentation, increases in peak flow, channel stability, risk of rain-on-snow events, and increases in stream water temperature. Please disclose the locations of seeps, springs, bogs and other sensitive wet areas, and the effects on these areas of the project activities. Where livestock are permitted to graze, we ask that you assess the

26

present condition and continue to monitor the impacts of grazing activities upon vegetation diversity, soil compaction, stream bank stability and subsequent sedimentation. Livestock grazing occurs in the Project area and causes sediment impacts, trampled or destabilized banks, increased nutrient loads in streams, and decreased density, diversity, and function of riparian vegetation that may lead to increased stream temperatures and further detrimental impacts to water quality.

The Project will violate the NEPA if there are no valid snag surveys done for the project area both within and outside proposed harvest units.

The project will violate the NEPA if there are no valid surveys

for old growth habitat within each project area, as identified by Green et al. 1992; old growth types need to be defined and quantified by timber types, such as lodgepole pine, Douglas-fir, mixed conifer, spruce, subalpine fir, and limber pine.

27

The project will likely violate the NEPA if the mitigation measures for MIS, sensitive species, and Montana Species of Concern (birds, mammals including bats) are not clearly defined, and demonstrated to be effective as per the current best science.

FAILURE TO REVIEW AND PROTECT CULTURAL AND HISTORICAL RESOURCES

Consultation with the State Historic Preservation Office (SHPO) must be completed prior to a decision being signed.

Any required protection measures provided from SHPO will be incorporated into my final decision.

Crucial to the preservation of the historical and cultural foundations of the nation, Section 106 of the National Historic Preservation Act (NHPA) and its implementing regulations, 36 C.F.R. Part 800 (PDF) (revised August 5, 2004) require Federal agencies to consider the effects of projects they carry out, approve, or fund on historic properties. Additionally, Federal agencies must

28

provide the Advisory Council on Historic Preservation (ACHP) opportunity to comment on such projects prior to the agency's final decision.

A Federal project that requires review under Section 106 is defined as an "undertaking." An undertaking means a project, activity or program funded in whole or in part under the direct or indirect jurisdiction of a Federal agency, including those carried out by or on behalf of a Federal agency; those carried out with Federal financial assistance; and those requiring a Federal permit, license, or approval.

Section 110 of the NHPA

Added to the NHPA in 1992, Section 110 requires Federal agencies to emphasize the preservation and enhancement of cultural resources. Section 110 directs agencies to initiate measures necessary to direct their policies, plans, and programs in such a way that federally-owned sites, structures, and objects of histori-

29

cal architectural or archaeological significance are preserved, restored, and maintained for the inspiration and benefit of the public. The agencies are also encouraged to institute (in consultation with the ACHP) procedures to assure Federal plans and programs contribute to the preservation and enhancement of non-Federally owned sites, structures, and objects of historical, architectural, and archaeological significance. Has the MT SHPO received this survey? The cultural surveys need to be done before the NEPA and NHPA process can be completed, which has not occurred. The project must be approved by the SHPO and the public needs to be given a chance to comment on this.

Did the Forest Service conduct NEPA analysis (i.e. an EA or EIS) for the Fire Plan the Forest is using for this project? If you don't the project will be in violation of NEPA, NFMA, and the APA.

30

Please provide a map showing the Wildland Urban Interface (WUI) and the locations of all homes in comparison to the project area. Please demonstrate that the WUI complies with the definition of the WUI found in the Healthy Forest Restoration Act.

Since the Forest Service did not conduct NEPA for the Fire Plan, please disclose the cumulative effects of Forest-wide implementation of the Fire Plan in the project EIS, or EA if you refuse to write an EIS, to avoid illegally tiering to a non- NEPA document. Specifically analyze the decision to prioritize mechanical, human-designed, somewhat arbitrary treatments as a replacement for naturally-occurring fire.

Did the Forest Service conduct ESA consultation for the Fire Plan?

31

Will the Forest Service be considering binding legal standards for noxious weeds in its revision of the Custer Gallatin Forest Plan?

How effective have BMPs been at stopping (i.e. preventing) new weed infestations from starting during logging and related road operations?

Is it true that new roads are the number one cause of new noxious weed infestations?

Why isn't the Forest Service considering a Forest Plan amendment in this Project to amend the Forest Plan to include binding legal standards that address noxious weeds?

Is it true that noxious weeds are one of the top threats to biodiversity on our National Forests?

How can the Forest Service be complying with NFMA's requirement to maintain biodiversity if it has no legal standards that address noxious weeds?

32

Will this Project address all Project area BMP needs, i.e. will the BMP road maintenance backlog and needs from this Project all be met by this Project?

The scoping notice was not clear if any MIS were found. What MIS did you find, how many and how did you look for these MIS?

How will the decreased elk security and thermal cover affect wolverines? Please formally consult with the US FWS on the impact of this project on wolverines. The U.S. District Court ordered the USFWS to reconsider if wolverines should be listed under ESA. Wolverines need secure habitat in big game winter range.

Which wildlife species and ecosystem processes, if any, does the fire-proofing in the proposed project benefit? Which species and

processes do fire-proofing harm?

What is your definition of healthier?

33

What evidence do you have that this logging will make the forest healthier for fish and wildlife? What about the role of mixed severity and high severity fire - what are the benefits of those natural processes?

How have those processes (mixed and high severity fire) created the ecosystems we have today?

Over how many millennia have mixed and high severity fire have been occurring without human intervention?

What beneficial ecological roles do beetles play? You didn't answer this in violation of NEPA, NFMA and the APA.

Can the forest survive without beetles?

Will all WQLS streams in the project area have completed TMDLs before a decision is signed?

Will this project leave enough snags to follow the Forest Plan requirements and the requirements of sensitive old growth species such as flammulated owls and goshawks?

34

Will this Project exacerbate existing noxious weed infestations and start new infestations?

Do unlogged old growth forests store more carbon than the wood products that would be removed from the same forest in a logging operation?

What is the cumulative effect of National Forest logging on U.S. carbon stores? How many acres of National Forest lands are logged every year? How much carbon is lost by that logging?

Is this Project consistent with "research recommendations (Krankina and Harmon 2006) for protecting carbon gains against the potential impacts of future climate change? That study recommends "[i]ncreasing or maintaining the forest area by avoiding deforestation," and states that "protecting forest from logging or clearing offer immediate benefits via prevented emissions."

35

Please list each visual quality standard that applies to each unit and disclose whether each unit meets its respective visual quality standard.

Weeds

Native plants are the foundation upon which the ecosystems of the Forest are built, providing forage and shelter for all native wildlife, bird and insect species, supporting the natural processes of the landscape, and providing the context within which the public find recreational and spiritual opportunities. All these uses or values of land are hindered or lost by conversion of native vegetation to invasive and noxious plants. The ecological threats posed by noxious weed infestations are so great that a former chief of the Forest Service called the invasion of noxious weeds "devastating" and a "biological disaster." Despite implementation of Forest Service "best management practices"

(BMPs), noxious weed infestation on the Forest is getting worse and noxious weeds will likely overtake native plant populations
36

if introduced into areas that are not yet infested. The Forest Service has recognized that the effects of noxious weed invasions may be irreversible. Even if weeds are eliminated with herbicide treatment, they may be replaced by other weeds, not by native plant species.

Invasive plant species, also called noxious weeds, are one of the greatest modern threats to biodiversity on earth. Noxious weeds cause harm because they displace native plants, resulting in a loss of diversity and a change in the structure of a plant community. By removing native vegetative cover, invasive plants like knapweed may increase sediment yield and surface runoff in an ecosystem. As well knapweed may alter organic matter distribution and nutrient through a greater ability to uptake phosphorus over some native species in grasslands. Weed colonization can alter fire behavior by increasing flammability: for example, cheatgrass, a widespread noxious weed on the Forest, cures early and leads to more frequent burning. Weed colonization can
37

also deplete soil nutrients and change the physical structure of soils.

The Forest Service's own management activities are largely responsible for noxious weed infestations; in particular, logging, prescribed burns, and road construction and use create a risk of weed infestations. The introduction of logging equipment into the Forest creates and exacerbates noxious weed infestations. The removal of trees through logging can also facilitate the establishment of noxious weed infestations because of soil disturbance and the reduction of canopy closure. In general, noxious weeds occur in old clearcuts and forest openings, but are rare in mature and old growth forests. Roads are often the first place new invader weeds are introduced. Vehicle traffic and soil disturbances from road construction and maintenance create ideal establishment conditions for weeds. Roads also provide obvious dispersal corridors. Roadsides throughout the project area are infested with noxious weeds. Once established along roadsides,
38

invasive plants will likely spread into adjacent grasslands and forest openings.

Prescribed burning activities within the analysis area would likely cumulatively contribute to increases in noxious weed distribution and populations. As a disturbance process, fire has the potential to greatly exacerbate infestations of certain noxious weed species, depending on burn severity and habitat type (Fire Effects Information System 2004). Soil disturbance, such as that resulting from low and moderate burn severities from prescribed fire and fire suppression related disturbances (dozer lines, drop spots, etc.), provide optimum conditions for noxious weed invasion. Dry site vegetation types and road corridors are extremely

vulnerable, especially where recent ground disturbance (timber management, road construction) has occurred. Units proposed for burning within project area may have closed forest service access roads (jammers) located within units. These units have the highest potential for noxious weed infestation and exacerbation through fire activities. Please provide an alternative that eliminates units that have noxious weeds present on roads within units from fire management proposals.

Please address the ecological, social and aesthetic impact of current noxious weed infestations within the project area. Include an analysis of the impact of the actions proposed by this project on the long and short term spread of current and new noxious weed infestations. What treatment methods will be used to address growing noxious weed problems? What noxious weeds are currently and historically found within the project area? Please include a map of current noxious weed infestations which includes knapweed, Saint Johnswort, cheat grass, bull thistle, Canada thistle, hawkweed, hound's-tongue, oxeye daisy and all other Category 1, Category 2 and Category 3 weeds classified as noxious in the MONTANA COUNTY NOXIOUS WEED LIST. State-listed Category 2 noxious weed species yellow and orange hawkweeds are recently established (within the last 5 to 10 years) in Montana and are rapidly expanding in established areas. They can invade undisturbed areas where native plant communities are intact. These species can persist in shaded conditions and often grow underneath shrubs making eradication very difficult. Their stoloniferous (growing at the surface or below ground) habit can create dense mats that can persist and spread to densities of 3500 plants per square mile (Thomas and Dale 1975). Are yellow and orange hawkweeds present within the project area?

Please address the cumulative, direct and indirect effects of the proposed project on weed introduction, spread and persistence that includes how weed infestations have been and will be influenced by the following management actions: road construction including new permanent and temporary roads and skid trails proposed within this project; opening and decommissioning of roads represented on forest service maps; ground disturbance and traffic on forest service template roads, mining access routes, and private roads; removal of trees through commercial and pre-commercial logging and understory thinning; and prescribed burns. What open, gated, and decommissioned Forest Service roads within the project area proposed as haul routes have existent noxious weed populations and what methods will be used to assure that noxious weeds are not spread into the proposed action units?

Noxious weeds are not eradicated with single herbicide treatments. A onetime application may kill an individual plant but

dormant seeds in the ground can still sprout after herbicide treatment. Thus, herbicides must be used on consistent, repetitive schedules to be effective.

What commitment to a long-term, consistent strategy of application is being proposed for each weed infested area within the proposed action area? What long term monitoring of weed populations is proposed?

42

When areas treated with herbicides are re-seeded on national forest land, they are usually reseeded with exotic grasses, not native plant species. What native plant restoration activities will be implemented in areas disturbed by the actions proposed in this project? Will disturbed areas including road corridors, skid trails, and burn units be planted or reseeded with native plant species?

The scientific and managerial consensus is that prevention is the most effective way to manage noxious weeds. The Forest Service concedes that preventing the introduction of weeds into un-infested areas is "the most critical component of a weed management program." The Forest Service's national management strategy for noxious weeds also recommends "develop[ing] and implement[ing] forest plan standards" and recognizes that the cheapest and most effective solution is prevention. Which units within the project area currently have no noxious weed populations within their boundaries? What minimum standards

43

are in the Custer Gallatin National Forest Plan to address noxious weed infestations? Please include an alternative in the DEIS that includes land management standards that will prevent new weed infestations by addressing the causes of weed infestation. The failure to include preventive standards violates NFMA because the Forest Service is not ensuring the protection of soils and native plant communities. Additionally, the omission of an EIS alternative that includes preventive measures would violate NEPA because the Forest Service would fail to consider a reasonable alternative.

Rare Plants

The ESA requires that the Forest Service conserve endangered and threatened species of plants as well as animals. In addition to plants protected under the ESA, the Forest Service identifies species for which population viability is a concern as "sensitive species" designated by the Regional Forester (FSM 2670.44). The response of each of the sensitive plant species to manage-

44

ment activity varies by species, and in some cases, is not fully known. Local native vegetation has evolved with and is adapted to the climate, soils, and natural processes such as fire, insect and disease infestations, and windthrow. Any management or lack of management that causes these natural processes to be altered may have impacts on native vegetation, including threatened and sensitive plants. Herbicide application - intended to

eradicate invasive plants - also results in a loss of native plant diversity because herbicides kill native plants as well as invasive plants.

Whitebark Pine ?

?

Not all ecosystems or all Rocky Mountain landscapes have experienced the impacts of fire exclusion. In some wilderness areas, where in recent decades natural fires have been allowed to burn, there have not been major shifts in vegetation composition and structure (Keane et al. 2002). In some alpine ecosystems, 45

fire was never an important ecological factor. In some upper subalpine ecosystems, fires were important, but their rate of occurrence was too low to have been significantly altered by the relatively short period of fire suppression (Keane et al. 2002). For example, the last 70 to 80 years of fire suppression have not had much influence on subalpine landscapes with fire intervals of 200 to several hundred years (Romme and Despain). Consequently, it is unlikely that fire exclusion has yet to significantly alter stand conditions or forest health within Rocky Mountain sub-alpine ecosystems. ?

Whitebark pine seedlings, saplings and mature trees, present in subalpine forests proposed for burning, would experience mortality from project activity. Whitebark pine is fire intolerant (thin bark). Fire favors whitebark pine regeneration (through canopy opening and reducing competing vegetation) only in the presence of adequate seed source and dispersal mechanisms (Clarks 46

Nutcracker or humans planting white-bark pine seedlings). White pine blister rust, an introduced disease, has caused rapid mortality of whitebark pine over the last 30 to 60 years. Keane and Arno (1993) reported that 42 percent of whitebark pine in western Montana had died in the previous 20 years with 89 percent of remaining trees being infected with blister rust. The ability of whitebark pine to reproduce naturally is strongly affected by blister rust infection; the rust kills branches in the upper cone bearing crown, effectively ending seed production.

Montana is currently experiencing a mountain pine beetle epidemic. Mountain pine beetle prefer large, older whitebark pine, which are the major cone producers. In some areas the few remaining whitebark that show the potential for blister rust resistance are being attacked and killed by mountain pine beetles, thus accelerating the loss of key mature cone-bearing trees.

47

Whitebark pine seedlings and saplings are very likely present in the subalpine forests proposed for burning and logging. In the absence of fire, this naturally occurring white-bark pine regeneration would continue to function as an important part of the subalpine ecosystem. Since 2005, rust resistant seed sources have been identified in the Northern Rockies (Mahalovich et al

2006). Due to the severity of blister rust infection within the region, natural whitebark pine regeneration in the project area is prospective rust resistant stock.

Although prescribed burning can be useful to reduce areas of high-density subalpine fir and spruce and can create favorable ecological conditions for whitebark pine regeneration and growth, in the absence of sufficient seed source for natural regeneration maintaining the viability and function of whitebark pine would not be achieved through burning.

Planting of rust-resistant seedlings would likely not be sufficient to replace whitebark pine lost to fire activities.

48

What surveys have been conducted to determine presence and abundance of whitebark pine regeneration? If whitebark pine seedlings and saplings are present, what measures will be taken to protect them? Please include an alternative that excludes burning in the presence of whitebark pine regeneration (consider 'Daylighting' seedlings and saplings as an alternative restoration method). Will restoration efforts include planting whitebark pine? Will planted seedling be of rust-resistant stock? Is rust-resistant stock available? Would enough seedlings be planted to replace whitebark pine lost to fire activities? Have white pine blister rust surveys been accomplished? What is the severity of white pine blister rust in proposed action areas? Please see the attached paper by Six et al. 2021 about logging and white bark pine.

It is not clear how vast acres of national forest lands can be intensively manipulated without a single wildlife management objective identified. This means that the agency cannot be meeting

49

the NFMA requirements to maintain a diversity of wildlife. And it also means that the stated purpose and need is invalid, and a violation of the NEPA, because vegetation and fuels management have a direct impact on wildlife, which means that wildlife management also has to be included as a purpose and need. In effect, the purpose and need of the project is to remove wildlife habitat, but this is never identified to the public, in violation of the NEPA.

The agency is violating the NEPA by using vague, unmeasurable terms to rationalize the proposed logging to the public.

How can the public measure "resiliency?" What are the specific criteria used to define resiliency, and what are the ratings for each proposed logging unit before and after treatment? How is the risk of fire as affected by the project being measured so that the public can understand whether or not this will be effective? How is forest health to be measured so that the public can see that this is a valid management strategy? What specifically con-

50

stitutes a diversity of age classes, how is this to be measured, and how are proposed changes measured as per diversity? How are diversity measures related to wildlife (why is diversity need-

ed for what species)? If the reasons for logging cannot be clearly identified and measured for the public, the agency is not meeting the NEPA requirements for transparency.

Please find attached the paper by Faison et al. 2023 titled, "The importance of natural forest stewardship in adaptation planning in the United States." They "argue that expensive management interventions are often unnecessary, have uncertain benefits, or are detrimental to many forest attributes such as resilience, carbon accumulation, structural complexity, and genetic and biological diversity. Natural forests (i.e., those protected and largely free from human management) tend to develop greater complexity, carbon storage, and tree diversity over time than forests that are actively managed; and natural forests often become less susceptible to future insect attacks and fire following these

51

disturbances. Natural forest stewardship is therefore a critical and cost effective strategy in forest climate adaptation."

Faison et al. 2023 shows that the project is not meeting the purpose and need of the project. Please find Faison et al. 2023 attached.

Please see the paper by Baker et al. 2023. This landmark study found a pattern of "Falsification of the Scientific Record" in government-funded wildfire studies.

Dr. Baker's paper is the best available science. Please explain why this project is not following the best available science. The Draft Decision Notice is in violation of NEPA.

This unprecedented study was published in the peer-reviewed journal Fire, exposing a broad pattern of scientific misrepresentations and omissions that have caused a "falsification of the scientific record" in recent forest and wildfire studies funded or authored by the U.S. Forest Service with regard to dry forests of

52

the western U.S. Forest Service related articles have presented a falsified narrative that historical forests had low tree densities and were dominated by low-severity fires, using this narrative to advocate for its current forest management and wildfire policies. However, the new study comprehensively documents that a vast body of scientific evidence in peer-reviewed studies that have directly refuted and discredited this narrative were either misrepresented or omitted by agency publications. The corrected scientific record, based on all of the evidence, shows that historical forests were highly variable in tree density, and included "open" forests as well as many dense forests. Further, historical wildfire severity was mixed and naturally included a substantial component of high-severity fire, which creates essential snag forest habitat for diverse native wildlife species, rivaling old-growth forests.

These findings have profound implications for climate mitigation and community safety, as current forest policies that are

53

driven by the distorted narrative result in forest management

policies that reduce forest carbon and increase carbon emissions, while diverting scarce federal resources from proven community wildfire safety measures like home hardening, defensible space pruning, and evacuation assistance.

"Forest policy must be informed by sound science but, unfortunately, the public has been receiving a biased and inaccurate presentation of the facts about forest density and wildfires from government agencies," said Dr. William Baker in their press release announcing the publication of their paper.

"The forest management policies being driven by this falsified scientific narrative are often making wildfires spread faster and more intensely toward communities, rather than helping communities become fire-safe," said Dr. Chad Hanson, research ecologist with the John Muir Project in the same press release.

"We need thinning of small trees adjacent to homes, not back-country management."

54

"The falsified narrative from government studies is leading to inappropriate forest policies that promote removal of mature, fire-resistant trees in older forests, which causes increased carbon emissions and in the long-run contributes to more fires" said, Dr. Dominick A. DellaSala, Chief Scientist, Wild Heritage, a Project of Earth Island Institute concluded in the press release. Please analyze how the project would effect wilderness quality lands.

Please see the attached map of the lands that would be protected in the Greater Yellowstone Ecosystem by the Northern Rockies Ecosystem Protection Act, S. 1198 (attached) and H.R. 2420 in the Greater Yellowstone Ecosystem. Would this project make any lands no longer qualify as wilderness?

Does the project area include any roadless lands? Section 605 of the Healthy Forest Recreation Act prohibits logging in inventoried Roadless Areas.

55

Please see the following article:

Montana researchers urge towns to focus on wildfire preparation

<https://missoulacurrent.com/research-wildfire-preparation/>

Laura Lundquist

(Missoula Current) For more than a decade, a small group of scientists have been trying to convince people that fireproofing their homes is far more effective than logging the forest when it comes to surviving wildfire. But few people are listening. In mid-December, six researchers published a paper in the Proceedings of the National Academy of Sciences journal warning that communities across the nation, but particularly those in the West, aren't prepared to survive an urban conflagration such as the one that devastated Lahaina, Hawaii, in August.

The paper, titled "Wildland-urban fire disasters aren't actually a wildfire problem," points out that, since 2016, communities

from Lahaina to Gatlinburg, Tenn., that have lost hundred of homes to fires have certain things in common: the fires occurred under extreme weather conditions - high winds and persistent drought - and most of the structures weren't fire-resistant.

"These problem fires were defined as an issue of wildfires that involved houses. In reality, they are urban fires initiated by wildfires. That's an important distinction - and one that has

56
big repercussions for how we prepare ourselves for future fires," the authors wrote.

The authors included three researchers from the Forest Science and Fire Sciences laboratories of the U.S. Forest Service Rocky Mountain Research Station in Missoula and one from Headwaters Economics in Bozeman.

In a 2014 paper in the Proceedings of the National Academy of Science, some of the same authors developed a community risk assessment that put the focus on improving the security of individual homes in a community, not the forest around them. The emphasis is placed on modifying the house and the home ignition zone, a region within 100 feet of a house where debris and vegetation should be eliminated or minimized to reduce the chance of fire getting close to the house.

The reason that urban conflagrations begin and spread is because wind pushes embers and heat from one unprotected building to another, overwhelming fire departments that normally train to fight fire in just one building. Conditions are made worse when buildings are close together, because radiant heat becomes a bigger factor, spreading fire quicker.

"Reducing the likelihood that a home will ignite interrupts the disaster sequence by enabling effective structure protection. New construction siting, design, construction materials, and landscaping requirements should take wildfire potential into account," the authors wrote in the December paper.

57

One of the paper's authors, Jack Cohen, is a fire-behavior analyst and heat transfer engineer who has spent 40 years investigating wildfires, particularly those that are linked to incidents where hundreds of homes burned. He has spent at least the past decade writing papers and giving talks about the need to focus on making homes less susceptible to wildfires, which are a natural process, especially in the arid West.

When asked why the researchers decided to submit the recent article that seeks to drum home points they already promoted a decade ago, Cohen said cities and agencies have done very little during that time period to put their recommendations into place.

"What prompted us this time was the Lahaina urban conflagration that was associated with a grassfire. It may be a repeated message on our part, but it's not being received very well. Not much has changed," Cohen said. "The federal and

state agencies still don't get it - they're still defining the problem as a wildfire control problem."

Since the 2014 paper, Cohen and other researchers have had to just watch as town after town has burned terribly but predictably, as if no one has read their research. In Gatlinburg and Pigeon Forge, Tenn., 2,460 buildings burned in a 2016 fire; in 2018, the Camp Fire led to the loss of almost 19,000 buildings in Paradise, Calif.; in December 2021, 1,084 buildings burned in Superior and Louisville, Colo. from a grass fire; and in November 2021, a grassfire sparked fires in 23 homes in Denton, Mont.

58

Each wildfire had very little connection to most of the burning buildings, Cohen said. A wildfire is the source of initial ignition, but from that point on, it's a series of structure fires that lead to more structure fires. For example, with the Four Mile Canyon Fire in Boulder, Colo., the state of Colorado and the Forest Service had completed a number of fuel treatments nearby that they touted as protective. But high winds carried fire brands to ignite the houses far from the fire. Cohen found that while 168 houses burned, a lot of vegetation around the houses didn't, "so the wildfire didn't sweep through town."

"In the past five years, a number of incidents with more than 100 houses burning have been initiated by grass fires, which burn quickly. The grass fires pass through and are gone while the community continued to burn," Cohen said. "What I've found, particularly over the past five or six years, is that extreme wildfire is not dependent on closed-canopy conifers that produce big flames. The only time these urban disasters occur is under extreme conditions. That typically means it's very windy."

Nothing about the Lahaina Fire surprised Cohen. Not even the overblown claims that a wildfire "roared through and destroyed the town." Again, the wildfire was over before the town really started to burn. The fire started as a grassfire fanned by high winds, and had Lahaina not been there, the fire would have burned through the buffel grass and guinea grass within a matter of minutes before it died out on the beach.

But Lahaina was there, a high-density community with several blocks of multi-story, largely-connected wooden structures.

59

That configuration caused buildings to catch fire either due to burning embers flying from other buildings or from catching fire due to the overwhelming heat from nearby buildings.

"The ignition initiated where the grassfire came down, and that was it - it was a conflagration," Cohen said. "You don't want to be in a high-density community when you can't control the fire. Thirteen of the 26 fatalities in the 1991 Oakland Hills Fire occurred in the street when two-story buildings were burning on both sides of the street and the road became blocked. The heat was untenable."

One house in Lahaina stood untouched and was dubbed "the miracle house." But Cohen said it was just a good example of the points he and his fellow authors have been trying to communicate about defensible space and being fire-adapted. The owners had recently renovated the house with a nonflammable roof. It had wood walls, but the nearest building was about 30 feet away - far enough to prevent radiant heat from starting a fire - and there was little debris on the grounds or the house to actively spread the fire.

"The home ignition zone works," Cohen said. "The home ignition zone came out of the modeling I did and then the crown fire experiments I did with wood walls to show the distance, the proximity required to produce an ignition was realistic. At the same time, California was cutting 300-foot clearances around communities, which means nothing to (airborne) burning embers, but it's way over (what's required) for radiant heat exposure."

60

Cohen and his colleagues hope their latest paper prompts more action from local governments. Cohen is hoping Missoula County can do a better job when it updates its Wildfire Protection Plan in the near future.

But more than likely, Cohen said, they'll be writing a similar paper in another few years, trying to make politicians and the public understand. It doesn't help that they're fighting some in their own agency, the Forest Service, who insist that logging, not home modification, will save communities.

"Fire is inevitable. But nobody's figuring it out," Cohen said.

"We're starting from the presumption that it's wildfire that spreads through a community that lays it to waste. We even have the agencies responding in that fashion by being obsessed with this notion of wildfire control. So they do fuel treatments to have safe firefighting. That's not only counter ecologically, it doesn't work."

Contact reporter Laura Lundquist at lundquist@missoulacurrent.com.

Please find, "Wildland-urban fire disasters aren't actually a wildfire problem," by Calkin et al. 2023 attached. Calkin et al. 2024 found that to protect homes from wildfire we have to harden homes, not cut and burn forests.

The project area should be within 100 feet of homes not on Forest Service and BLM lands unless a home is within 100 feet of Forest Service and BLM lands. The purpose and need are not

61

based on the best available science and is in violation of NEPA, NFMA and the APA.

Please see the attached paper titled: "Management for Mountain Pine Beetle Outbreak Suppression: Does Relevant Science Support Current Policy?" By Six et al. 2014.

The abstract states: While the use of timber harvests is generally accepted as an effective approach to controlling bark beetles

during outbreaks, in reality there has been a dearth of monitoring to assess outcomes, and failures are often not reported. Additionally, few studies have focused on how these treatments affect forest structure and function over the long term, or our forests' ability to adapt to climate change. Despite this, there is a widespread belief in the policy arena that timber harvesting is an effective and necessary tool to address beetle infestations. That belief has led to numerous proposals for, and enactment of, significant changes in federal environmental laws to encourage more timber harvests for beetle control. In this review,

62

we use mountain pine beetle as an exemplar to critically evaluate the state of science behind the use of timber harvest treatments for bark beetle suppression during outbreaks. It is our hope that this review will stimulate research to fill important gaps and to help guide the development of policy and management firmly based in science, and thus, more likely to aid in forest conservation, reduce financial waste, and bolster public trust in public agency decision-making and practice. The paper finds:

Mountain pine beetle can remain in non-outbreak phase for very long periods of time, even when forests are composed of suitable age classes of host trees and in a condition often considered to be highly susceptible and "unhealthy". Outbreaks occur only when multiple thresholds involving temperature, tree defenses, and brood productivity are surpassed that allow positive feedbacks to amplify across several scales [2,64].

63

While outbreak development is complex, the primary elements that must exist are an abundance of suitable hosts and a trigger [63]. Triggers for mountain pine beetle that allow population amplification and subsequent widespread outbreak initiation are warm temperatures and drought, conditions that often co-occur [65]. There can also be a substantial lag period, even several years, from the initiation of the abiotic factors that trigger an outbreak to when populations actually amplify [65,66]. However, once a threshold number of beetles is surpassed, the outbreak becomes self-perpetuating. They also state: The on-the-ground reality is that direct control efforts typically fall far below the levels needed to stabilize, let alone control, mountain pine beetle populations. In the above cited studies, rates of detection in mitigated stands ranged from 45%-79%. These situations are not unusual. Direct control treatments are laborious, extremely costly and time consuming, and require high levels of training. Logistical difficulties,

64

including proper seasonal timing, access, inclement weather, and lack of trained personnel, increase the odds that they will not be effective. The high financial cost of such efforts coupled with a volatile market for sawtimber, pulp and pellets further complicates the use of direct controls. Importantly, outbreak

development is extremely swift and the amount of mitigation required can rapidly outstrip the ability of managers to respond.

Six et al. 2014 also note:

The hypothesis that light has a strong effect on mountain pine beetle behavior, particularly in reducing attacks, has led to a new treatment called daylighting. This approach is currently being implemented on a broad scale by federal and western state agencies. Daylighting involves removing trees and vegetation from around trees that are targeted for retention and is believed to work by repelling beetles from the boles of trees by increasing light and solar radiation [117]. While widely recommended, the efficacy of this treatment is unknown; there are no published studies on its effects on bark beetles.

65

Six et al. 2014 found that beetles are selective in killing the least healthy trees but logging occurs without consideration of genetics.

Very importantly, the beetle exercises selectivity in the trees it kills. While extremely high numbers may override this selectivity, evidence is accumulating that, even under outbreak conditions, beetles choose trees that have particular qualities. Beetles commonly select trees for attack that exhibit lower growth rates, defenses, and higher water stress [58,74,77]. While these factors can be influenced both locally and regionally by site conditions and climate, much of the variation in these properties within individual stands that affect bark beetle choice likely has a genetic basis. Outbreaks can result in strong natural selection against trees with phenotypes (and likely genotypes) favorable for the beetle and for those that possess unfavorable qualities [58,77]. However, when humans thin forests, trees are removed according to size, species, and density, without consideration of genetics. Thus, trees best adapted to surviving beetle outbreaks are as likely to be removed as those that are not.

When humans thin forests, they typically manage for resistance and resilience, rather than adaptation which involves genetic change. It is very important to distinguish between resistance, resilience, and adaptation, as each have different goals and operate on different temporal scales [140]. Resistance is a short-term holding action where we try to maintain an existing state. Approaches focusing on resistance often require massive interventions and increasing physical and financial investments over time. Such approaches may set

66

forests up for future outbreaks [136] and even catastrophic failure as they surpass thresholds in a warming climate [140]. In contrast, practices that promote resilience attempt to allow forests the ability to adjust to gradual changes related to climate change and to recover after disturbance. However, like resistance, resilience is not a long-term solution. In the long

term, forests must be able to adapt to change. Adaptation involves genetic change driven by natural selection. Currently, much of forest management, including bark beetle management, focuses on resistance and resilience, mainly through direct and indirect management, respectively. However, neither approach allows for true adaptation. For long term continuity of our forests, it will be imperative to begin to incorporate this aspect of management into our approaches.

Six et al 2014 conclude: One of the biggest problems in assessing the utility of direct controls is a general lack of monitoring or post hoc assessments of the outcomes of implementing these practices. Despite decades of direct control and large-scale implementation of these practices, few rigorous studies on its efficacy have been done and there remains no agreement among scientists or foresters regarding its ability to reduce beetle populations or losses of trees. Studies conducted prior to the current outbreak have variously concluded that direct treatments

67

may merely act to delay infestation of susceptible stands [97], or that if used correctly, can be effective [98,99]. Many studies found that while some treatments slowed the rate of infestation, overall, they had little to no impact on mountain pine beetle populations [97,100-104].

The manner in which policy makers have accepted beetle timber harvest treatments as a panacea for responding to bark beetle outbreaks in North American forests raises a number of red flags. As ecosystems and places that have economic, social, and cultural value to human communities are altered by climate change, there is a risk that people will overreact because of a need to "do something" to respond to change, and to give themselves some sense of control over broader forces that appear to be out of control. That pressure, to "do something", might also interact with the uncertainty about which choices are effective and appropriate (as with beetle timber harvest treatments) to create an opportunity for political pressures to force the adoption of particular choices that benefit specific interest groups [143]. It is perhaps no accident that the beetle treatments that have been most aggressively pushed for in the political landscape allow for logging activities that might provide revenue and jobs for the commercial timber industry. The result is that the push to "do something," uncertainty, and political pressures might lead us to act to respond to climate change before we understand the consequences of what we are doing, in the end producing more harm than good.

68

Our argument here is not to forgo management, but rather that management should be led by science and informed by monitoring. Both direct and indirect management for bark beetles have their place. However, to manage our forests in a way that best ensures their long-term function while wisely using limited financial resources, policy makers and the public

need a clearer understanding of current science
Please take a hard look at the impact of this project on climate
change. ?

Thank you for your attention to these concerns.

Sincerely yours,

Mike Garrity

Alliance for the Wild Rockies

P.O. Box 505

69

Helena, Montana 59624

And on behalf of:

Sara Johnson Native Ecosystems Council

P.O. Box 125?

Willow Creek, MT 59760

?

and for

Steve Kelly

Council on Wildlife and Fish

P.O. Box 4641 ?

Bozeman, MT 59772 ?

?

Tel: (406) 586-4421

70

And for

Kristine Akland

Center for Biological Diversity

P.O. Box 7274

Missoula, MT 59807

kakland@biologicaldiversity.org

71