

Older eastern white pine trees and stands sequester carbon for many decades and maximize cumulative carbon

Susan A. Masino^{1, 2}, William R. Moomaw^{3, 4*}, Robert T. Leverett⁵

¹Psychology / Neuroscience, Trinity College, United States, ²Harvard Forest, Faculty of Arts and Sciences, Harvard University, United States, ³Tufts University, United States, ⁴Woodwell Climate Research Center, United States, ⁵The Eastern Native Tree Society (ENTS), United States

Submitted to Journal:
Frontiers in Forests and Global Change

Specialty Section:
Tropical Forests

Article type:
Original Research Article

Manuscript ID:
620450

Received on:
22 Oct 2020

Revised on:
11 Feb 2021

Journal website link:
www.frontiersin.org

Conflict of interest statement

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest

Author contribution statement

RTL chose site locations and individual trees, established measurement methods and protocols, did the on-site tree measuring, and performed the subsequent analysis. SAM analyzed and organized the content and the supplements and participated in drafting and finalizing the text. WRM framed the analysis in the context of other studies and the larger context of climate change, assisted with data analysis and presentation, and drafting and editing the text.

Keywords

proforestation, Intact forest, Ecological resilience, Carbon accumulation, Chronosequence, Old-growth and second-growth forest, Tree volume, ecological integrity

Abstract

Word count: 343

Pre-settlement New England was heavily forested, with trees exceeding 2 m in diameter. The forests have regrown since farm abandonment, representing what is arguably the most successful regional reforestation on record and identified recently in the "Global Safety Net." Temperate "old-growth" forest and remnant stands demonstrate that native tree species can live several hundred years and continue to add to forest biomass and structural and ecological complexity. Forests globally are an essential natural climate solution that accumulate carbon and reduce annual increases in atmospheric CO₂ by approximately 30%. Some studies emphasize young, fast-growing trees and forests while others highlight carbon storage and accumulation in old trees and intact forests. We addressed this directly within New England with long-term, accurate field measurements and volume modeling of individual trees and two stands of eastern white pines (Pinaceae: *Pinus strobus*) and compared our results to models developed by the U.S. Forest Service. Within this sample and species, our major findings complement and clarify previous findings and are three-fold: 1) beyond 80 years, an intact eastern white pine forest can accumulate carbon above-ground in living trees at a high rate and double the carbon stored in this compartment in subsequent years; 2) large trees dominate above-ground carbon and can continue to accumulate carbon; 3) productive stands can continue to sequester high amounts of carbon in live trees for well over 150 years. Because the next decades are critical in addressing the climate crisis, and most New England forests are less than 100 years old, a major implication of this work is that maintaining and accumulating carbon in some existing forests – proforestation – is a powerful regional climate solution. Furthermore, older and old-growth trees and forests are rare, complex, highly dynamic and biodiverse: dedication of some forests to proforestation will produce large carbon-dense trees and also protect ecosystem integrity, special habitats, and native biodiversity long-term. In sum, strategic policies to grow and protect suitable existing forests in New England will optimize a proven, low cost, natural climate solution that also protects and restores biodiversity across the landscape.

Contribution to the field

Forests are an essential natural climate solution for accumulating and storing atmospheric CO₂, and some studies emphasize young, fast-growing trees and forests whereas others highlight high carbon storage and accumulation rates in old trees and intact forests. To address this question directly within New England we leveraged long-term, accurate field measurements along with volume modeling of individual trees and intact stands of eastern white pines (*Pinus strobus*) and compared our results to models developed by the U.S. Forest Service. Our major findings complement, extend, and clarify previous findings and are threefold: 1) intact forests continue to sequester carbon and store high cumulative carbon above ground; 2) large trees dominate above-ground carbon storage and continue sequestering carbon for hundreds of years; 3) pine stands continue to sequester high amounts of carbon for well over 150 years. Because the next decades are critical in addressing the climate crisis, and the vast majority of New England forests are less than 100 years old, and can at least double their cumulative carbon, a major implication of this work is that maintaining and accumulating maximal carbon in existing forests – proforestation – is a powerful near-term regional climate solution.

Funding statement

Ethics statements

Studies involving animal subjects

Generated Statement: No animal studies are presented in this manuscript.

Studies involving human subjects

Generated Statement: No human studies are presented in this manuscript.

Inclusion of identifiable human data

Generated Statement: No potentially identifiable human images or data is presented in this study.

Data availability statement

Generated Statement: The original contributions presented in the study are included in the article/supplementary material, further inquiries can be directed to the corresponding author/s.

In review

Older eastern white pine trees and stands sequester carbon for many decades and maximize cumulative carbon

Robert T. Leverett¹, Susan A. Masino², William R. Moomaw^{3,*}

¹American Forests National Champion Tree Program, Washington, DC, USA; Native Tree Society (www.nativetreesociety.org); Friends of Mohawk Trail State Forest, Florence, MA, USA

²Trinity College, Hartford, CT; Harvard Forest (2018-2019), Petersham, MA, USA

³The Fletcher School and Global Development and Environment Institute, Tufts University, Medford, MA, USA; Woodwell Climate Research Center, Falmouth MA USA

Correspondence:

William R. Moomaw, Ph.D.

William.moomaw@tufts.edu

Abstract

Pre-settlement New England was heavily forested, with trees exceeding 2 m in diameter. The forests have regrown since farm abandonment, representing what is arguably the most successful regional reforestation on record and identified recently in the “Global Safety Net.” Temperate “old-growth” forest and remnant stands demonstrate that native tree species can live several hundred years and continue to add to forest biomass and structural and ecological complexity. Forests globally are an essential natural climate solution that accumulate carbon and reduce annual increases in atmospheric CO₂ by approximately 30%. Some studies emphasize young, fast-growing trees and forests while others highlight carbon storage and accumulation in old trees and intact forests. We addressed this directly within New England with long-term, accurate field measurements and volume modeling of individual trees and two stands of eastern white pines (*Pinaceae: Pinus strobus*) and compared our results to models developed by the U.S. Forest Service. Within this sample and species, our major findings complement and clarify previous findings and are three-fold: 1) beyond 80 years, an intact eastern white pine forest can accumulate carbon above-ground in living trees at a high rate and double the carbon stored in this compartment in subsequent years; 2) large trees dominate above-ground carbon and can continue to accumulate carbon; 3) productive stands can continue to sequester high amounts of carbon in live trees for well over 150 years. Because the next decades are critical in addressing the climate crisis, and most New England forests are less than 100 years old, a major implication of this work is that maintaining and accumulating carbon in some existing forests – proforestation – is a powerful regional climate solution. Furthermore, older and old-growth trees and forests are rare, complex, highly dynamic and biodiverse: dedication of some forests to proforestation will produce large carbon-dense trees and also protect ecosystem integrity, special habitats, and native biodiversity long-term. In sum, strategic policies to grow and protect suitable existing forests in New England will optimize a proven, low cost, natural climate solution that also protects and restores biodiversity across the landscape.

Keywords: carbon accumulation, proforestation, chronosequence, tree volume measurements, old-growth forest, ecological integrity, ecological resilience

Introduction

A global priority for the climate has long been reducing ongoing emissions of heat-trapping greenhouse gases (GHGs) produced by burning carbon-based fuels. While this is essential, it is not sufficient for halting the rise in global temperatures. It is necessary to also simultaneously increase carbon dioxide (CO₂) removal (CDR) and keep carbon stored within natural systems. Clearing and harvesting forests, draining and developing wetlands, and degrading soils account for one-third of all the CO₂ added to the atmosphere by humans since the beginning of the industrial revolution (Simmons and Matthews, 2016). Together, these ongoing actions continue to add approximately 1.6 PgC/year (1 Pg equals 1 Gt or 10¹⁵ grams or 1 billion metric tonnes; Friedlingstein et al., 2020). Burning wood for heat and electricity adds additional CO₂, and current forest management practices limit the potential of this natural solution to accumulate carbon above and below ground and keep it out of the atmosphere (Sterman et al., 2018).

Two recent Intergovernmental Panel on Climate Change (IPCC) reports identify the urgent and unprecedented imperative to simultaneously and rapidly reduce Carbon Dioxide Emissions and achieve additional Carbon Dioxide Removal (CDR) from the atmosphere (IPCC, 2018; 2019). These reports identify forests as playing a major role in accumulating carbon out of the atmosphere. However, for CDR the focus is primarily on afforestation (planting new forests) and reforestation (regrowing forests) and ignores the more rapid climate mitigation and adaptation benefits of additional growth by existing forests, termed “proforestation” (Moomaw et al., 2019).

Even achieving the goal of “zero net carbon” will only “probably” limit global average temperatures to 1.5°C (IPCC, 2018) above the pre-industrial global temperature and a significant increase above the current level (1.1°C). This additional temperature increase will result in greater disruption to the climate system and will accelerate ecological decline. To avoid ever-more serious consequences of a changed climate, the goal must be to become net carbon *negative* as soon as possible. Growing suitable existing forests is an effective and low cost means for reducing the atmospheric stock of carbon as others have noted (Fargione et al., 2018; Hudiburg et al., 2019; Moomaw et al., 2019; Mildrexler et al., 2020) and will be demonstrated by the findings reported in this paper. Natural regeneration of forests has recently been found to accumulate more carbon in the first 30 years than managed reforestation (Cook-Patton et al., 2020).

A second and perhaps even more urgent priority is the strong protection of intact biodiverse natural systems (Watson et al., 2018), as verified in the Global Assessment Report on Biodiversity and Ecosystem Services (Intergovernmental Science-Policy on Biodiversity and Ecosystem Services, 2019) and the recent “Global Deal for Nature” (Dinerstein et al., 2019). A global review with a dual focus on carbon and biodiversity identified regions that are part of a “Global Safety Net” (Dinerstein et al., 2020), and the safety net must be now be translated to local levels. This joint climate/biodiversity priority was also highlighted in the peer-reviewed declaration of a Climate Emergency signed by over 13,000 scientists in late 2019 and which highlighted proforestation as a global climate solution (Ripple et al., 2020).

There is scientific consensus that we can substantially close the gap between CO₂ emissions and removals by maximizing a range of nature-based solutions (Griscom et al., 2017; Fargione et al., 2018). Regarding biodiversity, the beneficial role of protected areas in supporting species abundance and diversity was confirmed in a global meta-analysis (Coetzee et al., 2014), and the benefit of protecting intact ecosystems was quantified by comparing the probability of extinction in the six major global regions. On average, “wilderness” reduces the rate of species’ extinction by half due to higher

rates of species loss in unprotected areas (Di Marco et al., 2019); the quantified benefit of wilderness in preventing extinction is even higher in regions, including the Eastern United States. Biodiverse intact forests can simultaneously provide long-term protection to natural processes and biodiversity, reduce extinction, and provide pathways for migration while accumulating atmospheric carbon moderating local and global temperature increases (Friedlingstein et al., 2020). Taken together, it is practical and possible to act immediately to protect ecosystems and prevent extinction while we maintain increased CDR rates and store and accumulate additional carbon in forests and forest soils.

Forest conservation studies tend to focus on high-biodiversity tropical forests (Mitchard, 2018), yet temperate forests are also biodiverse (Hilmers et al., 2018), benefit human health and well-being in highly populated areas (Karjalainen et al., 2010), and provide many essential ecosystem services (United States Forest Service, 2021). They also have a large additional potential for CDR that has been underestimated by 32% (Cook-Patton et al., 2020). New England Acadian Forests are the only region in the lower 48 United States identified as part of the “Global Safety Net” as a Tier 1 climate stabilization area (Dinerstein et al., 2020). Current forest CDR in the United States reduces annual net nation-wide greenhouse gas emissions by 11.6% (United States Environmental Protection Agency, 2018), with the potential for much more (Keeton et al., 2011; Moomaw et al., 2019). Houghton and Nassikas (2018) estimate the current gross carbon sink in forests recovering from harvests and in abandoned agriculture to be -4.4 PgC/year (negative means removal) globally, consistent with the IPCC 1.5°C report that identified forests as key to increasing accumulation rates. This potential carbon sink from recovering forests is nearly as large as the gap between anthropogenic emissions and removal rates, 5.1 PgC/year (Friedlingstein et al., 2020).

In the context of resource production and forest management, some forest carbon is stored in lasting wood products, and responsible forestry can provide a reliable wood supply from a semi-natural forest. However, multiple analyses have found that more carbon associated with timber harvests is lost to the atmosphere than is stored in the harvested wood products (Nunery and Keeton, 2010; Harris et al., 2016). For example, just 19% of the original carbon stock in Oregon forests in 1900 is in long lived wood products; approximately 16% is in landfills, and the remaining 65% is in the atmosphere as carbon dioxide (Hudiburg et al., 2019). Updated models indicate that the product substitution benefits of wood products are overestimated between 2 and 100-fold (Harmon, 2019) and any near-term carbon benefit relies on product substitution (Hudiburg et al., 2019; Leturcq, 2020). Biogenic emissions from harvesting in the United States are estimated to be 640 MtC/year or 85% of total forestry emissions, exceeding the commercial and residential building sectors, and fossil fuel emissions from harvesting add an additional 17% CO₂ to the atmosphere above biogenic emissions (Harris et al., 2016).

Strategic planning for responsible resource production can both mitigate these emissions and ensure a protected network of intact natural areas. For example, the US Climate Alliance underestimates the importance of “net carbon accumulation” in forests (United States Climate Alliance, 2021). Forests do accumulate net carbon now, but carbon above and below ground is far below historic levels and far below its potential (Law et al., 2018; Hudiburg et al., 2019). A critical and explicit goal is to increase and optimize carbon accumulation by utilizing some forests for responsible resource production as needed and protecting other forests for climate protection, long-term full biodiversity, science, and human health and well-being.

At a global level, if deforestation were halted, and existing secondary forests allowed to continue growing, a network of these intact forests would protect the highest number of species from extinction (Di Marco et al., 2019; World Wildlife Federation, 2020) and it is estimated that they could sequester ~120 PgC in the 84 years between 2016 and 2100 (Houghton and Nassikas, 2018). This is equivalent to about 12 years of current global fossil fuel carbon emissions. These global numbers are conservative as outlined in recent analyses (Cook-Patton et al., 2020) and they do not factor in the enhanced

regional CDR potential and high cumulative carbon that can be achieved with proforestation in such carbon-dense temperate forests of the Pacific Northwest (Law et al., 2018) and New England (Nunery and Keeton, 2010; Keeton et al., 2011; Moomaw et al., 2019; Dinerstein et al., 2020).

Because these global and regional projections can be difficult to translate locally, particularly over time, we focused on a detailed analysis of individual trees and stands in New England. Historically, between 80% and 90% of the New England landscape was heavily forested, and early chroniclers describe pre-settlement forests with many large, mature trees reaching 1 to 1.5 m in diameter (Whitney, 1996). Fast-growing riparian species like sycamores and cottonwoods could reach or exceed 2 m. Today, New England trees of this size are mostly found as isolated individuals in open areas, parks, and old estates. Old-growth forests (primary forests) and remnants are currently less than 0.2% of northern New England's landscape, and less than 0.03% in Southern New England. Ongoing attempts to document their value and identify their locations is underway (Davis, 1996; Kershner and Leverett, 2004; Ruddat, 2020). Secondary forests in New England consist mostly of smaller, relatively young trees (on average less than 100 years old). The U.S. Forest Service estimates that fewer than 7% of the nation's forests exceed 100 years in age.

Our goal in this study was to measure carbon directly in individual trees and in an "average" versus an older stand of eastern white pine (*Pinaceae: Pinus strobus*) in New England. Most forest carbon studies focus on large geographical areas, and utilize "net" carbon data gathered from LIDAR (Light Detection And Ranging) and satellite technology, as well as statistical modeling based on the US Forest Service methods. Upon examining these options we note that carbon estimates from different tools and models can lead to disparate results at the level of individual trees – and these errors can be extrapolated to stands (Leverett et al., 2020). Therefore, we capitalized on the extensive tree-measuring protocols and experience of the Native Tree Society (NTS) to conduct highly accurate direct field measurements and measure volume precisely in younger vs. older trees growing in stands (Native Tree Society, 2021). We used direct measurements to evaluate volume-biomass models from multiple sources and developed a hybrid – termed FIA-COLE – to capitalize on the strengths of each model. We calculated the live above-ground carbon (in tonnes) in individual eastern white pines and individuals of other species in pine stands using conservative assumptions and direct measurements in pines up to 190 years old.

2. Materials and methods

This paper centers primarily on 1) individual eastern white pines (*Pinaceae: Pinus strobus*), 2) a representative older pine stand in Western Massachusetts, named the *Trees of Peace (TOP)*: located in Mohawk Trail State Forest, Charlemont, MA), and 3) a nearby younger pine stand (~750 ft center to center from the *TOP*). Both stands regenerated naturally from pasture and they share abiotic conditions such as a similar elevation, soil type (Hinkley loamy), temperature and precipitation. The younger stand is slightly downslope, and neither shows evidence of major recent disturbance. In 1989 the *TOP* lost 6 trees in a storm. Currently the *TOP* has 76 pines covering 0.6 to 0.7 ha.

While not discussed in detail herein, we have also collected and analyzed data from NTS measurements in 38 other sites with white pines in the Eastern United States. Since 1990, NTS has taken thousands of on-site direct measurements of individual trees in dozens of stands of eastern white pines (See examples Supplement 1). Measurements are published on the society's website (Native Tree Society, 2021) and comprehensive measurement protocols were adopted from those developed by NTS (Leverett et al., 2020) and incorporated into the American Forests Tree Measuring Guidelines Handbook (Leverett and Bertollette, 2014). A brief description of the measurement methods and models is provided in section 2.1, Supplement 2 and in Leverett et al. (Leverett et al., 2020). Here, in

all cases, the best mathematical processes were applied, *e.g.* the sine instead of the tangent height method and the best statistical models.

In the pine stands, a point-centered plot was established with a radius of 35.89 m, covering 0.403 hectares (subsequently referred to as 0.4 ha), with the goal of evaluating a standard acre (radius: 117.75 ft), and thus relevant to forestry conventions in the U.S. Within the *TOP*, 44 mature white pine stems were tallied along with 20 hardwoods and eastern hemlocks greater than 10 cm in diameter at breast height (DBH, 4' 5" of 1.37 m from the ground). The measured acre had 50 pines in July 1989 when six trees were lost in a wind event. The pines are ~160 years old; the hardwoods and hemlocks are estimated to be between 80 and 100 years old.

2.1 Height and diameter direct measurement methodology

We quantified the volume of the trunk and limbs of each tree from heights and diameters measured with laser-based hypsometers, monoculars with range-finding reticles, traditional diameter tapes, and calipers (described in detail in Leverett et al., 2020). Each instrument was calibrated and independently tested for accuracy over a wide range of distances and conditions (see Supplement 2 for an example). Absolute accuracies of the two main infrared lasers were verified as ± 2.5 cm for distance, surpassing the manufacturer's stated accuracy of ± 4.0 cm. The tilt sensors were accurate to $\pm 0.1^\circ$, meeting the manufacturer's stated accuracy. The combination of these distance and angle error ranges, along with the most accurate trigonometric methods noted above (sine vs. tangent method), gave us height accuracies to within 10 to 15 cm on the most distant targets being measured and approximately half that on the closest targets. We distinguished the rated and/or tested accuracy of a particular sensor of an instrument (such as an infrared laser or tilt sensor) from the results of a measurement that utilized multiple sensors.

Tree heights were measured directly for each pine with a visible top, using the sine method (Supplement 2) whenever possible rather than the traditional tangent method. Our preference for the sine method is supported by NTS, the US Forest Service (Bragg et al., 2011) and American Forests (Leverett and Bertolette, 2014). The more traditional tangent method often over/under-estimates heights by treating the sprig being measured (interpreted as the top), as if it were located vertically over the end of the baseline. The heights of 38 white pines in the *TOP* with visible tops were measured directly using the sine method.

2.2 Use of a form factor and FIA-COLE in determining pine volume

To compute trunk volume directly from the base to the absolute top of a tree, diameters at base and breast height were measured with conventional calibrated tapes according to the procedures established and published by NTS. Diameters aloft were measured with the combination of laser range-finders and high performance monoculars with range-finding reticles. A miniature surveying device, the LTI Trupoint 300, was also used. Its Class II, phase-based laser is rated at an accuracy of ± 1.0 mm to clear targets and its tilt sensor is accurate to ± 0.1 degrees. In the *TOP*, we computed the volume of each pine's trunk and limbs using diameter at breast height, full tree height, trunk form, and limb factors. (See Supplement 3 for a discussion on the development of the form factor and its importance in measuring volume, with comparisons to other methods of measurement).

Detailed measurements of 39 sample trees established an average form factor (see NTS measurements in Supplement 3, Table S3.2). The volume of each sample tree was determined by dividing the trunk into adjacent sections, with the length of each section guided by observed changes in trunk taper and/or visibility. Each section was modeled as the frustum of a regular geometric solid (neiloid, cone, paraboloid; see Supplement 3 and Leverett et al., 2020, for formulas). The form factor for each pine

was computed by adding its section volumes to obtain total trunk volume and then dividing the result by the product of the pine's height and breast-high cross-sectional area. This produced an average factor that would fit the pines growing in a stand. We applied the average form factor to all pines included in the *TOP* as one determination of trunk volume.

For comparison to our direct volume measurements, we applied a hybrid volume-biomass model to compute trunk volumes for pines in the *TOP*. This hybrid allowed us to make use of the extensive analysis of the US Forest Service Forest Inventory and Analysis (FIA) program and database (which determines volume and biomass through the use of allometric equations; United States Forest Service, 2020) as well as the Carbon On-Line Estimator (COLE; National Council for Air Stream Improvement, 2020). This hybrid was termed FIA-COLE. See Supplement 4 for a full explanation of the variables and equations for defining trunk volume. We finalized volumes for the pines in the *TOP* by averaging our direct measurements with those of FIA-COLE.

For the total volume of the above-ground portion of a pine, we derived a factor for limbs, branches, and twigs as a proportion of the trunk volume using the FIA-COLE model (Supplement 5). That model includes all the branching in what is defined as the "top" in a biomass calculation and the limb factor for large trees is typically an additional 15-16%. We ran the model for each of the individuals in the *TOP* and calculated the volume. This was converted to biomass (density) and then to carbon mass using a conservative carbon mass fractional factor of 48%.

2.3 Analysis of individual pine trees and a representative stand

In addition to the *TOP*, and older exemplary pines, we quantified above-ground carbon in younger trees and a representative stand. To determine an "average" pine at 50 years we defined two populations: (1) trees at 50 years that are still alive today, and (2) trees that were alive at 50 years but are missing today. This allowed us to compute an average trunk size for the missing trees and the associated carbon. We also measured white pines from young to older ages to estimate growth rates and volumes. The number of pines alive at 50 years but not alive today was determined from stand density data coming from both field counts and FIA (United States Forest Service, 2020).

We extensively studied an ~80-year-old stand of pines adjacent to the *TOP* (Supplement 6) growing on a terrace located just downslope from the *TOP* in an area fairly well protected from wind and with similar abiotic conditions and adequate soil depth. This age is more representative of the average stand of eastern white pine in New England (60-80 years; United States Forest Service, 2019). We also considered the range of pines of known ages from stands within the vicinity and elsewhere. Where we could, we examined ring growth and height patterns for individual pines during their early years on a variety of sites in different geographical locations. In some cases, we examined stumps and measured the average ring width. In other cases, we measured trees and counted limb whorls to get age estimates.

We measured the tallest pine in the *TOP* over a long time-span (referred to as Pine #58, its research tag number). Pine #58 has been measured carefully and regularly over a period of 28 years. In 1992 the tree was 47.24 m tall and 2.93 m in circumference. Since then, it has been climbed 4 times, tape-drop-measured, and volume-determined. Pine #58 continues to grow and has enabled us to quantify the changes in carbon accumulation in a dominant tree over decades. See Supplement 7 for a detailed measurement history of Pine #58.

Live tree above-ground volumes were converted to mass using standard wood density tables (United States Department of Agriculture, 2009). The air-dried density for white pine is 385.3 kg/m³ (0.3853 tonnes/m³). As noted above, we calculated the amount of carbon in each pine conservatively as 48% of total air-dried weight, whereby a cubic meter of white pine trunk or limbs holds 0.18494 tonnes of

carbon. (at least 50% is used more commonly; the percentage of carbon content in different species ranges from ~47% to 52+% and there is evidence that pine is at the upper range (Nicodemus and Williams, 2004). Note that the carbon in a cubic meter of wood varies depending on the species and is usually greater in hardwoods (United States Department of Agriculture, 2009).

3. Results

Our measurements indicate that individual eastern white pines can accumulate significant above-ground volume/carbon up to at least 190 years, that this volume/carbon accumulation in an individual tree can accelerate beyond 100 years, and that a stand of pines can double its above-ground live carbon between ~80 and 160 years.

3.1 Analysis of dominant individuals and averages for stand-grown pines

As Pine #58 is the tallest and the largest tree (volume) in the Trees of Peace (*TOP*), its performance over time was analyzed in great detail. It started growing as part of a more tightly packed stand, but presently has ample space. Its circumference at breast height is 3.30 m, its height is 53.71 m, and its crown spread is approximately 15.5 m. Over a period of 26 years, beginning in 1992, Pine #58 has grown in circumference at an average rate of 1.39 cm per year and grown in height 23.71 cm per year. For a chronosequence, we assumed that Pine #58 grew a lot when it was young – an average of up to 61 cm per year in its first 50 years. Its trunk and limb volume was 23.33 m³ at the end of the 2018 growing season (Supplement 7).

Figure 1 shows the increase in height, circumference and volume of Pine #58 within each 50-year interval up to 150 years and includes a photo of the tree. Its estimated age is ~160 years, and we used a chronosequence to determine previous epochs. For dominant pines in stands on good sites, ring widths for the first 50 years average ~0.6 cm and thus a 1.88 m circumference at 50 years. (Note that we measured one exceptional pine at 2.13 m in circumference.) Heights of stands at age 50 depend largely on site characteristics and expressed as site index (the average height of a stand at 50 years). The average index for white pine in Massachusetts is approximately 20 m (William Van Doren, Massachusetts Department of Conservation and Recreation, *pers. comm.*). For Pine #58 we calculated a much higher index to assume rapid early growth in the first 50 years. Based on these principles, the change in circumference and growth in height were greatest in the first 50 years, and decreased in the next two 50-year periods, confirming young pines “grow more rapidly” in terms of annual height and radial increases. However, volume growth, and thus carbon accumulation, continued to increase in the epochs studied here. This is primarily because volume increases linearly with height but increases as the square of the diameter (see Figure 1 and Supplement 8).

As noted, we assumed Pine #58 had optimal rapid growth in the first 50 years. Even so, our analysis supports the conclusion that the pine accumulated the majority of its current carbon *after age 50* and at an increased rate during subsequent epochs. Pine #58 now stores 4.33 tC above ground and continues to grow. For comparison, the carbon sequestered in the trunk of the highest volume 50-year-old pine that we encountered (2.13 m circumference, 34.75 m height, and 0.4346 form factor) is 1.16 tC. Therefore, even in the best-case scenario Pine #58 would have acquired only a quarter of its current carbon by age 50. Note that the same crown area occupied by multiple younger trees cannot achieve the carbon in this larger tree (Robert T. Leverett, *unpublished observations*).

Up to a point, the carbon advantage gained by the older trees accelerates with their increasing age and size, a finding that has been affirmed globally (Stephenson et al., 2014). Figure 2 documents the average volume in individual pines in the stands at ~80 and ~160 years as well as several additional large pines. MSF Pine #1, the largest pine in Monroe State Forest, western Massachusetts, has a trunk

volume of 35.9 m³ at approximately 190 years (6.62 tC; Figure 2). Assuming its early years accumulated 1.16 tC at 50 years, which is the fastest growing 50-year old pine we measured in all sampled locations, the large pine added 5.46 tC between 50 and 190 years, or 1.95 tC per 50-year cycle after year 50. This is at least 1.68 times the rate of growth for the first 50 years. This compares to a 1.6 ratio for Pine #58. In both cases more than 75% of the carbon they sequestered occurred *after* their first 50 years even when assuming the most optimal growth observed during the first 50 years.

3.2 Stand measurements at ~80 and ~160 years

Detailed measurements were taken in comparable pine stands at ~80 and ~160 years (*TOP*). As noted, the average tree in each stand is shown in Figure 2, and the distribution of tree sizes in the *TOP* is shown in Figure 3A. The largest pine in the *TOP* holds 4.33 tC and the smallest holds 0.53, an eight-fold difference. A comparison of the stand density and above ground carbon at ~80 vs. ~160 yr are shown in Figure 3B.

Complete data for 76 individual pines in the *TOP* (the 0.4 ha primary plot plus additional trees in the stand) is provided in Supplement 9. Similar data were collected from 0.4 ha of the ~80-year old stand (Supplement 6). This age is more representative of the average stand of eastern white pine in New England. Average values for both stands are summarized in Table 1. As shown in Figure 2, we found an average of 0.66 tC per tree compared to 1.95 tC per tree in the *TOP*, a near tripling of carbon in the average individual pine in the older stand. We found a robust size distribution among the pines in the older stand (Figure 3A), as well as a lower stand density (fewer stems), and a higher level of carbon in the *TOP* (Figure 3B). Pines predominated both plots, and non-pine species added ~10% to the total above ground carbon in the *TOP* (Figure 3B).

We emphasize that all of our calculations are based on a conservative value for the carbon mass fractional factor in the pines (48%) and only include above ground live tree-based carbon – they do not include more labile sources of additional carbon in needles, leaves and understory plants, or the accumulation of carbon in downed woody debris in older stands. Our measurements also do not include the large store of underground carbon (the root system is typically estimated as an additional 15-20% of the above-ground tree volume, and total soil organic carbon can be an additional 50% or more (Birdsey and Heath, 1995). Therefore, the total carbon is considerably higher. Nevertheless, the live trees in the older stand hold twice the carbon of the younger stand: the above-ground tree-based carbon measured directly in the primary acre in the 80 year old stand is 46.9 tC and the 160-year-old stand is 94.4 tC, translating to 117.2 and 236.0 tC per hectare, respectively. Approximately 10% of the tree-based carbon in the older stand is non-pine whereas non-pine live tree carbon in the younger stand is negligible (Table 1).

4. Discussion

We found that above-ground carbon stored in individual eastern white pines (*Pinaceae: Pinus strobus*) and stands can continue to increase well beyond 150 years. A chronosequence coupled with decades of direct measurements of a dominant stand-grown individual pine in Massachusetts demonstrate that height and circumference increase rapidly during the first 50-year epoch with smaller increases in 50-year epochs thereafter. In contrast, volume (and therefore carbon) shows the smallest increment in the first 50 years and the biggest in the 50-yr epoch between 100 and 150 years. This superior carbon sequestration in older trees is consistent with recent reports of recent rapid sequestration of older oak trees in Massachusetts (Finzi et al., 2020) and the outsized forest accumulation in large trees (Stephenson et al., 2014; Mildrexler et al., 2020). Here, the largest pine measured in Massachusetts (by volume) achieved 6.62 tC at 190 years old, and we found very large pines at ages ranging up to 350 years at dozens of sites in the Eastern United States.

378 Using direct measurement of above-ground carbon in different-aged pine stands, we found that live
379 tree carbon can continue to increase in a pine stand up to at least 160 years. We found twice as much
380 above-ground live tree carbon in a measured research acre within the older vs. the younger stand. The
381 live pines in the older stand also exhibited marked size diversity and the stand had a higher tree species
382 diversity.

383 The representative stands in this analysis approximate the average pine forest age in New England
384 (~80 years old) and a comparable stand approximately twice that age. To determine the biomass and
385 above ground carbon in living trees as a function of tree size and age, we have used a combination of
386 direct measurements and a hybrid FIA-COLE (Forest Inventory and Analysis - Carbon On-Line
387 Estimator) volume and biomass model to quantify individual trees and stands of eastern white pine.
388 We found that individual trees continue accumulating carbon well past 150 years, and more than 75%
389 of the carbon in pines up to 190 years is gained after the first 50 years. Despite a lower stand density
390 (fewer stems), total above-ground carbon is greatest in older stands and continues to increase past 150
391 years. The carbon per hectare quantified in these stands aligns with previous averages for the region
392 and previous regional estimates that New England forests can accumulate between 2.3 and 4.2 times as
393 much carbon as they now contain on productive sites (Keeton et al., 2011). The total carbon stored is
394 much greater when below-ground carbon in roots, coarse woody debris, standing dead trees, smaller
395 plants and soils are included (Birdsey and Heath, 1995; Nunery and Keeton, 2010; Tomasso and
396 Leighton, 2014).

398 Forest managers stress the high accumulation rates of younger forests as important in absorbing
399 atmospheric CO₂. This is an important consideration for production forests as well as to help optimize
400 between growing a wood resource and accumulating carbon. Younger individual trees do not sequester
401 absolute amounts of carbon more rapidly than larger more mature trees, and we did not find evidence
402 for a significant benefit for a young stand compared to an older stand. We note this is a limited sample,
403 and we did not estimate rates of accumulation below 80 years (Table 1).

405 Multi-use forests provide a source of wood products and can support recreation but active management
406 practices limit forest carbon accumulation long-term. At a range of scales, chronic intervention
407 eliminates the ability for that forest to host the full biodiversity of some of our rarest species of plants,
408 animals, insects, fungi, lichens, reptiles and amphibians found in older and continuously forested areas
409 (McMullin and Wiersma, 2019; Moose et al., 2019) as well as climate-sensitive birds that may benefit
410 from mature or old-growth forests (Betts et al., 2017). These older unmanaged forests also have fewer
411 invasive species (Riitters et al., 2018).

413 The pine stands studied here grew from abandoned sheep pasture, and therefore were unlikely to have
414 been severely disturbed prior to natural regeneration. Site history influences growth and net carbon
415 accumulation, especially in the early years, since disturbed soil can continue to lose carbon for more
416 than a decade (Birdsey and Heath, 1995; Hamburg et al., 2019). We recognize that at some point the
417 above-ground carbon in living trees will no longer increase as the live trees in the stand eventually will
418 reach a steady state of death and renewal. Pines easily reach 200 years and some live 400 years; today
419 the *TOP* is less than halfway to that age and the younger stand is only ~25% of that lifespan. Previous
420 work shows that pine stands continue to add above ground carbon beyond 200 years (Seymour, 2011;
421 2016), and even when above-ground live carbon reaches asymptote, total forest carbon continues to
422 increase, even in some primary ("old-growth") forests (Mackey et al., 2015); after tree death or forest
423 disturbance there is a new growth as well as transfer of live carbon to dead wood and woody debris,
424 the litter layer, and into the soil. For example, 70 years after an old-growth (virgin) eastern hemlock
425 (*Tsuga canadensis*) and eastern white pine stand blew down (the 1938 Hurricane in New England) that
426 forest stored as much carbon as forests that were 250 years old (D'Amato et al., 2017).

There is no evidence of recent disturbance in either research plot herein. The older pine stand shows an increased prevalence and growth of trees of other species (including more carbon-dense hardwoods), and for multiple reasons it is unlikely it has reached maximal above-ground live carbon or total carbon. Rather, this forest appears to be transitioning into a phase where the structural diversity, species diversity and total carbon load will continue to rise. A goal for future research is a better understanding of tree and stand-level carbon accumulation and dynamics as well as many other ecological features in different forest types and in stands well beyond 150 years – a time when old-growth characteristics are starting to redevelop in eastern forests.

Public forests in New England are typically older than private forests (but still predominantly less than 100 years old), and provide the greatest possibility for future carbon-dense and biodiverse intact forests across the landscape. Native tree species can live for several hundred years (and in the case of eastern hemlock (*Tsuga canadensis*) and black gum (*Nyssa sylvatica*), up to and exceeding 500 years) (Whitney, 1996; Speduto et al., 2000). Despite the shortage of old and old-growth forests (and their proven resilience to disturbance (D'Amato et al., 2017), and the increased prevalence of natural disturbances (e.g. insect outbreaks, windstorms) creating forest diversity and forest openings (Oswalt 2019), a major focus across public land has been to make forests younger. These programs assert that these habitats prevent a suite of species from declining, that they sequester carbon more rapidly, and that they are more resistant to disturbance than their older counterparts (Anwar, 2001). This approach downplays the rate of the natural development of niches for multiple species (Zlonis and Niemi, 2014) and the accumulation of biodiversity in temperate forests during natural forest succession (Hilmers et al., 2018). It also overlooks cumulative forest carbon (Moomaw et al., 2019) as well as the superior resilience of older forests to the stresses of climate change (Thom et al., 2019). Comparing details of age and location (tropical, temperate, boreal, etc.) are important, as is evaluating the term “young” – in some cases it is considered up to 140 years (Pugh et al., 2019).

Our findings are consistent with Stephenson et al. (2014) who found that absolute growth increases with tree size for most of 403 tropical and temperate tree species, and a study of 48 forest plots found that in older forests, regardless of geographical location, half of all above-ground biomass (and hence carbon), is stored in the largest 1% of trees as measured by diameter at breast height (Lutz et al., 2018). An increase in carbon density per hectare was found as the age of the stand increased in the Northeast U.S. (Keeton et al., 2011), and a recent study in China found that forests with older trees and greater species richness had twice the levels of carbon storage than did less diverse forests with younger trees (Liu et al., 2018). Earlier work demonstrated that intact old growth forests in the Pacific Northwest contained more than twice the amount of sequestered carbon as did those that were harvested on a fixed rotation basis (Harmon et al., 1990).

Globally, forests are capable of accumulating twice as much atmospheric carbon, and the current deficit is due to a combination of conversion and management (Erb et al., 2018). Continuing current management in the Northeast will result in a large difference between the potential for land-based carbon and the current trajectory (Duveneck and Thompson, 2019). Meanwhile, natural regeneration and reforestation is a superior climate solution compared to managed reforestation and tree planting (Cook-Patton et al., 2020), and proforestation – growing existing natural forests – complements and extends natural regeneration as an ongoing climate solution by leveraging the accumulation potential in forests that are already established (Moomaw et al., 2019). Proforestation recognizes implicitly that older forests and large trees are critical to a global strategy for carbon accumulation and biodiversity protection (Lindenmayer and Laurance, 2016). Rapidly moving large stocks of atmospheric carbon as CO₂ into forests and reducing emissions are both essential to limiting the increase in global temperatures, and protecting intact and connected habitat is essential in preventing extinction. These time-sensitive dual goals and the importance of traditional indigenous land use are explicitly recognized internationally in the Global Deal for Nature, the Global Safety Net, and the recent

“Campaign for Nature” or “30 x 30” – i.e., protecting 30% of the planet’s land and water by 2030 (Campaign for Nature, 2021), and in the ambitious coalition goal of “Nature Needs Half” (Nature Needs).

An important additional implication of our study is that the estimated potential additional carbon dioxide (CO₂) removal (CDR) achieved by future growth of secondary forests as reported by Houghton and Nassikas (2018) is likely an underestimate because it does not account for high ongoing accumulation rates as trees age in regions with relatively young (compared to tree lifespan) forests like those of the Northeast United States. The global study of natural forest carbon accumulation by Cook-Patton et al. (2020) and the synthesis of quantified carbon and biodiversity by Moomaw et al. (2019) provide evidence for the power of natural forest processes throughout their growth and development. These reports and the current site-specific findings support the high regional contribution of carbon accumulation in the coming decades by Northeastern temperate forests and their designation as a Tier 1 climate stabilization region (Dinerstein et al., 2020).

While the IPCC clearly identified forests as essential for sequestering additional carbon for climate stability, it focused on production forests that are currently recovering from being harvested or on unforested areas where forests could be planted (afforestation). Bastin et al. (2019) proposes an afforestation project on 0.9 billion ha but acknowledges the relatively long time before large amounts of carbon would be sequestered. Global tree planting efforts are under way, but are presented too simplistically (Holl and Brancalion, 2020); for example, there is little data on how to plant an ecosystem, and tree planting efforts can suffer from numerous challenges, including high mortality (Cao et al., 2011). In contrast, growing existing forests is an established near-term strategy (Moomaw et al., 2019). Overall, afforestation and reforestation are valuable, but neither can keep as much carbon out of the atmosphere as proforestation in the next 50 years – the timeline when it is needed most to avoid irreversible consequences of a changed climate.

Although this study focused exclusively on above-ground live tree carbon accumulation, we emphasize that additional carbon exists and accumulates above and below ground. Other ecosystem services of proforestation also accrue, and the essential goal of protecting a “Global Safety Net” of nature extends explicitly beyond greenhouse gas emissions and mitigating the climate crisis (Dinerstein et al., 2020). Nevertheless, an accurate carbon-centric model of “business as usual” vs. proforestation must include comprehensive real-world carbon fluxes. Removing carbon from the forest releases carbon into the atmosphere, and in some cases a portion of the carbon is stored in wood and/or wood is substituting for other materials. Recent work shows that near-term carbon benefits associated with wood products and substitution have been overestimated based on outdated assumptions or neglecting or underestimating future accumulation (Harmon, 2019; Leturcq, 2020). Efforts should be made on consumption and conservation to ensure we protect primary forests and additional secondary forests where possible: carbon storage in forests is low-risk, high-capacity and practical – therefore preferable to experimental bioenergy with carbon capture and storage (BECCS) suggested by the IPCC report (Anderson and Peters, 2016; IPCC, 2018). Finally, letting existing secondary forests grow creates a network of nature that can provide equity, access to natural heritage, scientific discovery, and cumulative health benefits for people. Protecting and growing a network of suitable existing forests as a carbon sink in New England is cost-effective (Tomasso and Leighton, 2014) and does not compete directly with agriculture and other demands for land use.

The direct measurements at the tree and stand level in this paper are consistent with parameterized and other studies at larger scale in verifying that larger trees (Stephenson et al., 2014; Lutz et al., 2018) and stands of larger trees accumulate the most carbon over time compared to smaller trees (Mildrexler et al., 2020). They support the proforestation strategy of growing existing forests to achieve their natural capacity to accumulate carbon and achieve their ecological potential (Moomaw et al., 2019) to redress

the balance of carbon lost to the atmosphere from global forests due to human activity (Hudiburg et al., 2019). The important implication of these findings is that the trees and the forests that we need most for carbon storage and CDR to help limit near-term climate change are the ones that are already established.

Currently, plantations and forests managed for forest products account for 71% of all forest area globally (IPCC, 2019), more than sufficient for resource production. Strategic decisions can enable some of these forests to be dedicated to climate protection and research, and the remaining 29% should be protected wherever possible. This would be a major step toward the goal of “30 x 30” – with additional climate stabilization areas are needed beyond that. Together 30 x 30 plus climate stabilization will move us toward long-term protection of “half-earth” (Wilson, 2016). High levels of carbon accumulation and biodiversity protection are integral to resiliency in a changing climate – including the resiliency achieved by protecting species networks and interactions, genetic diversity and the potential for specific adaptive epigenetic changes (Hanlon et al., 2019). These complexities are poorly understood – science and technology is evolving, and new techniques can discover new species (Schulz et al., 2018) – and any areas, even on public land, lack a detailed ecological inventory due to resource constraints or a focus on other priorities. Meanwhile, intensive biodiversity inventories have yielded many hundreds of new species – often small species such microbes, lichen, fungi, algae and insects; *i.e.* Smokies Species Tally (Discover Life in America, 2021). Much more research is needed, and essential ecological processes develop and diversify at timescales far beyond a human lifetime.

In sum, the current findings ground-truth the capacity for a representative New England eastern white pine stand to at least double its above-ground live tree carbon in the coming decades, confirming previous chronosequencing of pine stands in the region (Seymour, 2011). We did not attempt to quantify or estimate the flux in other carbon compartments above or below ground. With a small fraction of New England (~3% overall, ~1% in Southern New England) prioritized for proforestation and natural processes, protection of a suitable network of land from unneeded intervention is urgent, and public land is the most logical place to start: funding to ensure evidence-based intervention and additional data collection will generate policies that protect the long-term public trust. At the same time, systems to support local wood use and reuse are equally needed to ensure the highest and best use of this resource, protect local expertise and jobs, and reduce emissions associated with the forest industry; in some states it is the largest source of emissions (Law et al., 2018). Comprehensive education, information and compensation programs should be established to provide private landowners a range of options based on numerous ecosystem services, including maximal carbon and biodiversity accumulation, with the goal of optimizing natural solutions that address the Climate Emergency immediately (Ripple et al., 2020). Failing to protect natural systems erodes the wealth and well-being that is essential to meet this unprecedented challenge and avoid “a ghastly future” (Bradshaw et al., 2021).

Acknowledgments

We acknowledge the critical technical assistance of Jared D. Lockwood in measuring pines in the *TOP*, the younger pine stand, and elsewhere as needed. We thank Ray Asselin for photographing pines for further analysis, and for ring and whorl counts to establish ages. We thank Monica Jakuc Leverett for invaluable assistance in editing and revising the original draft, thank Dr. David Ruskin for tireless efforts throughout the process, and thank Dr. Edward Faison for constructive suggestions. We also thank the reviewers for constructive comments. Supported by Trinity College (SAM), a Charles Bullard Fellowship in Forest Research (SAM), a Faculty Research Grant from the NASA Connecticut Space Grant Consortium (RTL, SAM) and the Rockefeller Brothers Fund (WRM).

580 **Author contributions statement**

581

582 RTL chose site locations and individual trees, established measurement methods and protocols, did the
583 on-site tree measuring, and performed the subsequent analysis. SAM analyzed and organized the
584 content and supplements and participated in drafting and finalizing the text. WRM framed the analysis
585 in the context of other studies and the larger context of climate change, assisted with data analysis and
586 presentation, and drafting and editing the text.

587

588 **Conflict of interest statement**

589

590 This work was not carried out in the presence of any personal, professional or financial relationships
591 that could potentially be construed as a conflict of interest.

592

In review

593 **Table 1.** Summary of key measurements within a 160-year pine stand (*TOP*) and a comparable ~80
 594 year old stand (2018 – 2019 values).
 595

Individual Values	<u>~160 year old 0.4 hectare</u>	
	Circumference at breast height (avg)	2.36 m
	Diameter at breast height (avg)	0.75 m
	Height (avg)	45.10 m
	Tree Volume (trunk + limbs; avg)	10.47 m ³
	Above-ground carbon per tree (avg)	1.95 tC
	<u>~80 year old 0.4 hectare</u>	
	Circumference at breast height (avg)	1.56 m
	Diameter at breast height (avg)	0.50 m
	Height (avg)	38.4 m
	Tree volume (trunk + limbs; avg)	3.58 m ³
	Above-ground carbon per tree (avg)	0.66 tC
Stand Values	<u>Full Stand at ~160 years</u>	
	Number of pines	76
	Above-ground pine-based carbon	146.84 tC
	Above-ground non-pine carbon	14.90 tC
	Total above-ground tree carbon	161.74 tC
	<u>Research Acre ~160 years (0.4 hectare)</u>	
	Number of pines	44
	Above-ground pine-based carbon	85.8 tC
	Above-ground non-pine carbon	8.6 tC
	Total above-ground tree carbon	94.4 tC
	<u>Research Acre ~80 years (0.4 hectare)</u>	
	Number of pines	71
	Total above-ground pine-based carbon (negligible non-pine carbon)	46.86 tC

596

Figure Legends

Figure 1. Changes in circumference, height and volume of a stand-grown individual eastern white pine (Pine #58) in three 50-y intervals. *Upper panels* – **A:** Change in circumference during 0-50, 50-100 and 100-150 y. **B:** Change in height between 0-50, 50-100 and 100-150 y. **C:** Change in above-ground tree volume (trunk plus limbs) between 0-50, 50-100 and 100-150 y. *Lower panels* – **D:** Cumulative circumference at 50, 100 and 150 y compared to cumulative above-ground volume. **E.** Cumulative height at 50, 100 and 150 y compared to cumulative above-ground volume. On each lower panel initial slopes were matched to reflect the rapid change in circumference and height during the first 50-y interval. Note that volume is a proxy for above-ground carbon. Values for circumference, height and volume of Pine #58 were determined by a combination of direct measurement and chronosequence and described in the text and in Supplement. **F.** Pine #58 (center) being readied for climbing and measuring.

Figure 2. Tonnes of above-ground carbon (tC) in an “average” eastern white pine in a measured research acre (green locants) and in five individual trees (A - E) measured directly on site at three separate locations in Massachusetts. Average tC and standard deviation is based on pines in a stand at ~80 years (0.66 ± 0.38 tC) and ~160 years (1.95 ± 0.73 tC) as described in the text. Direct measurement of tC is shown for individual trees in western Massachusetts at these ages and locations: A, B - ~190 years (MSF #1 and #2, Monroe State Forest); C - ~160 years (Pine #58, Mohawk Trail State Forest; more details of Pine #58 shown in Figure 1); D - ~150 years (Totem, Northampton, MA); E - ~120 years (BB #2, Broad Brook, Florence, MA).

Figure 3. Carbon distribution, stand density and cumulative carbon in predominantly eastern white pine stands at ~80 and 160 years. These two stands were regrown from land previously used as pasture (i.e. not recovering from a harvest at time zero). **A.** Distribution of above-ground carbon (tC) among 76 eastern white pines of different sizes in the full *TOP* stand at ~160 years old. The majority contained 1-3 tC. **B.** Stand density and above-ground carbon measured directly on site in a research acre of eastern white pine at ~80 and 160 years. Stand density (# of stems) declined while above-ground carbon increased. The older stand includes some non-pine species that added to the number of stems and total carbon (open locants).

References

- Anderson, K., and Peters, G. (2016). The trouble with negative emissions. *Science* 354, 182-183.
- Anwar, A. (2001). *Does the age of a tree affect carbon storage?* [Online]. GISS Institute on Climate and Planets, National Aeronautics and Space Administration. Available: <https://icp.giss.nasa.gov/research/ppa/2001/anwar/>.
- Bastin, J.F., Finegold, Y., Garcia, C., Mollicone, D., Rezende, M., Routh, D., Zohner, C.M., and Crowther, T.W. (2019). The global tree restoration potential. *Science* 365, 76-79.
- Betts, M.G., Phalan, B., Frey, S.J.K., Rousseau, J.S., and Yang, Z. (2017). Old-growth forests buffer climate-sensitive bird populations from warming. *Divers. Distrib.* 24, 439-447.
- Birdsey, R.A., and Heath, L.S. (1995). "Carbon changes in U.S. forests," in *Productivity of America's Forests and Climate Change. Gen. Tech. Rep. RM-GTR-271*, ed. L.A. Joyce. (Fort Collins, CO: U.S.D.A. Forest Service, Rocky Mountain Forest and Experiment Station: 56-70).
- Bradshaw, C.J.A., Ehrlich, P.R., Beattie, A., Ceballos, G., Crist, E., Diamond, J., Dirzo, R., Ehrlich, A.H., Harte, J., Harte, M.E., Pyke, G., Raven, P.H., Ripple, W.J., Saltr , F., Turnbull, C., Wackernagel, M., and Blumstein, D.T. (2021). Underestimating the challenges of avoiding a ghastly future. *Front. Conserv. Sci.* 1, 615419.
- Bragg, D.C., Frelich, L.E., Leverett, R.T., Blozan, W., and Luthringer, D.J. (2011). "The sine method: An alternative height measurement technique", F.S. U.S. Department of Agriculture, Southern Research Station. (Asheville, North Carolina: Res. note SRS-22).
- Campaign for Nature (2021). Available: <https://www.campaignfornature.org/>.
- Cao, S., Chen, L., Shankman, D., Wang, C., Wang, X., and Zhang, H. (2011). Excessive reliance on afforestation in China's arid and semi-arid regions: Lessons in ecological restoration. *Earth-Sci. Rev.* 104, 240-245.
- Coetzee, B.W., Gaston, K.J., and Chown, S.L. (2014). Local scale comparisons of biodiversity as a test for global protected area ecological performance: a meta-analysis. *PLoS One* 9, e105824.
- Cook-Patton, S.C., Leavitt, S.M., Gibbs, D., Harris, N.L., Lister, K., Anderson-Teixeira, K.J., Briggs, R.D., Chazdon, R.L., Crowther, T.W., Ellis, P.W., Griscom, H.P., Herrmann, V., Holl, K.D., Houghton, R.A., Larrosa, C., Lomax, G., Lucas, R., Madsen, P., Malhi, Y., Paquette, A., Parker, J.D., Paul, K., Routh, D., Roxburgh, S., Saatchi, S., Van Den Hoogen, J., Walker, W.S., Wheeler, C.E., Wood, S.A., Xu, L., and Griscom, B.W. (2020). Mapping carbon accumulation potential from global natural forest regrowth. *Nature* 585, 545-550.
- D'Amato, A.W., Orwig, D.A., Foster, D.R., Barker Plotkin, A., Schoonmaker, P.K., and Wagner, M.R. (2017). Long-term structural and biomass dynamics of virgin *Tsuga canadensis*-*Pinus strobus* forests after hurricane disturbance. *Ecology* 98, 721-733.
- Davis, M.B. (1996). *Eastern Old-Growth Forests: Prospects For Rediscovery And Recovery*. Washington DC: Island Press.
- Di Marco, M., Ferrier, S., Harwood, T.D., Hoskins, A.J., and Watson, J.E.M. (2019). Wilderness areas halve the extinction risk of terrestrial biodiversity. *Nature* 573, 582-585.
- Dinerstein, E., Joshi, A.R., Vynne, C., Lee, A.T.L., Pharand-Deschenes, F., Franca, M., Fernando, S., Birch, T., Burkart, K., Asner, G.P., and Olson, D. (2020). A "Global Safety Net" to reverse biodiversity loss and stabilize Earth's climate. *Sci. Adv.* 6, eabb2824.
- Dinerstein, E., Vynne, C., Sala, E., Joshi, A.R., Fernando, S., Lovejoy, T.E., Mayorga, J., Olson, D., Asner, G.P., Baillie, J.E.M., Burgess, N.D., Burkart, K., Noss, R.F., Zhang, Y.P., Baccini, A., Birch, T., Hahn, N., Joppa, L.N., and Wikramanayake, E. (2019). A Global Deal For Nature: Guiding principles, milestones, and targets. *Sci. Adv.* 5, eaaw2869.
- Discover Life in America (2021). Available: <https://dlia.org/smokies-species-tally>.
- Duveneck, M.J., and Thompson, J.R. (2019). Social and biophysical determinants of future forest conditions in New England: Effects of a modern land-use regime. *Glob. Environ. Change* 55, 115-129.

679 Erb, K.-H., Kastner, T., Plutzer, C., Bais, A.L.S., Carvalhais, N., Fetzel, T., Gingrich, S., Haberl, H.,
680 Lauk, C., Niedertscheider, M., Pongratz, J., Thurner, M., and Luyssaert, S. (2018).
681 Unexpectedly large impact of forest management and grazing on global vegetation biomass.
682 *Nature* 553, 73-76.

683 Fargione, J.E., Bassett, S., Boucher, T., Bridgman, S.D., Conant, R.T., Cook-Patton, S.C., Ellis, P.W.,
684 Falcucci, A., Fourqurean, J.W., Gopalakrishna, T., Gu, H., Henderson, B., Hurteau, M.D.,
685 Kroeger, K.D., Kroeger, T., Lark, T.J., Leavitt, S.M., Lomax, G., McDonald, R.I., Megonigal,
686 J.P., Miteva, D.A., Richardson, C.J., Sanderman, J., Shoch, D., Spawn, S.A., Veldman, J.W.,
687 Williams, C.A., Woodbury, P.B., Zganjar, C., Baranski, M., Elias, P., Houghton, R.A., Landis,
688 E., McGlynn, E., Schlesinger, W.H., Siikamäki, J.V., Sutton-Grier, A.E., and Griscom, B.W.
689 (2018). Natural climate solutions for the United States. *Sci. Adv.* 4, eaat1869.

690 Finzi, A.C., Giasson, M.-A., Barker Plotkin, A.A., Aber, J.D., Boose, E.R., Davidson, E.A., Dietze,
691 M.C., Ellison, A.M., Frey, S.D., Goldman, E., and Al, E. (2020). Carbon budget of the Harvard
692 Forest Long-Term Ecological Research site: pattern, process, and response to global change.
693 *Ecol. Monogr.* 90, e01423.

694 Friedlingstein, P., O'Sullivan, M., Jones, M.W., Andrew, R.M., Hauck, J., Olsen, A., Peters, G.P.,
695 Peters, W., Pongratz, J., Sitch, S., Le Quéré, C., et al. (2020). Global carbon budget 2020.
696 *Earth Syst. Sci. Data* 12, 3269-3340.

697 Griscom, B.W., Adams, J., Ellis, P.W., Houghton, R.A., Lomax, G., Miteva, D.A., Schlesinger, W.H.,
698 Shoch, D., Siikamäki, J.V., Smith, P., Woodbury, P., Zganjar, C., Blackman, A., Campari, J.,
699 Conant, R.T., Delgado, C., Elias, P., Gopalakrishna, T., Hamsik, M.R., Herrero, M., Kiesecker,
700 J., Landis, E., Laestadius, L., Leavitt, S.M., Minnemeyer, S., Polasky, S., Potapov, P., Putz,
701 F.E., Sanderman, J., Silvius, M., Wollenberg, E., and Fargione, J. (2017). Natural climate
702 solutions. *Proc. Natl. Acad. Sci. USA* 114, 11645-11650.

703 Hamburg, S.P., Vadeboncoeur, M.A., Johnson, C.E., and Sanderman, J. (2019). Losses of mineral soil
704 carbon largely offset biomass accumulation 15 years after whole-tree harvest in a northern
705 hardwood forest. *Biogeochemistry* 144, 1-14.

706 Hanlon, V.C.T., Otto, S.P., and Aitken, S.N. (2019). Somatic mutations substantially increase the per-
707 generation mutation rate in the conifer *Picea sitchensis*. *Evol. Lett.* 3, 348-358.

708 Harmon, M.E. (2019). Have product substitution carbon benefits been overestimated? A sensitivity
709 analysis of key assumptions. *Environ. Res. Lett.* 14, 065008.

710 Harmon, M.E., Ferrell, W.K., and Franklin, J.F. (1990). Effects on carbon storage of conversion of
711 old-growth forests to young forests. *Science* 247, 699-702.

712 Harris, N.L., Hagen, S.C., Saatchi, S.S., Pearson, T.R.H., Woodall, C.W., Domke, G.M., Braswell,
713 B.H., Walters, B.F., Brown, S., Salas, W., Fore, A., and Yu, Y. (2016). Attribution of net
714 carbon change by disturbance type across forest lands of the conterminous United States.
715 *Carbon Balance Manag.* 11, 24.

716 Hilmers, T., Friess, N., Bassler, C., Heurich, M., Brandi, R., Pretzsch, H., Seidl, R., and Muller, J.
717 (2018). Biodiversity along temperate forest succession. *J. Appl. Ecol.* 55, 2756-2766.

718 Holl, K.D., and Brancalion, P.H.S. (2020). Tree planting is not a simple solution. *Science* 368, 580-
719 581.

720 Houghton, R.A., and Nassikas, A.A. (2018). Negative emissions from stopping deforestation and forest
721 degradation, globally. *Glob. Chang. Biol.* 24, 350-359.

722 Hudiburg, T.W., Law, B.E., Moomaw, W.R., Harmon, M.E., and Stenzel, J.E. (2019). Meeting GHG
723 reduction targets requires accounting of all forest emission sectors. *Environ. Res. Lett.* 14,
724 095005.

725 Intergovernmental Panel on Climate Change (2018). "Summary for Policymakers," in *Global warming*
726 *of 1.5°C. An IPCC Special Report on the impacts of global warming of 1.5°C above pre-*
727 *industrial levels and related global greenhouse gas emission pathways, in the context of*
728 *strengthening the global response to the threat of climate change, sustainable development,*
729 *and efforts to eradicate poverty*, eds. V. Masson-Delmotte, P. Zhai, H. O. Pörtner, D. Roberts,

730 J. Skea, P. R. Shukla, A. Pirani, W. Moufouma-Okia, C. Péan, R. Pidcock, S. Connors, J. B. R.
731 Matthews, Y. Chen, X. Zhou, M. I. Gomis, E. Lonnoy, T. Maycock, M. Tignor & T. Waterfield
732 (Geneva, Switzerland: World Meteorological Society).

733 Intergovernmental Panel on Climate Change (2019). "Summary for Policymakers," in *Global Warming*
734 *of 1.5°C. An IPCC Special Report on the impacts of global warming of 1.5°C above pre-*
735 *industrial levels and related global greenhouse gas emission pathways, in the context of*
736 *strengthening the global response to the threat of climate change, sustainable development,*
737 *and efforts to eradicate poverty*, eds. V. Masson-Delmotte, P. Zhai, H.-O. Pörtner, D. Roberts,
738 J. Skea, P.R. Shukla, A. Pirani, W. Moufouma-Okia, C. Péan, R. Pidcock, S. Connors, J.B.R.
739 Matthews, Y. Chen, X. Zhou, M.I. Gomis, E. Lonnoy, T. Maycock, M. Tignor & T. Waterfield
740 (Geneva, Switzerland: World Meteorological Society).

741 Intergovernmental Science-Policy on Biodiversity and Ecosystem Services (2019). *Global Assessment*
742 *Report on Biodiversity and Ecosystem Services*, eds. E.S. Brondizio, J. Settele, S. Díaz, HT
743 Ngo [Online]. Available: <https://ipbes.net/global-assessment>.

744 Karjalainen, E., Sarjala, T., and Raitio, H. (2010). Promoting human health through forests: overview
745 and major challenges. *Environ. Health Prev. Med.* 15, 1-8.

746 Keeton, W.S., Whitman, A.A., McGee, G.C., and Goodale, C.L. (2011). Late-successional biomass
747 development in northern hardwood-conifer forests of the Northeastern United States. *Forest*
748 *Sci.* 57, 489-505.

749 Kershner, B., and Leverett, R.T. (2004). *The Sierra Club Guide to the Ancient Forests of the*
750 *Northeast*. San Francisco: Sierra Club Books.

751 Law, B.E., Hudiburg, T.W., Berner, L.T., Kent, J.J., Buotte, P.C., and Harmon, M.E. (2018). Land use
752 strategies to mitigate climate change in carbon dense temperate forests. *Proc. Natl. Acad. Sci.*
753 *USA* 115, 3663-3668.

754 Leturcq, P. (2020). GHG displacement factors of harvested wood products: the myth of substitution.
755 *Sci. Rep.* 10, 20752.

756 Leverett, R.T., and Bertolette, D. (2014). *American Forests Champion Trees Measuring Guidelines*
757 *Handbook* [Online]. Available: [https://www.americanforests.org/wp-](https://www.americanforests.org/wp-content/uploads/2014/12/AF-Tree-Measuring-Guidelines_LR.pdf)
758 [content/uploads/2014/12/AF-Tree-Measuring-Guidelines_LR.pdf](https://www.americanforests.org/wp-content/uploads/2014/12/AF-Tree-Measuring-Guidelines_LR.pdf).

759 Leverett, R.T., Ruskin, D.N., and Masino, S.A. (2020). Direct measurement of trunk volume in forest
760 trees: Focus on white pine and comparison to a statistical method. *bioRxiv*,
761 doi.org/10.1101/2020.03.18.995985.

762 Lindenmayer, D.B., and Laurance, W.F. (2016). The unique challenges of conserving large old trees.
763 *Trends Ecol. Evol.* 31, 416-418.

764 Liu, X., Trogisch, S., He, J.-S., Niklaus, P.A., Bruelheide, H., Tang, Z., Erfmeier, A., Scherer-
765 Lorenzen, M., Pietsch, K.A., Yang, B., Kühn, P., Scholten, T., Huang, Y., Wang, C., Staab, M.,
766 Leppert, K.N., Wirth, C., Schmid, B., and Ma, K. (2018). Tree species richness increases
767 ecosystem carbon storage in subtropical forests. *Proc. Biol. Sci.* 285, 20181240.

768 Lutz, J.A., Furniss, T.J., Johnson, D.J., Davies, S.J., Allen, D., Alonso, A., Anderson-Teixeira, K.J.,
769 Andrade, A., Baltzer, J., Becker, K.M.L., Blomdahl, E.M., Bourg, N.A., Bunyavejchewin, S.,
770 Burslem, D.F.R.P., Cansler, C.A., Cao, K., Cao, M., Cárdenas, D., Chang, L.-W., Chao, K.-J.,
771 Chao, W.-C., Chiang, J.-M., Chu, C., Chuyong, G.B., Clay, K., Condit, R., Cordell, S.,
772 Dattaraja, H.S., Duque, A., Ewango, C.E.N., Fischer, G.A., Fletcher, C., Freund, J.A., Giardina,
773 C., Germain, S.J., Gilbert, G.S., Hao, Z., Hart, T., Hau, B.C.H., He, F., Hector, A., Howe,
774 R.W., Hsieh, C.-F., Hu, Y.-H., Hubbell, S.P., Inman-Narahari, F.M., Itoh, A., Janík, D.,
775 Kassim, A.R., Kenfack, D., Korte, L., Král, K., Larson, A.J., Li, Y.D., Lin, Y., Liu, S., Lum, S.,
776 Ma, K., Makana, J.-R., Malhi, Y., McMahon, S.M., Mcshea, W.J., Memiaghe, H.R., Mi, X.,
777 Morecroft, M., Musili, P.M., Myers, J.A., Novotny, V., De Oliveira, A., Ong, P., Orwig, D.A.,
778 Ostertag, R., Parker, G.G., Patankar, R., Phillips, R.P., Reynolds, G., Sack, L., Song, G.-Z.M.,
779 Su, S.-H., Sukumar, R., Sun, I.-F., Suresh, H.S., Swanson, M.E., Tan, S., Thomas, D.W.,
780 Thompson, J., Uriarte, M., Valencia, R., Vicentini, A., Vrška, T., Wang, X., Weiblen, G.D.,

781 Wolf, A., Wu, S.-H., Xu, H., Yamakura, T., Yap, S., and Zimmerman, J.K. (2018). Global
782 importance of large-diameter trees. *Glob. Ecol. Biogeogr.* 27, 849-864.

783 Mackey, B., Dellasala, D.A., Kormos, C., Lindenmayer, D., Kumpel, N., Zimmerman, B., Hugh, S.,
784 Young, V., Foley, S., Arsenis, K., and J.E.M., W. (2015). Policy options for the world's
785 primary forests in multilateral environmental agreements. *Conserv. Lett.* 8, 139-147.

786 McMullin, R.T., and Wiersma, Y.F. (2019). Out with OLD growth, in with ecological continNEWity:
787 new perspectives on forest conservation. *Front. Ecol. Environ.* 17, 176-181.

788 Mildrexler, D.J., Berner, L.T., Law, B.E., Birdsey, R.A., and Moomaw, W.R. (2020). Large trees
789 dominate carbon storage in forests east of the Cascade crest in the U.S. Pacific Northwest.
790 *Front. For. Glob. Change* 3, 594274.

791 Mitchard, E.T.A. (2018). The tropical forest carbon cycle and climate change. *Nature* 559, 527-534.

792 Moomaw, W.R., Masino, S.A., and Faison, E.K. (2019). Intact forests in the United States:
793 Proforestation mitigates climate change and serves the greatest good. *Front. Forests Global*
794 *Change* 2, 27.

795 Moose, R.A., Schigel, D., Kirby, L.J., and Shumskaya, M. (2019). Dead wood fungi in North America:
796 an insight into research and conservation potential. *Nat. Conserv.* 32, 1-17.

797 National Council for Air Stream Improvement (2020). United States Department of Agriculture Forest
798 Service. COLE: Carbon on Line Estimator.

799 Native Tree Society (2021). Available: <https://www.nativetreesociety.org/>, <https://www.ents-bbs.org>.

800 Nature Needs Half (2021). Available: <https://natureneedshalf.org/>.

801 Nicodemus, M.A., and Williams, R.A. (2004). "Quantifying aboveground carbon storage in managed
802 forest ecosystems in Ohio", in: *Proceedings of the 14th Central Hardwoods Forest Conference*
803 *GTR-NE-316*.

804 Nunery, J.S., and Keeton, W.S. (2010). Forest carbon storage in the northeastern United States: Net
805 effects of harvesting frequency, post-harvest retention, and wood products. *Forest Ecol.*
806 *Manag.* 259, 1363-1375.

807 Pugh, T.a.M., Lindeskog, M., Smith, B., Poulter, B., Arneth, A., Haverd, V., and Calle, L. (2019). Role
808 of forest regrowth in global carbon sink dynamics. *Proc. Natl. Acad. Sci. USA* 116, 4382-4387.

809 Riitters, K., Potter, K., Iannone, B., Oswalt, C., Guo, Q., and Fei, S. (2018). Exposure of protected and
810 unprotected forest to plant invasions in the eastern United States. *Forests* 9, 723.

811 Ripple, W.J., Wolf, C., Newsome, T.M., Barnard, P., and Moomaw, W.R. (2020). World scientists'
812 warning of a climate emergency. *BioScience* 70, 8-12.

813 Ruddat, J. (2020). *Documenting and Protecting New England's Old-Growth Forests* [Online].
814 Available: <https://digitalcommons.wpi.edu/iqp-all/5763>.

815 Schulz, F., Alteio, L., Goudeau, D., Ryan, E.M., Yu, F.B., Malmstrom, R.R., Blanchard, J., and
816 Woyke, T. (2018). Hidden diversity of soil giant viruses. *Nat. Commun.* 9, 4881.

817 Seymour, R. (2011). Influence of age on productivity and carbon sequestration of Eastern white pine.
818 *Northeastern States Research Cooperative*. Available: [https://nsrforest.org/project/influence-](https://nsrforest.org/project/influence-age-productivity-and-carbon-sequestration-eastern-white-pine)
819 [age-productivity-and-carbon-sequestration-eastern-white-pine](https://nsrforest.org/project/influence-age-productivity-and-carbon-sequestration-eastern-white-pine)

820 Seymour, R. (2016). Managing an aging resource: Influence of age on leaf area index, stemwood
821 growth, growth efficiency, and carbon sequestration of eastern white pine. *Northeastern States*
822 *Research Cooperative*. Available:
823 <https://nsrforest.org/sites/default/files/uploads/seymour11full.pdf>.

824 Simmons, C.T., and Matthews, H.D. (2016). Assessing the implications of human land-use change for
825 the transient climate response to cumulative carbon emissions. *Environ. Res. Lett.* 11, 035001.

826 Sperduto, D.D., Nichols, W.F., Crowley, K.F., and Bechtel, D.A. (2000). Black Gum (*Nyssa sylvatica*
827 Marsh) in New Hampshire. *New Hampshire Natural Heritage Inventory*, Concord, New
828 Hampshire.

829 Stephenson, N.L., Das, A.J., Condit, R., Russo, S.E., Baker, P.J., Beckman, N.G., Coomes, D.A.,
830 Lines, E.R., Morris, W.K., Rüger, N., Álvarez, E., Blundo, C., Bunyavejchewin, S., Chuyong,
831 G., Davies, S.J., Duque, Á., Ewango, C.N., Flores, O., Franklin, J.F., Grau, H.R., Hao, Z.,

832 Harmon, M.E., Hubbell, S.P., Kenfack, D., Lin, Y., Makana, J.-R., Malizia, A., Malizia, L.R.,
833 Pabst, R.J., Pongpattananurak, N., Su, S.-H., Sun, I.-F., Tan, S., Thomas, D., Van Mantgem,
834 P.J., Wang, X., Wiser, S.K., and Zavala, M.A. (2014). Rate of tree carbon accumulation
835 increases continuously with tree size. *Nature* 507, 90-93.

836 Sterman, J.D., Siegel, L., and Rooney-Varga, J.N. (2018). Reply to comment on 'Does replacing coal
837 with wood lower CO₂ emissions? Dynamic lifecycle analysis of wood bioenergy'. *Environ. Res.*
838 *Lett.* 13, 128003.

839 Thom, D., Golivets, M., Edling, L., Meigs, G.W., Gourevitch, J.D., Sonter, L.J., Galford, G.L., and
840 Keeton, W.S. (2019). The climate sensitivity of carbon, timber, and species richness covaries
841 with forest age in boreal-temperate North America. *Glob. Chang. Biol.* 25, 2446-2458.

842 Tomasso, L.P., and Leighton, M. (2014). The impact of land use change for greenhouse gas inventories
843 and state-level climate mediation policy: A GIS methodology applied to Connecticut. *J.*
844 *Environ. Protec.* 5, 52176.

845 United States Climate Alliance (2021). Available: <http://www.usclimatealliance.org/>.

846 United States Department of Agriculture (2009). *Specific Gravity and Other Properties of Wood and*
847 *Bark for 156 Tree Species Found in North America* [Online]. Available:
848 <https://www.fs.usda.gov/treearch/pubs/34185>.

849 United States Environmental Protection Agency (2018). *Sources of greenhouse gas emissions*
850 [Online]. Available: <https://www.epa.gov/ghgemissions/sources-greenhouse-gas-emissions>.

851 United States Forest Service (2019). *EVALIDator Version 1.8.0.01* [Online]. Available:
852 <https://apps.fs.usda.gov/Evalidator/evaluator.jsp>.

853 United States Forest Service (2020). *Forest Inventory Analysis* [Online]. Available:
854 <https://www.fia.fs.fed.us/>.

855 United States Forest Service (2021). *Ecosystem Services* [Online]. Available:
856 <https://www.fs.fed.us/ecosystemservices/>.

857 Watson, J.E.M., Evans, T., Venter, O., Williams, B., Tulloch, A., Stewart, C., Thompson, I., Ray, J.C.,
858 Murray, K., Salazar, A., Mcalpine, C., Potapov, P., Walston, J., Robinson, J.G., Painter, M.,
859 Wilkie, D., Filardi, C., Laurance, W.F., Houghton, R.A., Maxwell, S., Grantham, H., Samper,
860 C., Wang, S., Laestadius, L., R Runting, R.K., Silva-Chávez, G.A., Ervin, J., and Lindenmayer,
861 D. (2018). The exceptional value of intact forest ecosystems. *Nat. Ecol. Evol.* 2, 599-610.

862 Whitney, G.G. (1996). *From Coastal Wilderness to Fruited Plain: A History of Environmental Change*
863 *in Temperate North America from 1500 to the Present*. Cambridge University Press.

864 Wilson, E.O. (2016). *Half-Earth: Our Planet's Fight for Life*. New York, New York: Liveright
865 Publishing Corp.

866 World Wildlife Federation (2020). *Living Planet Report 2020* [Online]. Available:
867 <https://livingplanet.panda.org/en-us/>.

868 Zlonis, E., and Niemi, G. (2014). Avian communities of managed and wilderness hemiboreal forests.
869 *Forest Ecol. Manag.* 328, 26-34.

870

Figure 1.TIF




Figure 2.TIF




Figure 3.TIF

