

Data Submitted (UTC 11): 8/24/2022 10:19:04 PM

First name: Debbie

Last name: Kelly

Organization:

Title:

Comments: The piñon pine and juniper forests found in the red rock area are subject to temperature extremes and limited moisture availability. These forests are often the sole woodland provider of wildlife habitat, vegetative cover, watershed protection, and traditional food and medicine gathering in dryland and arid BLM-managed lands across the West.

Single-leaf piñon pine trees can reach ages of up to 600 years and juniper can reach ages of up to 1,600 years. These historic forests and associated undisturbed biological soil crusts store a disproportionate amount of carbon in dryland ecosystems, and are more resistant to disruptions caused by climate change.

Old-growth piñon-juniper forests are home to more than 70 bird species and are often the only suitable habitat for many of these species, whose populations are currently declining rapidly under intense pressure from climate change, development, and drought.

Although they cover hundreds of millions of acres and provide irreplaceable habitat and ecosystem benefits, piñon-juniper forests are among the least studied and most ignored North American forest types. The BLM should ensure that its inventory and criteria for mature and old-growth forests include piñon pine and juniper forests, which are the largest forest type by acreage managed by the agency.

Please discontinue the mechanical destruction of these forests and keep them intact for wildlife, climate change and future human liveability.