

Data Submitted (UTC 11): 3/30/2024 3:39:35 PM

First name: Larry

Last name: Evans

Organization:

Title:

Comments: Dear Planning Team,

Thank you for your work in updating and revising the forest plan in accordance with best scientific understandings.

I have noticed that there is no mention of fungi among the species of special concern. Referring to USDA PNW-GTR-371, Managing Forest Ecosystems to Conserve Fungus Diversity and Sustain Wild Mushroom Harvests, May 1996, and to Handbook to Additional Fungal Species of Special Concern in the Northwest Forest Plan, PNW-GTR-572, January 2003, I would recommend as a minimum these 20 fungal spp of special concern. All of these are endemic to Region 1, several are species complexes containing yet undescribed members. These species are largely mycorrhizal and all are dependent upon the mixed coniferous forests of the northern Rockies. They all sequester carbon, nitrogen, and water in their life processes and are a primary source of amino acids for the rest of the ecosystem.

*Russula brevipes*, a keystone species in forest ecosystems around the Pacific northwest and the Rocky mountains, boasts more interconnecting relationships than any other, being not only a mycorrhizal associate of most conifers but also several species of heterotrophs, including plant species of special concern, and hosts the ascomycete species *Hypomyces lactifluum*, which is the commercially traded lobster mushroom.

2. *Herichium abetinus*, the conifer-loving species of lions mane or bears head mushroom, is rare to infrequent throughout our region. It's European counterpart was redlisted and collection of wild *Herichium* in England for example is punishable by jail time. Related species are being cultivated commercially after discovery of medicinal neurogenic compounds produced by these fungi.

3. *Geopyxis carbonicola* group, a key element in every northern boreal ecosystem, the fire cup forms mycorrhizae with virtually every plant in every conifer forest. In Alaska, I saw that the limits of the forest exactly corresponded to the boundaries of the presence of *Geopyxis*, which fruits after a forest fire.

4. *Rhizopogon* spp. are hypogeous truffles related to *Suillus* spp. that form subsurface fruiting bodies which are energy and protein rich, drought-resistant, and mycorrhizal with lodgepole, juniper, Doug fir, and Ponderosa pine. These are a key protein source for elk, squirrels, and gestating deer, Trappe et.al. reported deer scat collections during August were found to contain primarily undigested *rhizopogon* spores.

5. *Lactarius deliciosus* group are mycorrhizal with conifers and deciduous shrubs, and while they are food for invertebrates, these highly nutritious mushrooms are often ignored by forest mammals here. In Europe the corresponding species, which I have sampled, is much sought after for its vitamin A and C rich mushrooms. Recent studies show that agroforestry plantations of oak and related *Lactarius indigo* produce more protein more efficiently than conventional farming.

6. *Cantharellus* spp. aka chanterelles are a group of highly prized commercially valuable mushrooms that are mycorrhizal with fir and spruce and cannot be cultivated. Rich in proteins and vitamins, they are indigestible raw, and largely ignored by wildlife. Their presence indicates primarily undisturbed habitat, and is a good indicator of forest health. The mycelium are easily killed by soil compaction from heavy equipment.

7. *Morchella* sp are native highly prized commercial species. So called fire morels appear after a forest fire and annually attract pickers from around the continent. Individual fires have yielded morel harvests valued in the millions of dollars, much of which are sold to European and Asian markets.

8. *Suillus* spp are a keystone species complex and primary source of amino acids in our ecosystems, these mycorrhizal species are key to the survival of all pine seedlings and form myriad relationships with other plants, fungi, and heterotrophs. *Suillus* mushrooms are the ubiquitous spongy slippery jacks, and are eaten by humans, ursine and ungulate mammals, and invertebrates.

9. *Cortinarius caperata*, aka Rozites, one of many members of this little studied family of mycorrhizal fungi, has

shown antiviral properties. (Spatafora, et. al.) one of few Cortinarius eaten by people, but wildlife consume a wide variety and enormous volume of these mushrooms.

10. Ramaria spp., the coral mushroom group, mycorrhizal with fir and pine, spring food for elk, a key protein source for wildlife in the spring. Needs full canopy and BCR rich forest soils, hyphae easily damaged by compaction. Some spp eaten by humans.

11. Hygrophorus spp, the waxy caps, are an important source of food for wildlife. they form mycorrhizae with all conifers, as well as deciduous shrubs, and even grasses. These high energy mushrooms are important for bears emerging from hibernation and overwintering deer and elk.

12. Fomitopsis pinicola, the red belt conk, is essential for soil building in our region as it creates brown cuboidal rot, which is the primary vehicle for water retention in our western forests. A traditional medicine, the conks can also be felted into a leathery material for hats and handicrafts. Previously dismissed as common, it's infrequency mirrors a decline in forest health.

13. Fomitopsis officinalis, aka agarikon, aka quinine conk, has been highly valued for its medicinal value since the days of the second world war, when the government purchased conks shot off the sides of trees with .22s by locals for use as anti malarial drugs. Paul Stamets recently published evidence of its efficacy against several viruses, from smallpox to influenza. It is rare and endemic to mature Doug fir, larch, and fir.

14. Boletus edulis group are mycorrhizal with conifers in our area and reliably produce volumes of delicious edible mushrooms. Their presence is also an indicator of a healthy forest, and often associated with

15. Amanita muscaria group often found in association with B. edulis, these toxic and medicinal mushrooms have a long history in many cultures. An integral part of the ecosystem, they form mycorrhizae with conifers and are food for mammals, birds, and invertebrates.

16. Tricholoma murrilli, aka matsutake, aka pine mushroom, enjoys enormous popularity in Asia, where it is widely considered a delicacy. Price surges in the past made matsutake hunting a lucrative lifestyle, now somewhat diminished by alternative sources. Still a commercially important mushroom and export, mycorrhizal with pine and fir in harsh environments.

17 Sparassis radicata aka cauliflower mushroom, a highly prized edible, indicative of old growth habitat, easily destroyed habitat disturbances, increasingly rare.

18 Pycnoporus sp. local to the Lolo National Forest have demonstrated superior lignase activity, producing a bright, high tensile pulp that matches or exceeds that produced by conventional chemical processes. It is now being used by pulp mills in Argentina. Information about bio pulping can be found at science direct/pycnoporus. Also used to break fever in traditional medicine.

19 Truffles, including Tuber spp. are critical mycorrhizal partners and a vital component of the diets of small mammals in the Northern Rockies. Rosentreader et. al (blm biologist) reported that 85% the protein needs of flying squirrels were sourced from fungi. Some species are of commercial interest. Truffle populations are known to be impacted, and the species mix changed, by silvicultural practices. The removal of the host tree, high soil temperatures, and soil compaction will heavily impact production of truffles (p136)

pnw gtr 772, april 2009

Our organization has recorded more than 40 species of hypogeous fungi during Western Montana Mycological Association events and forays since 1992. One such species list was previously submitted.

Regards,

Larry Evans